



Northern Research Station | General Technical Report NRS-203 | July 2021

CLIMATE ADAPTATION ACTIONS FOR URBAN FORESTS AND HUMAN HEALTH



Abstract

Urban areas can be particularly vulnerable to climate change due to extensive impervious cover, increased pollution, greater human population densities, and a concentration of built structures that intensify impacts from urban heat, drought, and extreme weather. Urban residents are at risk from a variety of climate stressors, which can cause both physical and mental harm. Urban forests and tree cover provide a critical role in helping cities address climate change by supporting greenhouse gas mitigation, reducing the impacts of extreme heat and altered climate that impair human health, and helping communities to adaptively respond through engagement with nature. At the same time, urban forests are vulnerable to changes in climate and in need of robust strategies to adapt to those changes.

As climate change impacts increase, efforts to “green” cities and adapt urban forests to changing conditions take on greater importance to support human health and well-being. Urban forest managers and allied professionals are looking for information to reduce climate risks to urban forests and secure their benefits for people and ecosystems. This report, *Climate Adaptation Actions for Urban Forests and Human Health*, synthesizes adaptation actions to address climate change in urban forest management and promote human health and well-being through nature-based solutions. It compiles and organizes information from a wide range of peer-reviewed research and evidence-based reports on climate change adaptation, urban forest management, carbon sequestration and storage, and human health response to urban nature.

This report includes the Urban Forest Climate and Health Adaptation Menu, which presents information and ideas for optimizing the climate and human health outcomes of urban forestry projects and provides professionals who are working at the intersection of climate, public health, and urban forestry with resources to support climate adaptation planning and activities. Notably, it *does not* provide specific recommendations or guidance for any particular place; rather, it offers a range of action opportunities at different scales that can be incorporated into either comprehensive or specific climate adaptation initiatives. The Menu can be used with an existing, tested adaptation process to help managers consider climate risks and explore the benefits and drawbacks of potential adaptation actions within the context of a particular situation or project. It also can be useful for generating productive discussions about community needs and values to guide planning, education and outreach, research, or changes in policy or infrastructure within communities.

This publication was previously released as a postprint after undergoing technical and policy review. This final version includes a completed layout, changes to wording and grammar, and limited corrections to content. The most current version of the publication will always be available from the Digital Object Identifier (DOI): <https://doi.org/10.2737/NRS-GTR-203>.

The use of trade, firm, corporation, or program names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the U.S. Department of Agriculture or the Forest Service of any product or service to the exclusion of others that may be suitable.

Manuscript received for publication 26 April 2021

Published by:

USDA FOREST SERVICE

ONE GIFFORD PINCHOT DRIVE

MADISON, WI 53726

Visit our homepage at <https://nrs.fs.fed.us>

July 2021

CLIMATE ADAPTATION ACTIONS FOR URBAN FORESTS AND HUMAN HEALTH

Maria K. Janowiak, Leslie A. Brandt, Kathleen L. Wolf, Mattison Brady, Lindsay Darling, Abigail Derby Lewis, Robert T. Fahey, Kristen Giesting, Eboni Hall, Molly Henry, Maisie Hughes, Jason W. Miesbauer, Kailey Marcinkowski, Todd Ontl, Annamarie Rutledge, Lydia Scott, and Christopher W. Swanston

About the Authors

Maria K. Janowiak is the deputy director of the Northern Institute of Applied Climate Science with the USDA Forest Service, Northern Research Station, 410 MacInnes Drive, Houghton, MI 49931, maria.janowiak@usda.gov.

Leslie A. Brandt is a climate change specialist with the Northern Institute of Applied Climate Science, USDA Forest Service, Eastern Region, 1992 Folwell Avenue, St. Paul, MN 55108, leslie.brandt@usda.gov.

Kathleen L. Wolf is a research social scientist with the College of the Environment, School of Environmental and Forest Sciences, University of Washington, Box 352100, Seattle WA 98195, kwolf@uw.edu.

Mattison Brady is a student climate change adaptation specialist with the Northern Institute of Applied Climate Science and Michigan Technological University, 1400 Townsend Drive, Houghton, MI 49931, mbrady1@mtu.edu.

Lindsay Darling is a geographic information systems administrator with the Morton Arboretum, 4100 Illinois Route 53, Lisle, IL 60532, ldarling@mortonarb.org.

Abigail Derby Lewis is director of the conservation tools program in the Keller Science Action Center at the Field Museum of Natural History, 1400 S. Lake Shore Drive, Chicago, IL 60605, aderby@fieldmuseum.org.

Robert T. Fahey is an assistant professor at the University of Connecticut, Department of Natural Resources and the Environment, and the Center for Environmental Sciences and Engineering, 1376 Storrs Road, Storrs, CT 06269, robert.fahey@uconn.edu.

Kristen Giesting is a soil conservationist with the Natural Resources Conservation Service, 1333 East Liberty Circle, Greensburg, IN 47240, kristen.giesting@usda.gov.

Eboni Hall is the senior manager of urban forestry education at American Forests, 1220 L St NW, Suite 750, Washington, D.C. 20005, ehall@americanforests.org.

Molly Henry is the senior manager of climate and health at American Forests, embedded in Division of Forest Environment at the Rhode Island Department of Environmental Management, 235 Promenade St, Suite 394, Providence, RI 02908, mhenry@americanforests.org.

Maisie Hughes is the senior director of urban forestry at American Forests, 1220 L St NW, Suite 750, Washington, D.C. 20005, mhughes@americanforests.org.

Jason W. Miesbauer is an arboriculture scientist with the Morton Arboretum, 4100 Illinois Route 53, Lisle, IL 60532, jmiesbauer@mortonarb.org.

Kailey Marcinkowski is a climate change education specialist with the Northern Institute of Applied Climate Science and Michigan Technological University, 1400 Townsend Drive, Houghton, MI, 49931, kfmarcin@mtu.edu.

Todd Ontl is a climate change adaptation specialist with the Northern Institute of Applied Climate Science and Michigan Technological University, 1400 Townsend Drive, Houghton, MI, 49931, taontl@mtu.edu.

Annamarie Rutledge is a climate change outreach specialist with the Northern Institute of Applied Climate Science and Michigan Technological University, 1400 Townsend Drive, Houghton, MI 49931, amrutled@mtu.edu.

Lydia Scott is director of the Chicago Region Trees Initiative, located at the Morton Arboretum, 4100 Illinois Route 53, Lisle, IL 60532, lscott@mortonarb.org.

Christopher W. Swanston is a research ecologist with the USDA Forest Service, Northern Research Station, the director of the Northern Institute of Applied Climate Science, and the director of the USDA Northern Forests Climate Hub, 410 MacInnes Drive, Houghton, MI 49931, christopher.swanston@usda.gov

Acknowledgments

Many individuals contributed to the guidance, review, and editing of this document. We would first like to extend our thanks to the authors of the first forest and urban forest menus upon which this work is based: Angela (Kerber) Levernier of Dupage County, Illinois; and Stephen Handler, Patricia Butler-Leopold, and Danielle Shannon of the Northern Institute of Applied Climate Science. We express gratitude to Sarah Hines, Sarah Hurteau, and Natalie van Doorn for the insightful and productive feedback in reviewing this document. We also are grateful to the hundreds of managers who have demonstrated the practical application of earlier iterations of our adaptation products, as well as the Rhode Island Urban Forests for Climate and Health Stakeholder Group, who helped us formulate this new menu of actions. This work would not be possible without their interest and feedback.

The Doris Duke Charitable Foundation provided generous support to this project as part of a broader investment to optimize urban forests for climate and public health outcomes.

Cover image created by Kailey Marcinkowski using USDA photo by Lance Cheung.

Contents

Introduction	1
What Is The Urban Forest Climate and Health Menu?	4
An Adaptation Planning Process	5
How to Read this Menu	7
Description of Menu Items	11
Strategy 1: Activate social systems for equitable climate adaptation, urban forest, and human health outcomes	11
Strategy 2: Reduce the impact of human health threats and stressors using urban trees and forests.	15
Strategy 3: Maintain or increase extent of urban forests and vegetative cover.....	20
Strategy 4: Sustain or restore fundamental ecological functions of urban ecosystems.....	25
Strategy 5: Reduce the impact of physical and biological stressors on urban forests	29
Strategy 6: Enhance taxonomic, functional, and structural diversity	38
Strategy 7. Alter urban ecosystems toward new and expected conditions.....	43
Strategy 8: Promote mental and social health in response to climate change.....	52
Strategy 9: Promote human health co-benefits in nature-based climate adaptation.....	55
Glossary	59
Literature Cited	65
Appendix 1.....	101
Appendix 2.....	103
Appendix 3.....	109

INTRODUCTION

Climate change is having a profound impact on cities across the world, and these changes are expected to accelerate in coming decades.^{1,2} Urban areas can be particularly vulnerable as extensive impervious cover and concentration of built structures intensify impacts from urban heat, drought, and extreme weather.^{3,4} In addition, human population densities in cities increase the risk of human illness and injury from climate influences. Urban forests, defined here as all publicly and privately owned trees within an urban area, provide many ecosystem services. These include helping reduce the impacts of climate change on people and human communities, including reducing the urban heat island effect and moderating stormwater runoff (Table 1). Urban forests are increasingly being recognized for their value in protecting and enhancing human health and safety generally, and particularly in the face of a changing climate.

Urban forests are an integral element of green infrastructure; that is, the “interconnected network of green space that conserves natural ecosystem values and functions and provides associated benefits to human populations.”⁵ Urban forests include built and cultural settings such as streetscapes and yards, parks, cemeteries, school grounds, corporate campuses, greenways, and unmanaged green spaces, as well as what are defined here as natural areas, such as patches of native forests, open woodlands, savannahs, and barrens. The vitality and interconnectedness of these urban forest elements are critical in supporting landscape-scale ecological processes (such as carbon sequestration, wildlife migration, and pollination) and long-term ecological functions of urban landscapes.⁶ Further, the urban forest contributes to the human habitats of cities, the places and conditions that offer necessary respite from busy lives.⁷ Trees and landscapes, if well planned and designed, can improve human health and quality of life in many ways.

Urban forests face unique challenges relative to forests outside of cities, such as increased pollution, restricted rooting conditions, and altered soils. These challenges affect the ability of cities to establish, maintain, and improve tree growth and forest canopy cover while simultaneously managing the maintenance costs, infrastructure damage, and nuisance complaints associated with urban trees.⁸ Climate change increases many stressors on urban forests and vegetation, with cascading effects on human and community health.

Urban forests and vegetation can be used proactively to address and alleviate many of the effects of climate change on cities and residents (Fig. 1). Urban forests can help reduce the severity of climate change by reducing energy use for heating and cooling and by sequestering carbon. They also can support urban climate adaptation goals, such as heat reduction and improved air quality, which have direct human health implications. Bringing nature closer to people to promote human health is another opportunity of urban forestry and climate response. Urban residents face challenges of crowding, interpersonal stressors, and safety concerns.¹⁰ Interactions with trees—from streetscapes to forested reserves—can counter these experiences and promote mental, social, and physical health benefits.¹¹

Table 1.—Examples of interactions among climate change, urban forests, and human health and the ways in which urban forests reduce climate impacts on people and communities

Climate Impact	Potential Impacts on Urban Forests	Potential Impacts on People and Communities	Ways to Reduce Impacts on People and Communities
Warmer temperatures and more days with extreme heat	<ul style="list-style-type: none"> ▶ Physiological stress on trees ▶ Range expansion of pests, disease, and invasive plant species in response to warmer conditions ▶ Potentially increased production of volatile organic compounds 	<ul style="list-style-type: none"> ▶ Heat stress leading to illness or mortality ▶ More low air quality days that exacerbate asthma, cardiovascular, and other illnesses ▶ Increased energy utility demand for cooling and peak use failure 	<ul style="list-style-type: none"> ▶ Reduce extreme heat by providing canopy cover, shade, and moisture ▶ Reduce energy use for cooling ▶ Improve air and water quality ▶ Become thermal refuges for urban dwellers
Fewer days with extreme cold	<ul style="list-style-type: none"> ▶ Shifts to milder hardiness zones and altered plant habitat suitability ▶ Reduced mortality of diseases and pests caused by extreme cold 	<ul style="list-style-type: none"> ▶ Reduced demand for energy to heat homes ▶ Potentially fewer deaths from extreme cold ▶ Increased survivability in disease-transmitting insects 	<ul style="list-style-type: none"> ▶ Strategically placed trees further reduce wind and improve passive solar, thus reducing energy use ▶ Improve conditions for physical activity
Altered precipitation creating wetter conditions in some seasons	<ul style="list-style-type: none"> ▶ Wet conditions may favor some plant pathogens 	<ul style="list-style-type: none"> ▶ Increased mold exposure leading to upper respiratory symptoms ▶ Shifts in water quality and quantity 	<ul style="list-style-type: none"> ▶ Enhance filtration of pollutants and improved water quality ▶ Canopy interception of precipitation and buffering rain cycles
More frequent heavy precipitation events	<ul style="list-style-type: none"> ▶ Increased stormwater runoff and localized flooding ▶ Mortality of trees in flood-prone areas ▶ Soil saturation and slope failures 	<ul style="list-style-type: none"> ▶ Disaster-related injury and death ▶ Reduced water quality ▶ Mold and property damage ▶ Disruption to food systems 	<ul style="list-style-type: none"> ▶ Reduce runoff from forest interception and absorption of rain ▶ Root systems prevent erosion and property damage
Elevated risk of drought or aridification	<ul style="list-style-type: none"> ▶ Mortality of drought-susceptible trees ▶ Reduced forest growth and health ▶ Increased stress on forests and shifts to non-forest vegetation 	<ul style="list-style-type: none"> ▶ Reduced water supply ▶ Disruption to food systems ▶ Increased fire risk ▶ Increased dust, smoke, and fine particulates in air 	<ul style="list-style-type: none"> ▶ Enhance moisture retention in landscapes helps buffer shifts in precipitation ▶ Improve water quality and storage by healthy soils ▶ Interception and filtering of particulates
Increases in carbon dioxide (CO₂), a greenhouse gas	<ul style="list-style-type: none"> ▶ Increased pollen production 	<ul style="list-style-type: none"> ▶ More allergens ▶ Increased respiratory illness and asthma ▶ Human physiology stressor⁹ 	<ul style="list-style-type: none"> ▶ Reduce impacts from selection of lower-allergen or female trees ▶ Reduce localized CO₂ levels

Urban forest managers and allied professionals, such as urban planners and public health officials, are seeking tools and information to help reduce climate risks to urban forests and promote their beneficial functions. Included in this report is the Urban Forest Climate and Health Adaptation Menu (hereafter referred to as “Menu”), which provides information and ideas for optimizing the climate and human health outcomes of urban forestry projects and gives professionals working at the intersection of climate, health, and urban forestry resources to support climate adaptation planning and activities. Multiple agencies and organizations, along with professionals across multiple disciplines, can collaborate to increase climate and human health benefits. Further, attention to environmental justice and equity is important in these processes. Forestry programs and activities can help address health disparities across underserved populations, as well as inequities in distribution of trees and green spaces across communities. Community engagement that respects local knowledge while incorporating community needs and traditions can support urban forestry projects that promote tree and human health.



Figure 1.—These are some of the benefits that forested urban ecosystems may provide for climate adaptation, carbon mitigation, and human health.

What Is The Urban Forest Climate and Health Menu?

One of the major challenges in responding to climate change is translating broad, science-based concepts for climate response into specific, tangible actions that can be implemented within a community or neighborhood. The scientific and technical literature is replete with conceptual frameworks,¹²⁻¹⁴ compiled adaptation strategies,¹⁵⁻¹⁷ and tools to support management decisions.¹⁸⁻²⁰ However, more specific information is needed to help natural resources managers and community leaders identify actions suitable for particular landscapes and projects. The Menu addresses this challenge by providing a synthesis of adaptation actions that are nature-based solutions to address climate change in urban forest management and improve human health and well-being.

The Menu compiles and organizes information from a wide range of peer-reviewed research and evidence-based reports on climate change adaptation, urban forest management, and human health response to urban nature. Actions to increase carbon sequestration and storage in urban forests also are included when possible (Box 1), as these activities can

Box 1: Trees and Climate Mitigation

Forests are increasingly recognized for their potential as “natural climate solutions” for land-based carbon mitigation, given the need to prevent the most severe impacts of climate change in the near term.^{1, 22, 23} Urban forests can support greenhouse gas reductions by reducing energy use,^{24, 25} sequestering and storing carbon within trees and soils,²⁶ and providing material for wood products (Fig. 2). Ecosystems that are adapted to changing and more variable conditions also can provide increasingly important benefits for carbon mitigation.²⁷ Actions to increase both adaptation and mitigation benefits can have synergistic effects regarding climate change²⁸ and human health benefits. When possible, the Menu discusses key benefits and tradeoffs for carbon mitigation.




Figure 2.—Urban forest carbon mitigation benefits. ²⁴⁻²⁶

reduce the overall impact of climate change on humans and urban ecosystems. This work draws heavily on research and practice from the temperate regions of North America; thus, some adaptation actions may not be appropriate in different climate zones (e.g., semi-desert or desert) or in situations where forest or tree cover is not desirable or feasible.

The Menu builds on previous resources for climate response, in particular, an urban forests menu that was published in “Forest Adaptation Resources: Climate Tools and Approaches for Land Managers.”²¹ That report and its resources were developed for use in the Upper Midwest and Northeast using information from a comprehensive literature review of adaptation actions at numerous scales and locations. Subsequent testing, review, and feedback from practitioners was used to refine this adaptation Menu, which has been used by urban forestry professionals to develop real-world adaptation projects (www.forestadaptation.org/demos). Our report incorporates new considerations of human health, carbon mitigation, and human dimensions that are integral to urban forest management and expands the scope to temperate regions within North America. The effort was led by the Northern Institute of Applied Climate Science, the USDA Forest Service, American Forests, and the University of Washington.

An Adaptation Planning Process

The Menu is designed to be used in conjunction with the Adaptation Workbook,²¹ a tool that provides a structured, adaptive approach for integrating climate change considerations




Figure 3.—The Adaptation Workbook²¹ describes an assessment and decision process that is used in conjunction with vulnerability assessments, local knowledge, and adaptation strategies menus. The results are site-specific actions that address explicit management and conservation objectives under a range of potential future climates. A brief description of the Adaptation Workbook process is presented in [appendix 1](#) and in the “Climate & Health Action Guide.”²⁹

into planning, decision-making, and implementation of urban forest resources (Fig. 3). The Adaptation Workbook is a step-by-step process that helps users consider the potential effects of climate change and design land management and conservation actions that can help prepare for changing conditions. It can accommodate a wide variety of geographic locations, ownership types, ecosystems and land uses, management goals, and project sizes.

Together, the Menu and the Adaptation Workbook help managers consider climate risks and explore the benefits and drawbacks of potential adaptation actions within the context of a particular situation or project. While the Workbook was developed to be applied in natural resource management, including urban forestry, the process also can be adopted for planning of human health systems. Further, the “Climate & Health Action Guide”²⁹ is available online as an entry point to the Adaptation Workbook.²¹

The Adaptation Workbook and growing list of resources have been used together in hundreds of real-world natural resources management projects. Other menus address resource areas such as agriculture,³⁰ forest carbon management,²⁷ recreation,³¹ forested watersheds,³² open

Box 2: Climate and Health Adaptation on a Neighborhood Scale

A team of urban forestry professionals and community partners in Rhode Island used the Urban Forest Climate and Health Adaptation Menu, the Adaptation Workbook, and other resources to evaluate climate change impacts and outline adaptation efforts for a real-world project. The Providence Parks Department and the Providence Neighborhood Planting Program are working to engage residents and neighborhood stakeholders in developing and implementing community-driven tree-planting and stewardship solutions focused on climate adaptation and human health in Upper and Lower South Providence.

Tree canopy cover in Upper and Lower South Providence is threatened by increased temperatures and precipitation, more frequent extreme weather events, and altered soil moisture. Neighborhoods in this region are disproportionately burdened by the impacts of climate change and environmental injustice.³⁵ A number of interrelated factors, such as a high coverage of impervious surfaces, low tree canopy cover, and proximity to a major highway and industrial port, result in negative impacts on health and well-being; these include urban heat island effect, flooding, and poor air quality. The local urban forest managers used the Menu to identify adaptation actions, including:

- ▶ Increase and improve tree canopy
- ▶ Select climate-adapted tree species
- ▶ Increase ground cover biodiversity
- ▶ Install curbside bioswale tree filter pits
- ▶ Assess and maintain existing tree stock

[Appendix 2](#) presents a case study of this adaptation project. [Appendix 3](#) includes a tree species list for Rhode Island that provides information on climate vulnerability, carbon benefits, and health services and disservices for more than 120 tree species.

wetlands,³³ and culturally relevant indigenous perspectives.³⁴ Many of these are presented online as adaptation demonstrations that provide relevant case studies of real-world adaptation projects. To test the relevance of this Menu, community forestry partners from Rhode Island used it and Adaptation Workbook together in the development of an adaptation demonstration for a South Providence neighborhood (Box 2; also, [appendix 2](#)).

How to Read this Menu

The Menu offers nature-based solutions that address climate change and promote human health and well-being in urban forest management and conservation. The Menu is organized hierarchically into strategies, approaches, and tactics (Fig 4, Box 3).

STRATEGY is defined as a broad adaptation response that is applicable across a variety of socio-ecological systems, natural resources and sites, hydrologic and ecological conditions, and overarching management goals.

APPROACH is a detailed adaptation method selected in response to a specific issue, site condition, or management objective that further describes how strategies could be employed.

TACTICS are prescriptive actions designed to be useful for local community or site conditions and management objectives. Tactics are the most specific adaptation response, being practical actions that can be tailored to unique situations and fit the needs of particular species, ecosystem type, site conditions, management objectives, and other factors. For communities and health, tactics span choices to engage in planning, reduce risk, protect vulnerable populations, and design for optimal vegetation placement. Examples of tactics are provided for each approach.

The strategies, approaches, and tactics are derived from a wide variety of peer-reviewed

CONCEPT

Strategy: A strategy is a broad adaptation response that is applicable across a variety of resources and sites

Example Strategy (8): Promote mental and social health in the face of climate change

Approach: An approach is an adaptation response that is more specific to a resource issue or geography

Example Approach (8.2): Encourage community and social cohesion to support climate change

Tactic: The most specific adaptation response, providing prescriptive direction about actions that can be applied to specific situations

Example Tactic: Create community gathering spaces and stewardship programs in green spaces

ACTION

Figure 4.—The hierarchical relationship of strategies, approaches, and tactics relevant to urban forestry, with an example of each. These serve as “stepping stones” for moving from broad concepts to actions that can be applied to a project, policy, or planning effort.

research and evidence-based reports. The Menu helps to translate these sources and their concepts into targeted and prescriptive tactics for responding to climate change in communities and offers opportunities at different scales that can be incorporated into either comprehensive or specific climate adaptation initiatives. Notably, it does not provide specific recommendations or guidance for any particular place or situation, and not all strategies and approaches will work together (Table 2). Like any menu, the Urban Forest Climate and Health Adaptation Menu presents options to the user; however, some options will appear more suitable and appropriate than others. Actions that work well in one setting or community type may not work in another.

The Menu items can be applied in various combinations to achieve desired outcomes and can build upon any current management actions that support long-term urban forest sustainability and resilience, as well as more livable human communities. In addition to actionable choices, the Menu can be useful for brainstorming adaptation actions and generating productive discussions about community needs and values. Menu choices can be used to guide planning, education and outreach, and research, as well as changes in policy or infrastructure within communities.

Table 2.—Considerations for using Urban Forest Climate and Health Adaptation Menu

Menu Provides:	Menu Does Not:
<ul style="list-style-type: none"> ▶ A broad spectrum of climate adaptation actions that can help sustain healthy ecosystems and resilient green infrastructure. ▶ A platform for discussing climate change-related challenges and adaptation methods for projects from site to citywide scale. ▶ An actions framework offering managers choices for new programs and actions that align with their specific management goals and objectives. ▶ Approaches to address climate-related human health threats and additional nature-based solutions for wellness, especially for disadvantaged communities. ▶ A framework that promotes collaborative discussions and partnerships among urban forestry, planning, public health, and other allied professionals. ▶ Examples of tactics that can be used to implement a strategy or approach, yet are flexible and adaptable to local conditions and programs. 	<ul style="list-style-type: none"> ▶ Make recommendations or set guidelines for management decisions. It is up to the manager and stakeholders to decide how this information is used. ▶ Express preference for any strategies or approaches within a particular community or place, as these will depend on site-specific factors, local knowledge, and management goals. ▶ Provide an exhaustive set of tactics. Managers and stakeholders are encouraged to consider additional actionable tactics appropriate for their projects. ▶ Apply equally to all regions. The Menu will be most applicable to temperate regions and areas capable of supporting trees. ▶ Cover all topics that pertain to urban forestry and human health. Additional menus related to recreation, wildlife management, and watershed management, among others, can be found at www.forestadaptation.org/strategies. A recent review¹¹ provides additional urban forestry and human health information.

BOX 3: Urban Forest Climate And Health Adaptation Menu

Strategy 1: Activate social systems for equitable climate adaptation, urban forest, and human health outcomes

Approach 1.1: Address socio-ecological systems in early, comprehensive response.

Approach 1.2: Integrate urban forestry in climate planning and policy.

Approach 1.3: Address climate and health challenges of disadvantaged communities and vulnerable populations.

Strategy 2: Reduce the impact of human health threats and stressors using urban trees and forests

Approach 2.1: Reduce extreme temperatures and heat exposure.

Approach 2.2: Improve urban air quality conditions.

Approach 2.3: Anticipate and reduce human health impacts of hazardous weather and disturbance events.

Strategy 3: Maintain or increase extent of urban forests and vegetative cover

Approach 3.1: Minimize forest loss and degradation.

Approach 3.2: Maintain existing trees through proper care and maintenance.

Approach 3.3: Restore and increase tree, forest, and vegetative cover.

Approach 3.4: Sustain sites and ecosystems that provide high value across the landscape.

Strategy 4: Sustain or restore fundamental ecological functions of urban ecosystems

Approach 4.1: Maintain or restore soils and nutrient cycling in urban areas.

Approach 4.2: Maintain or restore hydrologic processes in urban forests.

Approach 4.3: Restore or maintain fire in fire-adapted ecosystems.

Strategy 5: Reduce the impact of physical and biological stressors on urban forests

Approach 5.1: Reduce impacts from extreme rainfall and enhance water infiltration and storage.

Approach 5.2: Reduce risk of damage from extreme storms and wind.

Approach 5.3: Reduce risk of damage from wildfire.

Approach 5.4: Maintain or improve the ability of forests to resist pests and pathogens.

Approach 5.5: Prevent invasive plant establishment and remove existing invasive species.

Approach 5.6: Manage herbivory to promote regeneration, growth, and form of desired species.

Strategy 6: Enhance taxonomic, functional, and structural diversity

Approach 6.1: Enhance age class and structural diversity in forests.

Approach 6.2: Maintain or enhance diversity of native species.

Approach 6.3: Optimize and diversify tree species selection for multiple long-term benefits.

Approach 6.4: Maintain or enhance genetic diversity.

Strategy 7. Alter urban ecosystems toward new and expected conditions

Approach 7.1: Favor or restore non-invasive species that are expected to be adapted to future conditions.

Approach 7.2: Establish or encourage new species mixes.

Approach 7.3: Introduce species, genotypes, and cultivars that are expected to be adapted to future conditions.

Approach 7.4: Disfavor species that are distinctly maladapted.

Approach 7.5: Move at-risk species to more suitable locations.

Approach 7.6: Promptly revegetate and remediate sites after disturbance.

Approach 7.7: Realign severely altered systems toward future conditions.

Strategy 8: Promote mental and social health in response to climate change

Approach 8.1: Provide nature experiences to ease stress and support mental function.

Approach 8.2: Encourage community and social cohesion to support climate response.

Strategy 9: Promote human health co-benefits in nature-based climate adaptation

Approach 9.1: Co-design large-scale green infrastructure and built systems to promote health.

Approach 9.2: Provide micro-scale nature experiences to promote health and healing.

DESCRIPTION OF MENU ITEMS

Strategy 1: Activate social systems for equitable climate adaptation, urban forest, and human health outcomes

This strategy addresses how the goals and programs for urban forest ecological adaptation can fit into the broader policy- and decision-making processes of local governments and organizations. It encourages action and engagement at the broadest scale of governance and policy within jurisdictions and by agencies. Effectively addressing climate change is complex, from identifying causes to understanding effects. Although urban forests are not the dominant land use or land cover in most cities, urban trees can deliver practical solutions for climate mitigation and adaptation, while delivering important co-benefits—including human health.³⁶ This is especially important as most Americans now live in urbanized areas, a term that encompasses cities and towns of all sizes. The success of climate strategies depends on the dedicated activity of stakeholders, passionate champions, and innovations within social systems from the local to regional level. An action model has been used to promote sustainable urban forests and includes three general activities: understand the vegetation resource, engage urban forest stakeholder communities, and enact quality resource management.³⁷ A focus on the urban forest resource and its management is important, alongside equal commitment to community involvement and public outreach. Effectively engaging urban residents and leadership can help sustain urban forest, climate, and environmental justice goals for the health of both trees and humans.

Approach 1.1: Address socio-ecological systems in early, comprehensive response

Addressing both climate and health implications at the earliest stages of planning can be helpful in ensuring a cohesive and comprehensive response.^{38,39} Urban forest projects may range from a site-scale memorial tree planting to a tree-planting campaign across an entire city. Combined analysis using big data (such as remote sensing), on-the-ground measures, civic science inputs, and social media records can inform better understanding of urban situations and climate patterns from local to regional scales.⁴⁰ Planning and working effectively at all scales involves partnership and collaboration, engaging professional expertise, local knowledge, and civic leadership and pride embedded within communities.⁴¹ Organizations and institutions that once regarded trees as beyond their scope are now becoming more attuned to nature-based solutions.⁴² While complex and sometimes time consuming, a commitment to incorporating a socio-ecological outlook in urban forestry planning and management can lead to more trees planted and stewarded, broader cross-sectoral support, and dedicated resources.⁴³

EXAMPLE ADAPTATION TACTICS

- ▶ Engage the public, decision-makers, and thought leaders in discussions of shared visions about urban forest planning and management, particularly on topics of climate and human health.⁴⁴

- ▶ For all tree and forest projects, evaluate potential to address multiple climate-based human health impacts⁴⁵ using a co-benefits outlook.⁴⁶
- ▶ Collaborate with public health and other agencies that work on climate planning to promote the urban forest as an upstream public health solution that addresses root causes of health problems rather than symptoms.^{47, 48}
- ▶ Synchronize urban forest planning and management across jurisdictions and departments to address emergent climate-based community health risks and threats.⁴⁹
- ▶ Identify, assess, and acknowledge disparities in canopy, parks, and green space distribution⁵⁰ as part of tree canopy analysis.⁵¹
- ▶ Co-develop community-driven projects that increase nearby nature, including experiences of trees and forests, an evidence-supported social determinant of health, and a path to tree equity.^{52, 53}
- ▶ Involve and engage communities in development of tree projects that reflect community and cultural values, while addressing local climate impacts and associated health effects.^{54, 55}

Approach 1.2: Integrate urban forestry in climate planning and policy

Many government agencies promote green infrastructure strategies and nature-based solutions in climate planning and policy.^{56, 57} Including urban forestry in planning and policy development helps ensure that trees are integrated into long-term climate response and that urban forestry professionals are included in these decisions. Connecting trees to climate change mitigation can also lead to more sustained and consistent urban forestry support.⁵⁸ There are multiple dimensions of planning that can include urban forestry and address climate and health. Comprehensive and strategic plans that serve as blueprints for most city initiatives and governance can include climate goals. Additionally, explicit climate mitigation and adaptation plans can be developed at the level of local government, ideally also addressing disaster preparedness. Further, municipalities may incorporate provisions for climate and health into urban forest management plans, as they provide guidance for sustainable urban forests and promote best practices. Planning entails processes of developing, drafting, and stakeholder vetting of initiatives, with reliance on public education, outreach, and engagement.⁴⁹ The resulting public dialog about planning and policy fosters improved awareness of emergent concerns and solutions by the public and local leaders, which can extend to urban forestry.⁵⁹ Incorporating trees and urban forests in all plans that reference climate can help promote climate-resilient communities, protect all people from the health impacts of climate change, and optimize health and health equity outcomes.

EXAMPLE ADAPTATION TACTICS

- ▶ Within and across local jurisdictions, assess tree canopy and other green spaces to analyze and ascertain tree distribution as a baseline for planning.⁶⁰
- ▶ Analyze disparities of green infrastructure distribution, especially in relationship to patterns of health morbidity (meaning rates of diseases) and mortality.⁶¹⁻⁶³

- ▶ Include climate and health interactions, with attention to environmental justice and equity, within municipal or county urban forest management plans.⁶⁴⁻⁶⁶
- ▶ Conduct a Health Impact Assessment of any urban forestry program or management plan on the community to local jurisdiction scale.⁶⁷
- ▶ Prepare Urban Forest Emergency Management Plans or Storm Mitigation Plans at the local government level to anticipate and respond to extreme weather and potential hazards from trees.⁶⁸
- ▶ Collaborate across city departments and agencies—including parks, transportation, public works, utilities, and school districts—to include trees, open space, or other components of green infrastructure in comprehensive climate action planning.^{69, 70}
- ▶ Facilitate the inclusion of trees and urban forests in all foundational urban planning (such as comprehensive plans and strategic plans) and associated departmental or sector-specific plans (such as tree retention in development, permitting policy, and transportation).⁵⁴
- ▶ Incorporate trees, natural areas, parks, and the urban forest in local government capital planning and budgeting that is dedicated to climate response and resilience.⁷¹
- ▶ Include urban forest, climate, and health interactions in more localized planning, such as Tribal or community plans,⁷² and also in implementation, such as code and ordinances.
- ▶ Include green infrastructure and urban forest parameters when developing metrics and benchmarking performance of climate policies or programs and addressing health outcomes.^{49, 73}



Forested urban paths and trails promote physical activity, a pathway to cardiovascular and respiratory health. Courtesy photo by Guy Kramer, used with permission.

- ▶ Expand planning and policy from single communities to multi-jurisdictional regional collaborations, supporting expansion of climate adaptation functions and services to the ecosystem scale.^{74, 75}

Approach 1.3: Address climate and health challenges of disadvantaged communities and vulnerable populations

Certain communities and populations are disproportionately at risk of impacts from climate change. Such risks are due to both historic legacies of discrimination (such as redlining^{76, 77}) and ongoing social, economic, and health challenges. Health concerns touch both place and person. Disadvantaged communities are often situated physically and economically such that they may experience “first and worst” climate change consequences. Native and Tribal communities are disproportionately affected by these consequences, as are residents living in lower-income or marginalized neighborhoods. Insufficient economic funding and infrastructure to respond to these threats can jeopardize these vulnerable communities.⁷⁸ In addition, there are population groups of specific demographics (such as elder or young age, or those with pre-existing health conditions) that are generally more vulnerable to changing conditions and environmental health threats.⁷⁹ Some climate conditions affect both disadvantaged communities and vulnerable populations disproportionately,⁸⁰ so attention to both human and tree health impacts is important. Equity is becoming a guiding principle in urban forestry, as there are disparities in the distribution of tree canopy in many cities.^{62, 81} Lower-income communities often have less tree canopy cover than nearby communities of greater affluence.^{82, 83} Successfully addressing these inequities is more complex than simply planting trees, however; careful consideration of potential positive and negative consequences for residents across the project location, including people having particular health sensitivities, may lead to more nuanced planning and implementation of tree planting.⁸⁴ A carefully planned urban forest project can generate a wide range of health benefits; however, urban greening, such as tree planting, also can be associated with dynamics of gentrification that can result in displacement of long-established residents.⁸⁵⁻⁸⁷ Careful, responsible urban forestry and urban greening planning includes important goals of community engagement, as well as involvement and collaboration with those who know their communities best.

EXAMPLE ADAPTATION TACTICS

Work to understand local climate-based human health vulnerabilities, possibly including a formal vulnerability analysis to define, identify, classify, and prioritize response to risk.^{61, 88}

- ▶ Apply the principles and practices of Community Based Climate Adaptation,^{89, 90} an inclusive approach that engages residents of disadvantaged communities early in projects and integrates local knowledge in the design and implementation of climate actions.
- ▶ Identify and prioritize sites for urban forest projects where vulnerable populations are concentrated, such as schools, elder care facilities, or health treatment centers.^{91, 92}

- ▶ Plan and select tree projects that can initiate and anchor the creation of public and civic nature spaces, and in so doing, address known disparities of parks and green space distribution and access.^{65, 93}
- ▶ Conduct outreach to residents of disadvantaged communities to understand their experience with urban forestry programs, determine needs, and initiate collaborations on improvements.^{53, 94}
- ▶ Engage residents and community organizations in planning and stewardship of tree projects so that benefits are sustained.⁹⁵
- ▶ Collaborate with community members and organizations to evaluate and implement culturally significant, relevant, and symbolic choices for green space vegetation, including trees and other vegetation.⁹⁶⁻⁹⁹
- ▶ Address potential gentrification or displacement consequences of urban forest enhancements within marginalized communities.^{85, 100-104}
- ▶ Involve local businesses and hire community residents to develop green infrastructure jobs and career pathways that are place-based and support beneficial climate and health outcomes.¹⁰⁵⁻¹¹⁰

Strategy 2: Reduce the impact of human health threats and stressors using urban trees and forests

This strategy describes how the urban forest can address multiple direct climate-related impacts in communities. Climate change is altering fundamental processes and conditions that sustain the lives and health of local trees and people. Communities are feeling the resulting consequences; there also are multiple public health implications for people within most cities.¹¹¹ Rising temperatures lead to more frequent heatwaves, with impacts amplified in urban settings, leading to heat-related illness and mortality.¹¹² Heat and other climate influences combine to impact air quality, while increased levels of particulates and allergens contribute to cardiovascular and respiratory illnesses.¹¹³ Extreme weather sets up conditions for increased flooding and storm events, along with more frequent and intense wildfires. Sea level rise introduces risks for coastal properties and residents. These disruptive events and changes can directly impact basic life support, such as drinking water contamination or loss of homes, and can introduce indirect health effects, such as psychological trauma and grieving.¹¹⁴ Nature-based adaptation strategies can be used to both prevent and respond to these interrelated human and ecosystem health threats. Strategically planned and implemented urban tree, forest, and greening projects present important opportunities to mitigate and accelerate recovery from extreme weather and disturbance events.¹¹⁵

Approach 2.1: Reduce extreme temperatures and heat exposure

One of the primary public health concerns regarding climate change is the increased intensity and frequency of heat waves. For instance, the number of days hotter than 100 °F in cities may increase nearly threefold in the United States by 2050, and the duration of the longest extreme heat events in an average year will double.¹¹² Extreme heat events cause

more deaths in the United States annually than all other weather-related causes combined (including hurricanes, lightning, tornadoes, and floods), and the effects of these events are most pronounced in urban areas.¹¹⁶ Increased heat is associated with higher rates of both heat stroke and hyperthermia; mortality rates may be even greater than reported because heat events trigger serious complications from pre-existing health conditions.¹¹⁷ Some people are at higher risk of heat-related illness, including those who are elderly, very young, disabled, poor, or live alone; those with existing cardiovascular medical conditions; and those living in urban areas having high air pollution or in buildings without air conditioning. Urban greening, particularly tree cover, is a heat-mitigation approach.¹¹⁷ Thermal comfort is improved when people spend time in tree-covered spaces.¹¹⁸ Trees have been found to reduce the risk for heatstroke and heat-related ambulance calls during extreme heat events.^{119,}¹²⁰ Groups of trees cool the immediate, and sometimes extended adjacent, area by way of evapotranspiration and shading, and also by affecting air movements and heat exchange.¹²¹ In warmer climates, shaded surfaces can range from 25 °F to 35 °F cooler than the peak temperatures of exposed surfaces.^{122, 123} Planning for placement of specific vegetation types can help control actual and localized effects, potentially creating a more amenable environment for communities.^{124, 125}


EXAMPLE ADAPTATION TACTICS

- ▶ Plan tree projects to reduce ambient summer temperatures in and around higher density residential areas and civic spaces serving more frequent visitors and users (such as transit centers, playgrounds, and plazas).^{126, 127}
- ▶ Plan projects to increase tree canopy cover and shading of heat-absorbing impervious surfaces, such as roads, sidewalks, large parking areas, and roof areas.^{123, 128-130}
- ▶ Plan tree planting or conservation to create patches, as tree clusters provide greater shading and transpiration benefits, though analysis of microclimate conditions is important (such as solar aspect, topography, and wind).^{127, 131}
- ▶ Promote and conserve large trees using local government policy or code and best management practices, as larger trees provide greater area and density of shade for cooling effects and human comfort.¹³²
- ▶ Create thermal refuges in frequently used public spaces (e.g., residential courtyards and neighborhood parks) through careful tree selection and arrangement, and incorporate built elements (e.g., shade structures, pools or fountains, and spray or mist systems) to provide additional respite during high heat events.¹³³⁻¹³⁶
- ▶ Design and locate tree projects to reduce extreme heat in under-resourced neighborhoods and near facilities that serve vulnerable populations, such as schools, hospitals, or elder care facilities.^{135, 137-140}
- ▶ Incorporate native grasses and other understory plants in spaces too small for trees to reduce urban heat island effects and use organic mulch instead of rock mulch to prevent the heat load of parkway strips, parking lot aprons, and other commercial planting spaces.

- ▶ Work closely with local or state health departments on efforts to monitor and report heat-related illness and other climate-related human health impacts to facilitate forest and public health data integration and planning.

Approach 2.2: Improve urban air quality conditions

Climate change is contributing to reduced air quality in many places by modifying weather patterns that increase and focus pollutants, enhance storm and disturbance events that raise particulate levels, and elevate the release of pollen and volatile organic compounds.^{141, 142} Poor air quality, most notably from fine particulates, compromises human respiratory and cardiovascular systems.¹⁴³ Ground-level ozone (generated by volatile organic compounds) and particle pollution can have a range of adverse effects on human health and are responsible for extensive human respiratory illness and mortality each year.¹⁴⁴ Higher pollen concentrations and longer pollen seasons, combined with other aeroallergens, can increase the prevalence and severity of allergic disease.¹⁴⁵ Pollen allergenicity is a seasonal health concern that interacts with broader environmental conditions, such as temperature, humidity, and other air pollutants, to negatively influence human health.¹⁴⁶ Although tree pollen may contribute to illness, some household allergens^{147, 148} and other plants (such as ragweed, which produces the most allergenic pollen¹⁴⁹) pose equal or greater risk. All of these conditions can disproportionately harm older people and children.¹⁵⁰ Awareness and consideration of these complexities are critical when planning health benefit pathways using the urban forest. Reduced asthma has been associated with exposure to natural areas and biodiversity,¹⁵¹ including in children.¹⁵² In some case, trees can intercept particles on leaf and limb surfaces, reducing the amount of respiratory irritants in the air.¹⁵³ They can also absorb



The High Line rail trail and public park in New York City is an example of urban greening that achieves many goals and adaptations. U.S. Department of Agriculture photo.

and remove polluting gases, such as ozone and nitrous oxide.^{154, 155} Careful planning of plant selection, planting density and location, and management can increase health benefits while reducing risks.¹⁵⁶

EXAMPLE ADAPTATION TACTICS

- ▶ Use tree projects at local and citywide scales to mitigate sources of air pollutant compounds and particulates to help protect people from negative health effects.¹⁵⁷⁻¹⁶⁰
- ▶ Select tree species having specific traits when planting to improve air quality, as different leaf structures and surfaces enable better capture at different times of the year,¹⁶¹⁻¹⁶⁵ with coniferous trees being more effective for overall particulate capture.¹⁶⁶
- ▶ Plant trees near major emissions sources, such as industrial or manufacturing sites, to reduce air pollution. Consider species selection, tree size, and leaf area, as well as the position of plantings relative to nearby buildings and other features that can influence wind direction and speed.¹⁶⁷
- ▶ Plan tree projects in alignment with transportation plans to provide “green screens” for high-speed, high-volume transportation corridors, as these are concentrated sources of particulates and emissions that can drift into adjacent residential areas and facilities that serve vulnerable populations (such as schools).^{160, 168, 169}
- ▶ Plan for the interactions of microclimate conditions of wind speed and direction, ventilation patterns, and adjacent structures (especially in urban canyons) to avoid elevating concentrations of aerial particulates due to tree plantings.^{170, 171}
- ▶ Plan tree projects to reduce locally recognized sources of plant-based allergens, such as using fewer male trees to reduce pollen or avoiding species that cause increased allergenic response.¹⁷²⁻¹⁷⁴
- ▶ Select and manage trees to reduce biogenic volatile organic compound emissions to limit contributions to air pollution.¹⁷⁵⁻¹⁷⁷

Approach 2.3: Anticipate and reduce human health impacts of hazardous weather and disturbance events

Climate change is exacerbating the incidence of episodic events and disturbances, such as coastal storms, hurricanes, tornadoes, flooding, and wildfires. These events can take a devastating toll on health within a community,¹⁷⁸ ranging from loss of shelter to reduced potable water, limited food access, and increased infectious disease transmission.^{179, 180} People in urban areas can be at greater risk given extensive transportation and utility infrastructures that are vulnerable to damage, along with higher density residential housing that may be hard-hit by property damage and personal injury. Additionally, increased incidence and severity of wildfire amplifies risk of tragic personal and property loss, especially in the wildland-urban interface, and severely compromises air quality over large areas. Urbanized areas are more prone to certain disturbance events (such as tornadoes

or floods¹⁸¹) than rural or wildland areas because of changing weather cycles. Managed landscapes and forests can be incorporated into emergency planning to prepare for these large, infrequent disturbances.¹⁸² Analysis and preparation before a storm event will help communities minimize harm to trees, property, and people. While trees can become hazards during high-wind events such as hurricanes and tornadoes, proper preparations can make trees more wind-resistant and storm recovery more effective.¹⁸³ Planning and management can also reduce tree and property damage during ice storms.¹⁸⁴ In addition, green infrastructure and natural resources are increasingly valued as buffers that protect human populations against weather-related acute shocks. For instance, maritime forests and forested swamps can be part of hybrid nature and infrastructure systems for coastal defense from storms.^{185, 186} This approach complements, and can be used in conjunction with, approaches 5.2 and 5.3, which describe actions for reducing extreme weather and wildfire impacts on trees and forests.

EXAMPLE ADAPTATION TACTICS

- ▶ Assess and prioritize areas that are vulnerable to extreme events, such as neighborhoods located in storm- or flood-prone areas.¹⁷⁸
- ▶ Conduct hazard and resilience assessments in ways that acknowledge and respond to historic socio-cultural inequities.¹⁸⁷
- ▶ Create, test, and implement urban forestry best practices for tree management prior to a disturbance event, an immediate response system after disturbance, and a mid-term recovery process.^{188, 189}
- ▶ Participate in local emergency preparedness initiatives to engage residents in developing community-based planning and response processes for their neighborhood urban forest.^{187, 190}
- ▶ Engage and educate community residents in tree management best practices and ongoing stewardship on both private properties (such as homeowners' yards) and public lands (e.g., municipal parks and community gardens) to minimize damages and loss during disturbance events.¹⁹¹
- ▶ In areas prone to high winds, implement best practices to manage trees (such as species selection and pruning) to minimize property damage and human injury during storm events.¹⁹²
- ▶ Educate property owners and community members to promote adoption of Stormwise practices of forest and management in urban and suburban areas that are at risk for extreme storms.^{193, 194}
- ▶ Educate property owners and community members to promote adoption of Firewise USA® practices of tree planting and management in urban and wildland urban interface areas that are at risk for wildfires.¹⁹⁵

Strategy 3: Maintain or increase extent of urban forests and vegetative cover

This strategy addresses the foundational role of healthy tree and vegetative cover in urban areas to reduce climate impacts to human health and ecosystem function, while maintaining carbon sequestration and other ecosystem services. Urban forests and other forms of vegetative cover can ameliorate many climate-related threats and help mitigate climate change through reduced energy demands and carbon sequestration. Increasing the extent of urban forests not only allows for enhanced climate adaptation and carbon mitigation benefits for all communities, it is also a primary opportunity for addressing the environmental inequities in our cities. For example, urban trees and greenspace can help reduce the urban heat effect that is exacerbated by climate change.^{123, 196} Green infrastructure, when properly planned and sited, can help minimize urban heat islands through increased shading and evaporative cooling.^{57, 197} Further, carbon mitigation in urban areas reaches its highest levels where tree canopy densities are highest, while simultaneously reducing energy usage in buildings.^{25, 198} Actions that retain or increase tree canopy cover, while preserving the integrity of these ecosystems in the face of climate change, can have some of the most significant benefits for maintaining human health, climate mitigation, and other environmental benefits into the future. Activities within this strategy seek to sustain or enhance the long-term benefits of urban forests on human well-being and ecosystem function by minimizing loss and fragmentation of historically forested areas, maintaining current tree and forest cover, and increasing tree canopy.

Approach 3.1: Minimize forest loss and degradation

Conversion of historically forested areas to other land uses, fragmentation, and degradation threaten the climate adaptation, carbon mitigation, human health, and other benefits these forests provide. Development pressures are a major threat to urban forests; from 2009 to 2014, forest loss due to urbanization was estimated at 175,000 acres per year in the United States.¹⁹⁹ Over the same time period, pavement and other impervious surfaces increased by 167,000 acres annually. Actions to minimize forest loss, landscape fragmentation, and ecosystem degradation can be fundamental to protecting ecosystem services, or the benefits people receive from nature. In some states, such as New Jersey and Maryland, municipalities have created policies or ordinances to reduce forest loss from development and reforest affected areas.^{200, 201} Additionally, planning for new urban infrastructure and site development that takes advantage of existing green infrastructure, including desirable trees, shrubs, and grass cover, will reduce the time and resources needed for vegetation establishment. Making efficient use of existing tree canopy allows for immediate benefits for climate and health for communities.

EXAMPLE ADAPTATION TACTICS

- ▶ Develop urban forest regulations, policies, or plans that reduce land-use change and land disturbance and identify requirements for remediation of disturbed sites.²⁰²

- ▶ Minimize the amount of land disturbed by urban site development and locate hardscaped areas, such as roads, sidewalks, and parking spaces, to minimize negative impacts to existing vegetation.²⁰³
- ▶ Retain and protect existing trees during development of urban green spaces.^{204, 205}
- ▶ Restrict development in priority areas by acquiring property for preserves or using conservation easements on private land holdings to protect natural land cover or maintain corridors between existing natural areas.
- ▶ Implement protective guidelines, such as best management practices and tree heritage and protection ordinances, to avoid unintentional loss of trees during development.^{204 - 207}


Approach 3.2: Maintain existing trees through proper care and maintenance

Urban forests will continue to face increasing pressure from climate change. More active management and investment to promote tree health, survivability, and longevity may be necessary to reduce hazards to human health and safety, and to ensure long-term health and continued provisioning of environmental benefits such as carbon sequestration. In developed urban sites, sustained maintenance of urban trees includes pruning, watering, mulching, and other soil improvements, along with pest and disease monitoring and management, and protection from extreme weather. Although these activities are costly, accounting for roughly a third of urban forest budgets, research has shown the costs of not maintaining trees can be even greater.²⁰⁸ For example, insecticide treatments, especially for large, long-lived trees that can store up to 1,000 times more carbon relative to smaller trees in some locations,²⁰⁹ can be more a cost-effective method for maintaining a tree versus the cost of removing and replacing a tree killed by emerald ash borer.^{210, 211} Street trees, in particular, face challenging growing conditions, with annual mortality rates at 3 to 5 percent, causing the average lifespan of a street tree to be 15 years or less.²¹² Some maintenance activities, such as watering, may become even more necessary as droughts become more frequent or severe.²¹³ In urban natural areas or forest patches, certain activities, such as hazardous tree removal, prescribed burning, and non-native invasive species removal, can be used to maintain or improve forest condition. In addition, some localities are beginning to incorporate traditionally rural silvicultural practices in their urban natural areas management as an integrated forest health and climate change strategy.²¹⁴

EXAMPLE ADAPTATION TACTICS

- ▶ Water individual trees in susceptible locations during extreme droughts and heat waves to reduce mortality.²¹⁵
- ▶ Prune street trees to establish strong branching structure with a central leader to reduce the center of gravity. Remove structural defects in mature canopy trees.

- ▶ Ensure that trees are planted properly, such as at an appropriate depth and with no root girdling, to optimize growth rates and make trees less susceptible to drought and other stressors.²¹⁵
- ▶ Remove mowed turf from the rooting zone of trees and replace it with organic mulch or other plants that require less water and nutrients, also protecting the tree from mower damage.
- ▶ Implement silvicultural practices in urban natural areas to improve the health of the entire forest community, such as thinning forest stands to increase growing space for the remaining trees.
- ▶ Manage stand density and age in forested areas to reduce risks of property damage and hazards from severe weather events, using Stormwise^{193, 194} or other forest management practices.
- ▶ Ensure that newly planted trees have sufficient soil volume to support the tree at maturity and enough space to grow without interfering with underground or overhead utilities.
- ▶ Utilize stormwater to supplement irrigation of street trees and urban forest vegetation.



Diverse tree plantings provide the greatest levels of ecosystem services if given proper care and maintenance. USDA Forest Service Eastern Region photo.

Approach 3.3: Restore and increase tree, forest, and vegetative cover

Efforts to increase urban tree cover and forest canopy are growing across many parts of the United States and globally in recognition of the benefits to human health, climate adaptation, carbon sequestration, and environmental quality. A growing body of research points to the potential of increasing forest cover as a natural climate solution across all ecosystem types²² and in urban areas, in particular.^{23, 216, 217} Urban forestry is highlighted as an important mechanism for increasing stored carbon, as trees in urban areas can have significant biomass and carbon sequestration.¹⁹⁸ Carbon sequestration rates in individual trees within urban areas can exceed those in natural forests due to greater foliar biomass and reduced competition from lower tree densities, as well as irrigation and fertilization²¹⁸—and a changing climate may be further accelerating these growth rates in some urban areas if there is also sufficient moisture.^{219, 220} Trees can have an additional important influence on carbon mitigation in urban zones by reducing the energy requirements for building heating in winter (due to wind protection) and summer cooling (from tree shading).¹⁵⁵ Increasing tree and forest cover in urban areas takes advantage of opportunities to increase canopy in places where trees are not currently present or abundant and will not interfere with other uses of the sites, which may include increasing tree density in green spaces where trees already exist, such as in urban riparian zones,²²¹ as well as afforestation on abandoned industrial or previously developed sites.

EXAMPLE ADAPTATION TACTICS

- ▶ Create parks and green spaces on abandoned or underutilized spaces, such as brownfields and vacant lots.^{222, 223}
- ▶ Plant trees on abandoned land that was cleared for agriculture, mining, or other reversible uses.
- ▶ Add street trees or other vegetation to help “green” areas that currently have low canopy, recognizing many of these areas have been historically disenfranchised.²²⁴
- ▶ Plant trees in strategic locations, such as upwind of areas most prone to extreme heat or in positions to provide maximum building shading or cooling benefits.^{57, 197}
- ▶ Integrate trees as part of low-impact development or stormwater runoff projects.²²⁵
- ▶ Replant forests following disturbances in urban parks.
- ▶ Allow passive reforestation on land that had been cleared for agriculture or other uses and retain tree species that grow quickly and provide cover and shade on sites where passive reforestation is occurring.
- ▶ Use fast-growing native tree species to rapidly create a privacy screen or shade for outdoor spaces or buildings, ensuring proper species selection for long-term sustainability on the site.

- ▶ Establish trees adjacent to urban streams^{221, 226} to create or expand riparian areas and help reduce impacts from extreme heat and floods.
- ▶ Ensure planting stock used for projects has been grown using techniques that provide healthy, vigorous root systems, such as in gravel beds,²²⁷ and provide planted trees with adequate soil volume for root growth and anchoring.
- ▶ Use urban tree canopy inventories and remotely sensed data to identify new and existing areas that can serve as wildlife corridors or greenways and inform reforestation efforts.
- ▶ Construct wildlife corridors between natural areas to mimic habitat and ecosystem structure of the natural habitat (e.g., by maintaining or promoting oak (*Quercus spp.*) canopy cover in a residential development between two oak woodland natural areas) while avoiding the creation of ecological traps in these areas.²²⁸
- ▶ Plant native shrubs, grasses, or herbaceous plants in places where there is insufficient space for mature trees, such as parkway strips, medians, or other narrow or restricted planting zones.
- ▶ Create pollinator habitat with native shrubs and herbaceous plants along roadways or transportation corridors that are inappropriate tree-planting areas or are adjacent to forest edges.

Approach 3.4: Sustain sites and ecosystems that provide high value across the landscape

Urban forests provide numerous ecosystem services and benefits, and some sites or ecosystems may provide disproportionately high benefits for biodiversity, carbon storage, or other services. Some urban sites currently support high levels of biodiversity and contain geophysical characteristics that are likely to sustain and promote diversity even as the climate continues to change. Urban areas have the potential to host a significant percentage of locally occurring native species, including endangered species and other species of concern. These sites may be quite rare, however, because many pre-urban forest ecosystems have been fragmented, degraded by human use and invasive species, or developed into other land uses.²²⁹ Remnant forest ecosystems can provide a suitable habitat for relict populations of species that were previously more widespread^{12, 230} and act as refugia by providing habitat for species lost from surrounding areas due to human-caused disturbance. Sites with several topographically related microclimates and local permeability may provide the best chance for species responding to climate change.²³¹ Where forest carbon is valued, remnant forests may provide the greatest carbon densities and sequestration rates in developed areas,²³² while riparian areas may provide good opportunities for maintaining carbon mitigation both in biomass and in soils.²³³ Restoration or reclamation projects may be needed to increase the representation of these habitats on the landscape or maintain the values associated with these systems. This approach places additional emphasis on targeted efforts to maintain and restore ecosystems that have been identified as high value, which extends efforts under Approach 3.1: Minimize forest loss and degradation.

EXAMPLE ADAPTATION TACTICS

- ▶ Identify areas with high diversity (species, topography, soils, or other factors) or other desirable attributes that can be set aside as natural areas or reserves, perhaps with the support of conservation easements, public tax funds used for municipal acquisition, or similar tools.
- ▶ Protect existing habitat remnants from loss, conversion, or invasion from nonnative plants, particularly if they are in areas that may provide future climate refugia.²³⁴
- ▶ Restore unique habitats that may be less susceptible to climate change or use reclamation efforts to create new patches of such habitats on suitable sites.
- ▶ Identify and protect areas of high geophysical or topographic diversity with the expectation that these areas may provide a range of climatic options to species with diverse requirements.²³⁴
- ▶ Identify urban plantings within developed landscapes that could serve as climate refugia.
- ▶ Establish and support development and management ordinances and regulations that protect and reduce impacts to high-quality forest remnants and ecological features.
- ▶ Use urban tree canopy inventories and remotely sensed data to identify new and existing areas that can serve as wildlife corridors or greenways.
- ▶ Manage natural areas that serve as wildlife corridors to promote their maximum habitat value (e.g., by removing invasive species) and prioritizing management in those locations.²³⁵
- ▶ Manage riparian corridors within otherwise highly developed landscapes to provide habitat value and ecosystem services.^{236, 237}


Strategy 4: Sustain or restore fundamental ecological functions of urban ecosystems

This strategy emphasizes ecological processes and functions to preserve the capacity of systems to cope with changing and more variable climate conditions. Complex interactions among shifting climate, vegetation, and landforms may result in changes in ecosystem hydrology, soil quality, and nutrient cycling.²³⁸ Ecosystems in urban environments are shaped not only by these changes, but also by the people who have shaped the landscape in the past, along with those who currently live, work, and recreate in them. Challenges to maintaining natural ecosystem functions in urban areas include impermeable surfaces, air and water pollution, frequent human disturbance, and altered soil characteristics. Climate change impacts can exacerbate these issues via extreme events and disturbances on ecosystems that may already be under stress or otherwise disrupted.²³⁹ For example, urban forests in coastal areas are increasingly susceptible to the impacts from sea-level rise, coastal flooding, salinization, and storm damage. Climate change impacts, either alone or interacting with other stressors, can impair the health and productivity of urban trees, thereby reducing

human health benefits for communities and the carbon mitigation capacity of urban forests. This strategy seeks to sustain or enhance ecological functions to reduce the potential negative impacts of a changing climate on urban forests.

Approach 4.1: Maintain or restore soils and nutrient cycling in urban areas

Urban soils provide a critical foundation for the health and productivity of urban forests. Poor soils and growing conditions cause most urban tree problems,²⁴⁰ and these less favorable conditions will be exacerbated by climate change.²⁴¹ Urban soils vary across a continuum from undisturbed to highly engineered;^{241, 242} soils in disturbed sites can lack essential nutrients and commonly include detrimental elements, such as chemicals, concrete, asphalt, and other foreign matter, that limit the long-term viability of a tree.^{242, 243} Rising temperatures can increase drought conditions and alter nutrient cycling in all forests; the urban heat effect can intensify these impacts. Extreme rain events increase the potential for greater stormwater runoff and erosion.²³⁹ Minimizing impacts to soils and restoring natural function can increase the benefits of urban soils. For example, although soil characteristics such as carbon content can vary widely in urban landscapes, urban soils store large amounts of carbon in organic matter²⁴⁴ and have a substantial capacity to sequester carbon, especially in residential areas or other locations with lower levels of disturbance and other significant management inputs.²⁴⁵ This approach focuses on preserving and restoring natural soil processes as a way to sustain urban trees and forests; it complements other approaches focused on maintaining urban ecosystems and reducing the impacts from extreme weather.²⁴⁶



Urban parks and natural areas can provide important ecological functions and human health benefits. Courtesy photo by Sophie Nito, used with permission.

EXAMPLE ADAPTATION TACTICS

- ▶ Analyze soil conditions prior to tree-planting activities to determine whether conditions are sufficient for healthy tree growth and identify deficiencies that need to be addressed.
- ▶ Improve growing conditions for revegetation or restoration efforts by adding organic matter amendments, such as mulch or biochar,²⁴⁷⁻²⁴⁹ which can help improve drainage, pH, soil carbon storage, and rooting conditions.
- ▶ Identify areas that have had minimal soil disturbance and include high-quality soil conditions as a consideration when creating reserves or undeveloped areas.
- ▶ Prevent or reduce soil erosion in areas prone to soil loss during heavy rainfall events, particularly when soils are exposed following disturbance.

Urban natural areas

- ▶ Remove invasive species that negatively affect soil processes or alter nutrient levels, such as European buckthorn (*Rhamnus cathartica*).²⁵⁰
- ▶ Add organic soil amendments (e.g., mulch, biochar) to urban sites undergoing restoration or revegetation.^{247, 248}
- ▶ Inoculate soil with mycorrhizal fungi to increase organic matter and improve nutrient cycling and moisture retention.²⁵¹

Developed urban sites

- ▶ Provide adequate root volume in tree planters while ensuring soil conditions (texture, pH, nutrient levels) match tree requirements.^{252, 253}
- ▶ Provide and develop adequate soil volume, texture, structure, and organic matter to support healthy tree growth.²⁵³⁻²⁵⁵
- ▶ Remove and replace the soil if toxicity or chemical levels are too high.
- ▶ Install a layer of mulch over the root zone of the tree to help retain moisture and mimic a natural growing environment.^{253, 256}
- ▶ Rebuild the soil profile following development.^{257, 258}

Approach 4.2: Maintain or restore hydrologic processes in urban forests

Climate change is altering precipitation patterns and increasing the frequency and intensity of rainfall and storms in many areas.²³⁹ Urban forests and vegetation help to maintain urban hydrologic processes by intercepting, absorbing, and filtering rainfall and stormwater, which can reduce runoff and improve the quality of water reaching streams and lakes.²⁵⁹ Likewise, vegetation can help retain soil moisture, which helps support urban forests and tree canopy and the associated benefits for human health and climate mitigation.²⁶⁰ Many cities are recognizing the value of this green infrastructure and increasing tree planting and other efforts to improve vegetative cover, especially in riparian areas. Riparian forests, wetlands, and floodplain forests serve important ecosystem functions, such as decreasing soil erosion, filtering water, and storing and recycling organic matter and nutrients,²⁶¹⁻²⁶³ along

with elevated carbon benefits.^{23, 233} Trees in riparian areas also provide shade, which helps to buffer stream temperatures. Forested riparian areas can serve as corridors for wildlife and plant species migrating across otherwise fragmented landscapes¹⁶ and provide substantial co-benefits for biodiversity and carbon storage.²³³ Urban wetlands, whether remnant, intentionally restored, or “accidental,” can also play an important role in hydrological functioning and other ecosystem services.^{263, 264} This approach focuses on maintaining or restoring natural ecosystems and features as elements of the urbanized landscape to protect water quality and cycling.

EXAMPLE ADAPTATION TACTICS

- ▶ Restore natural hydrology where appropriate by removing drain tiles or other remnant hydrological modifications.²⁶⁵
- ▶ Restore or reforest native communities and ecosystem components (e.g., natural groundcover, litter layer, coarse woody debris) in riparian areas, particularly those adjacent to developed areas, in order to reduce erosion and nutrient loading into adjacent water bodies.^{226, 265, 266}
- ▶ Adjust the location and design of trails in natural areas to minimize erosion under more intense surface runoff.^{267, 268}
- ▶ Restore or promote a diversity of riparian tree and plant species in order to increase stream shading, provide a source of woody debris, stabilize the soil, and provide habitat and connectivity for wildlife.²⁶⁵
- ▶ Manage water levels to supply proper soil moisture to vegetation adjacent to the stream during critical time periods, either by manipulation of existing dams and water control structures or restoration of natural dynamic water fluctuations.
- ▶ Locate new natural areas or green space in areas prone to ponding or flooding to add additional water storage capacity during extreme events.
- ▶ Direct water into natural features with herbaceous and woody plant cover to reduce runoff and nonpoint source pollution, while still providing outflow for excess water.
- ▶ Reforest floodplain forests with high degrees of mortality from emerald ash borer with a climate-adapted mix of trees.²¹⁴
- ▶ Take advantage of “accidental” wetland areas that arise in urban areas, using management to augment desired services and minimize disservices.²⁶⁴
- ▶ Connect elements of green infrastructure, such as planting beds, bioswales, rain gardens, and sequential stormwater treatments to natural systems.

Approach 4.3: Restore or maintain fire in fire-adapted ecosystems

Restoring natural fire regimes can help reduce ecosystem vulnerability to a changing climate, especially in areas that are susceptible to increases in wildfire under hotter, drier conditions.²⁶⁹ Using fire as a management tool can be difficult in urban settings due to

potential or perceived impacts to the built environment and public health.²⁷⁰ However, residents are generally supportive of prescribed fire if it can reduce risks of wildfire or increase native plant diversity in suburban and wildland-urban interface areas.²⁷¹⁻²⁷³ Where possible, fire can be an important management strategy in supporting ecosystem function and resilience. For example, even small prescribed fires in suburban forest patches can lead to increases in species richness and diversity.²⁷⁴ Prescribed fire can reduce wildfire risk and severity, improve tree survival, and prevent loss of forest conditions.²⁷⁵ Prescribed fire also can have long-term carbon mitigation benefits through reductions in risk of large-scale carbon losses from wildfires, although carbon may be reduced in parts of the forest (such as the forest floor). For example, conditions during implementation of prescribed fire typically result in low overstory tree mortality rates, preserving both carbon in live trees and the potential to sequester future carbon through tree growth.²⁷⁶ Additional care can also be taken to reduce potential impacts on public health through timing of prescribed fire to minimize air quality effects and providing advanced communication to community residents.^{277, 278} Where ecological or social constraints limit the application of prescribed fire, alternative management strategies (i.e., fire surrogates) can provide some benefits, but do not fully mimic the effects of prescribed fire.^{279, 280} This approach complements, and can be used in conjunction with, approach 5.3, which describes actions for reducing the risk of wildfire in ecosystems.

EXAMPLE ADAPTATION TACTICS

Urban natural areas

- ▶ Use prescribed fire to maintain fire-adapted ecosystems and reduce risk of fire spread into the wildland-urban interface.²⁸¹⁻²⁸³
- ▶ Use prescribed fire during suitable conditions (periods of low air pollution, low winds, low temperatures) to avoid negative impacts and potential for unwanted spread.
- ▶ Provide advance warning to community residents about prescribed fire activities and efforts to minimize adverse impacts from smoke.²⁷⁸
- ▶ Incorporate understory thinning, mowing, or other fire surrogate strategies to support native ecosystems in addition to prescribed fire or where fire management is not possible.²⁸⁴

Developed urban sites

- ▶ Manage fire-adapted urban trees and ecosystems using fire-surrogate treatments, such as understory thinning, mowing, hand-weeding, and appropriate herbicide application.

Strategy 5: Reduce the impact of physical and biological stressors on urban forests

Urban forests are experiencing increasing threats as a result of altered climate conditions and interactions with other environmental stressors.²³⁹ The stressors that affect urban

forests vary widely based on the impacts of climate change in a particular region or area, as well as local factors that influence exposure and sensitivity to climate change.^{239, 285} For example, cities in the western United States may be at greater risk of wildfire conditions given development in the wildland-urban interface,^{239, 286} while coastal cities are likely to be at greater risk of sea-level rise, coastal storms, and hurricanes.^{239, 287} Although the nature and severity of climate risks will vary for individual cities, there are many threats in common across urban areas. For example, climate change is expected to increase the impact of biological stressors, such as insect pests, pathogens, and invasive plant species,²⁸⁸⁻²⁹² urban areas may be especially prone to these stressors as a result of urban warming, human disturbance, and proximity to points of introductions.²⁹³


Approach 5.1: Reduce impacts from extreme rainfall and enhance water infiltration and storage

Urban forests can be used in combination with gray and green stormwater infrastructure to reduce harm from extreme rainfall to people, the built environment, and vulnerable soils. Natural and low-impact development techniques help to reduce stormwater conveyance, enhance groundwater recharge, and improve water quality by decentralizing flows and using soil and plants to capture and filter pollutants.²⁹⁴⁻²⁹⁷ Increasingly severe rainfall events interact with impervious surfaces to concentrate stormwater flows, often exceeding the capacity of urban gray infrastructure (e.g., pipes and sewers) to direct water to desired locations. Concentrated stormwater runoff can cause adjacent areas to erode, flood, destabilize stream channels, and impair water quality.^{295, 296, 298} Further, stormwater that is directed into water bodies through storm sewers bypasses vegetation and other natural features and can discharge untreated sewage and stormwater directly into surface waters, reducing water quality and creating human health risks. Trees and other green infrastructure can help manage excess stormwater volume at its source, reducing the burden on stormwater collection systems through canopy interception, evapotranspiration, and improved soil infiltration.²²⁵ Vegetation and soils can also reduce nutrient loading in stormwater, as runoff often contains nutrients, such as nitrogen and phosphorus, that are detrimental to surface water quality, yet important for plant growth.^{225, 299} Prioritizing the use of native plants and trees for this approach, when feasible, can help increase wildlife habitat for a wide range of species. Implementing green infrastructure approaches may be more effective in managing excess stormwater flows and far less expensive than upgrading gray infrastructure systems.²²⁵ While the primary benefits are hydrological regulation and water quality improvement, green infrastructure also provides notable carbon sequestration and storage benefits.³⁰⁰

EXAMPLE ADAPTATION TACTICS

- ▶ Incorporate a mixture of plant functional types to provide year-round interception and evapotranspiration benefits (e.g., deciduous and coniferous trees, turfgrass).^{225, 299}
- ▶ Manage the urban forest to increase leaf surface area; when planting, select species with greater leaf surface area and/or rough-surfaced leaves and bark.^{299, 301}

- ▶ Use ground covers and consider underplanting with smaller trees or shrubs to increase surface area for interception.²⁹⁹
- ▶ Retain as much tree canopy as possible to intercept rainfall; encourage canopy growth over impervious surfaces.²⁹⁹
- ▶ Provide appropriate care and maintenance for trees to ensure health and continued function.²⁹⁹
- ▶ Maximize belowground soil volume and use biological mulches below tree canopy to improve water infiltration and storage.²⁹⁹
- ▶ Select species based upon their intended use and transpiration capabilities; for example, species that transpire water in greater volumes (such as tulip tree [*Liriodendron tulipifera*], black gum [*Nyssa sylvatica*], birch [*Betula* spp.], dogwood [*Cornus* spp.], red maple [*Acer rubrum*], sycamore [*Platanus* spp.]) can be planted in areas that that receive stormwater.²⁹⁹
- ▶ Plan ahead to manage stormwater in the event of widespread tree loss due to storm damage or pest outbreaks.²²⁵
- ▶ Strategically grade soil where needed and avoid unnecessary soil disturbance to preserve soil porosity and natural drainages.



High density urban streetscapes need extra care to reduce the impacts of physical and biological stressors. USDA Forest Service photo.

Developed urban sites

- ▶ Integrate trees into other types of green infrastructure, such as rain gardens, to enhance their capacity and improve the regulation of soil moisture content.³⁰²
- ▶ Select water-tolerant tree species to plant in shallow, concave settings to collect runoff.²²⁵
- ▶ Improve soil quality and minimize compaction to increase rooting volume and infiltration capacity.
- ▶ Make use of structural soils systems (mix of mineral soil and coarse stone), such as Silva Cells™ or StormTree™, or suspended pavement over noncompacted soils.^{225, 303}
- ▶ Use permeable paving, suspended surfaces, or Silva Cells³⁰⁴ to enable runoff to collect and water trees.
- ▶ Incorporate permeable surfaces into designs, such as block pavers, porous asphalt, and concrete, to reduce hardening of surfaces and increase infiltration of storm flows.
- ▶ Attenuate and treat stormflows in depressional areas using bioretention systems to capture runoff, recharge groundwater, and reduce pollutant loads.²⁹⁶
- ▶ Divert and disperse stormwater off of impervious surfaces (such as walkways, roofs, roads, trails) to forests, densely vegetated areas, swales, and filter strips to increase water retention on site and enhance filtering of water.²⁹⁶
- ▶ Use vegetated shoulders and embankments of compacted soils to resist storm surge water flowing across a roadway.

Approach 5.2: Reduce risk of damage from extreme storms and wind

Climate change has increased the likelihood and severity of storms,³⁰⁵ which damage urban trees. Urban tree failures can cause severe property damage, electric outages,³⁰⁶ and human injuries or fatalities.³⁰⁷ Damage can also reduce the carbon mitigation capacity of urban forests and impair other ecosystem services.¹⁸⁹ In developed urban sites, these potential impacts make it essential for risk to be managed at the individual tree level.³⁰⁸ Factors such as tree form, size, condition, species, wind speed, pruning, and wood material properties affect tree resistance to storm damage.³⁰⁹⁻³¹² Urban foresters may need to balance the risk of tree failure, including the danger to people and property, with the loss of benefits when shade trees are removed. This is especially important for large trees, which provide greater benefits and take a long time to replace.³¹³ Although intensive hazard management is common in developed areas, it may also take place in urban natural areas where individual tree management is necessary or possible, such as in high-use areas or near infrastructure where damaged trees present hazards. Elsewhere, management approaches in natural areas are more likely to be focused on minimizing the impact of disturbances on tree communities. Public involvement and education are critical factors in wide and effective implementation of this approach.

EXAMPLE ADAPTATION TACTICS

- ▶ Develop disaster management plans and risk assessments to prepare for more frequent extreme storms.¹⁸⁹
- ▶ Create education and outreach programs for arborists, utilities, and the public to ensure tree removals are necessary.
- ▶ Monitor for hazard trees near potential targets (e.g., playgrounds, walking paths, or roads), and repair or replace these trees as quickly as possible.³¹⁴
- ▶ “Soften” the edges of forests and natural areas (i.e., reduce the edge influence at regenerating edges and minimize abrupt transitions) to reduce susceptibility to wind damage.³¹⁵
- ▶ Manage stand density and age in forested areas to reduce risks of property damage and hazards from severe weather events using [Stormwise](#) or other forest management practices.^{193, 194}
- ▶ Implement a predisturbance structural pruning program to improve tree health and ability to withstand extreme weather events.¹⁹²
- ▶ The main determinant of a tree’s ability to withstand extreme winds is a strong root system; when planting, provide ample area for future root growth and maintain adequate distance from restrictive pavement. If soil space is limited, select smaller-maturing species.³¹⁶
- ▶ Select species that are more tolerant of high winds to improve wind-resistance, plant trees in groups rather than as solitary individuals.^{192, 316}
- ▶ Use structural pruning to reduce safety and infrastructure issues by fostering mechanically strong branch structure.^{317, 318}
- ▶ Avoid planting trees susceptible to breakage in areas with high wind exposure or species susceptible to damage from ice loading.¹⁸⁹
- ▶ Develop programs for waste wood utilization to facilitate tree maintenance and forest management.³¹⁹
- ▶ Ensure nursery stock is properly pruned and has no circling roots.

Approach 5.3: Reduce risk of damage from wildfire

The increases in the frequency, size, and severity of wildfires in the United States have included greater incidence and risk of wildfires in the wildland-urban interface, especially in the western part of the country.³²⁰⁻³²⁴ This corresponds with an expansion of development in fire-prone areas in the last several decades.³²⁵ Methods for reducing risk to urban natural areas and surrounding communities may be similar, though on a smaller scale, than those employed in rural areas. However, due to their proximity to human communities and infrastructure, urban natural areas are of especially high priority in risk reduction.^{283, 320} Developed landscapes in the wildland-urban interface may incorporate somewhat different

principles of risk reduction to both property and vegetation through altering forest structure and composition,²⁸³ such as the Firewise USA® methods of creating defensible space around structures and using less-combustible landscaping.¹⁹⁵ In some situations, this approach can be used in conjunction with approach 4.3, which describes actions for using fire as a management tool in fire-adapted systems. Public involvement and education are critical factors in wide and effective implementation of this approach.

EXAMPLE ADAPTATION TACTICS

- ▶ Create, update, and implement a community wildfire protection plan: a plan that communities create in collaboration with emergency management and land management agencies to reduce wildfire risk.³²⁶
- ▶ Engage the public, decision-makers, and thought leaders in discussions about land use planning and regulations to reduce wildfire risk.³²⁷
- ▶ Use prescribed fire where possible or fire-surrogate treatments to manage the woody understory and other potential fuels to reduce the risk of wildfire.²⁸⁴
- ▶ Cooperate with appropriate authorities to train municipal fire fighters to respond quickly and appropriately to fires in natural areas.
- ▶ Promote treatments on private lands that reduce the risk of wildfire by creating defensible space around homes and other structures and by removing fuels,³²² including Firewise USA® practices.¹⁹⁵
- ▶ Avoid using highly flammable species in plantings near natural areas, such those containing pine or eucalyptus trees or chaparral shrubs.
- ▶ Avoid using highly flammable landscape materials (e.g., pine straw, shredded bark mulch) near buildings located near natural areas.



Careful tree selection supports sustainable urban forests that provide aesthetic and health benefits. USDA Forest Service Eastern Region photo.

Approach 5.4: Maintain or improve the ability of forests to resist pests and pathogens

Forest pests and pathogens create substantial challenges to urban trees and forests and their ability to provide many ecosystem services.³²⁸ These challenges are expected to be amplified by climate interactions with pest and pathogen ranges and life cycles.³²⁹ Insect pests, such as the mountain pine beetle in the western United States and the southern pine beetle and hemlock woolly adelgid in the eastern states, have expanded their ranges northward, at least partly in response to warmer climate conditions.^{289, 290, 329} The urban heat effect can further increase the risk in cities, particularly when urban heat also contributes to drought stress.²⁹³ Urban forests may also have higher levels of plant stress due to suboptimal growing conditions,³³⁰ sometimes combined with lower tree species diversity and associated reduction of ecosystem resistance to pests.²⁹³ Avoiding the introduction of pests and pathogens is often not possible. Reducing or eliminating stressors that might make a tree more susceptible to new or existing pests or pathogens will be important to maintaining forests in urban areas.³³⁰ While interventions to prevent tree mortality from pests and disease can be expensive, the cost is often less than tree removal and replacement.^{210, 211}

EXAMPLE ADAPTATION TACTICS

All urban sites

- ▶ Increase age diversity to help avoid large concentrations of declining trees that may be more susceptible to certain pests or pathogens.
- ▶ Monitor for new invaders so action can be taken before the pest or pathogen becomes established.
- ▶ Participate in a rapid response system for pest and pathogen detection, including training volunteers and local organizations to assist with identifying pests and pathogens.
- ▶ Use new remote sensing to detect pest and disease outbreaks.^{331, 332}
- ▶ Plant pest- and disease-resistant genotypes of native species when undertaking forest restoration projects.

Developed urban sites

- ▶ Select species and cultivars that are less susceptible to pests and pathogens.³³³
- ▶ Treat susceptible trees with pesticides and fungicides prior to infestation or use behavioral manipulation techniques to disrupt insects.
- ▶ Ensure adequate watering to avoid drought stress in susceptible trees.²⁹³
- ▶ Use new technology and data analysis techniques to monitor tree health at extremely fine scales, including for individual trees.^{334, 335}
- ▶ Apply sanitation practices, including removal of infected trees or use of pesticides, to limit spread.³³³

- ▶ Promote diversity across taxonomic levels by reducing the concentration of any one genus, species, or cultivar in order to reduce the risk from a selective pest or pathogen.³³³
- ▶ Encourage the use of best management practices that limit the spread or level of damage caused by pests or pathogens.

Approach 5.5: Prevent invasive plant establishment and remove existing invasive species

Invasive plant species—that is, species that are not native to an area and whose introduction causes harm—already pose significant problems in many urban forests, and climate change is likely to increase the rate of spread of invasive plant species in several ways.^{291, 292, 336} Urban areas are especially susceptible to introduction and spread of invasive plants due to the proximity of urban areas to the global transport linkages that are often points of introduction. Further, the conditions commonly found in urban areas give invasive plants a competitive advantage; this includes high levels of disturbance, nutrient loading, and warm or moderated microclimates. Warmer temperatures and altered climate conditions may remove existing climatic restraints and allow species to move into new areas, while increases in climate-related disturbances and changing ecosystem dynamics may create conditions that allow new species to invade more easily.²⁹¹ Early detection and rapid response will be very important as opportunities for new invaders increase.



Mixed species plantings can provide leafy, shaded streets and better resist pests and pathogens. USDA Forest Service Eastern Region photo.

EXAMPLE ADAPTATION TACTICS

- ▶ Manage and monitor natural area buffers to limit the spread of invasive plants to high quality or unaffected areas.
- ▶ Limit the spread of invasive species introduced through recreational activities (e.g., boot brushes at trailheads).
- ▶ Use stewardship groups and volunteers as part of a rapid response program to limit the spread of and remove invasive species.
- ▶ Avoid the use of known invaders in horticultural plantings (especially in and around natural areas) through partnerships with homeowners, municipalities, and park districts.
- ▶ Train local land managers, landowners, volunteers, and organizations to recognize possible threats and report them to appropriate local and state agencies.
- ▶ Plant a dense layer of larger native trees to shade out invasive plants in the understory.
- ▶ Remove existing invasive species using nonchemical treatments, such as directed flame or hot foam.^{337, 338}

Approach 5.6: Manage herbivory to promote regeneration, growth, and form of desired species

Changes in habitat quality and extent associated with climate change and increasing urbanization and fragmentation of natural areas are likely to alter the interaction of mammalian herbivores with managed systems, potentially resulting in ecosystem degradation.³³⁹⁻³⁴¹ Herbivory can cause substantial damage to desired plant species; for example, herbivory from white-tailed deer (*Odocoileus virginianus*) has been linked to reduced understory and native plant diversity, a lack of natural regeneration, increased drought stress, and enhanced susceptibility from invasive plant species in forest ecosystems.³⁴² It is difficult to predict how individual wildlife species will respond to climate change given complex interactions with the local environment and management.³⁴³⁻³⁴⁵ White-tailed deer populations are generally expected to remain high or increase as a result of more favorable climate conditions and also high behavioral plasticity.^{346, 347} Protecting desired species from herbivory can effectively reduce browse damage and prevent associated impacts of reduced growth and carbon sequestration. Managing herbivory also can be important in fostering resilience to other stressors that are exacerbated by climate change. This approach may be combined with tree-planting efforts or forest management activities that release advance regeneration or stimulate new regeneration.

EXAMPLE ADAPTATION TACTICS

- ▶ Use community-based management techniques to develop deer management plans based on management goals and community capacity.^{339, 348}
- ▶ Manage populations of deer or other herbivores using control methods.^{339, 349, 350}

- ▶ Apply repellent, tree tubes, bud caps, and other physical barriers to protect individual plants, especially during planting projects.
- ▶ Use fences to protect sensitive plant communities or restoration areas.
- ▶ Promote abundant regeneration of multiple species to supply more browse than herbivores are expected to consume.

Strategy 6: Enhance taxonomic, functional, and structural diversity

This strategy addresses the value of diverse ecosystems in supporting the adaptive capacity of urban areas under changing conditions. Uncertainty about the continued pattern and effects of climate change is a hallmark challenge of adaptation planning, and encouraging diversity in a range of ecosystem components is often considered a “no regrets” investment in ecosystem resilience.^{351, 352} Promoting species and structural diversity is as important in urban forests as in non-urbanized forest landscapes, if not more so. Urban areas are highly susceptible to introduction of nonnative pests and pathogens and often exhibit high occurrence of invasive plant species.^{203, 353} Examples of urban tree mortality demonstrate the role that species diversity can play in the face of pest introductions.^{354, 355} Widespread awareness has led to guidelines focused on diversification of the urban forest.³⁵⁵ However, urban areas contain sites with challenging growing conditions, and only a limited set of tree species may be able to tolerate such conditions.³⁵⁶ Species and structural diversity are especially important as both a climate adaptation and mitigation strategy because urban habitats (both natural areas and developed land uses) are likely to be stressed in the future in many ways, some of which will be unforeseeable.^{56, 357} A diverse set of species, carefully selected to match the urban environment, will be more likely to maintain adequate forest cover, carbon mitigation, and other ecosystem services under a changing and increasingly variable climate.

Approach 6.1: Enhance age class and structural diversity in forests

Diverse age structures can be beneficial in both developed and natural areas because trees are most vulnerable to specific stressors at different ages, an especially important consideration as climate exacerbates many forest stressors. For example, while both droughts and wind events are increasing in the changing climate, droughts typically are more damaging to seedlings than to mature trees, whereas older trees may be more susceptible to damage from wind events. Increasing age—and, perhaps correspondingly, size—diversity can increase the habitat value of the urban forest and spread out tree losses from natural mortality.^{12, 37} Further, greater diversity in tree ages can benefit forest carbon sequestration capacity.^{358, 359} Some urban forests are dominated by trees that established before the urbanized landscape; these legacy trees are now reaching the end of their lifespan.³⁶⁰ In natural areas, active forest management may be necessary to promote the development of multiple age classes, while maintaining greater amounts of dead wood, including snags, downed logs, and coarse woody debris, may increase carbon storage and enhance wildlife

habitat.^{361, 362} In contrast, managers often focus on individual trees in developed urban areas, which are removed upon death or damage and replanted as soon as resources permit. In developed sites, some planting and tree removal practices could help develop a more diverse tree age structure both within and among management units. In addition, pre-urban legacy trees often provide much of the carbon mitigation value, along with other ecosystem services and functional value (e.g., shading and habitat) of the urban forest,³⁶⁰ emphasizing the importance of preserving older and larger legacy trees. Preservation of these features will be essential to adapting the urban forest to future climates, as old trees may have superior genetics and may play a valuable role in helping species persist on the landscape.³⁶³

EXAMPLE ADAPTATION TACTICS

Urban natural areas

- ▶ Implement silvicultural practices in urban natural areas to promote multi-aged forests, such as single-tree or group selection methods.³⁶⁴
- ▶ Restore or create conditions that allow tree seedlings to thrive by removing nonnative species in the shrub layer and canopy trees while controlling herbivory.^{365, 366}
- ▶ Implement silvicultural practices to reduce competition around large, long-lived trees to maintain tree health, such as thinning dense stands from below or using prescribed fire in fire-adapted ecosystems.
- ▶ Retain and protect large, old, or long-lived tree species during forest management activities, such as invasive species treatment, removal of hazardous trees, and development of recreational trails.
- ▶ Retaining snags and downed trees when possible (i.e., those that pose no threat to people or infrastructure).



Careful plant selection, with attention to sense of place, creates appealing outdoor activity spaces. Courtesy photo by Guy Kramer, used with permission.

Developed urban sites

- ▶ Retain legacy trees from the pre-urban landscape during development or redesign of urban areas.
- ▶ Plant replacement trees in anticipation of mortality from pests and disease and before actual loss of canopy trees to diversify age structures.
- ▶ Rotate planting schedules so that removal and replanting is dispersed geographically, thereby avoiding complete tree removal or replanting within a single area (e.g., street or park) during a single year.
- ▶ Plant species of different average lifespans on the same block.
- ▶ Provide adequate clearance from structures, driveways, sidewalks, and other established trees when planting long-lived trees to avoid resource competition or future obstructions to growth.
- ▶ Improve growing conditions for large, old trees, such as improving soil infiltration or drainage conditions, installing irrigation and protective barriers during construction, enhancing soil fertility, or limiting soil compaction from pedestrian or automobile traffic.
- ▶ Enact tree preservation ordinances to protect older age classes of trees.

Approach 6.2: Maintain or enhance diversity of native species

Climate change threatens many native tree species and communities and is expected to reduce the ability of these systems to ameliorate climate impacts and sequester carbon over the long term.³⁶⁷⁻³⁶⁹ Although urban forests serve important ecological and social functions through a wide variety of native and nonnative plants, native species and ecosystems can provide elevated benefits in regard to some ecosystem services, such as pollination, relative to nonnative plants.³⁷⁰ Native plants that are adapted to both current urban conditions (which are often extreme) and anticipated future conditions can be used to restore and enhance species diversity in urban systems.^{37, 352, 371} Increasing the abundance of native species that are expected to persist into the future can have multiple important positive effects on the adaptation potential of the urban ecosystem.³⁷ For example, native species planted in urban locations can provide important habitat value for wildlife species; some areas, such as parks, may be able to emulate a functioning ecosystem to some degree and support functioning food webs.^{372, 373} Native plant ecosystems also may be able to provide migration corridors through intensely fragmented urban landscapes.³⁷⁴ It is important to recognize that it may be more difficult to promote native species diversity in urban systems due to more extensive planting and promotion of nonnative species, including invasive species.²⁰³ This approach may be used in conjunction with approach 7.3, which identifies the potential for novel species introductions.

EXAMPLE ADAPTATION TACTICS

Urban natural areas

- ▶ Restore or create conditions that allow for successful regeneration of a diverse mix of native species, which may require removing some existing trees to open the canopy.³⁷⁵

- ▶ Supplement natural regeneration by planting desired native species to add diversity.
- ▶ Identify and expand the size of urban natural areas to support a larger array of native species.²²⁹

Developed urban sites

- ▶ Identify native tree species for urban tree planting lists that provide important co-benefits, such as high value for wildlife³⁷ or adaptability to environmental or climate stressors.³⁶⁹
- ▶ Use native plant species as ground cover or horticultural plantings in the root zone of urban trees.

Approach 6.3: Optimize and diversify tree species selection for multiple long-term benefits


Warmer temperatures, altered precipitation patterns, and other changes in the climate are expected to affect the growth, productivity, and distribution of tree and plant species.³⁶⁸

³⁶⁹ Individual plants will respond differently based on the combination of functional traits (e.g., broadleaf versus conifer), species characteristics, and site-level characteristics, which increases the need to ensure that species are matched to both current and anticipated future conditions. The “right plant, right place” consideration that is already used in urban forestry can be extended to consider emerging hazards that may affect plant success during its entire lifespan.^{352, 355} Species selection for tree plantings in public spaces, such as streetscapes and parks, and also on private properties, can be used to enhance functional and taxonomic diversity and reduce the risk from any one species failing under future conditions. Current species selection criteria often include site tolerance, aesthetics, community input, maintenance required, and mature size. Climate adaptability can be added as an additional criterion through which to evaluate planting lists, as well as for desired co-benefits such as carbon storage, health benefits, or wildlife value (appendix 3). Species or functional traits, such as soil requirements, susceptibility to breakage, or phenology, can help identify trees that are less susceptible to extreme weather or other disturbances.¹⁸⁹ By altering lists of recommended species for planting or adjusting which species are planted at a given site, urban foresters can influence the taxonomic diversity of their urban forest and increase the adaptability of urban forests to a range of plausible future climates. Some cities are beginning to develop and use tools to optimize species selection at the site scale based on tree species characteristics (e.g., Baltimore, Maryland).³⁷⁶ This approach complements, and can be used in conjunction with, approaches 6.2 and 7.3, which employ native and introduced species, respectively.

EXAMPLE ADAPTATION TACTICS

- ▶ Include species that are less prone to ice and wind damage in forest restoration projects. Avoid planting trees susceptible to breakage in areas with high wind exposure.
- ▶ Select short-statured trees for planting under power lines to reduce susceptibility to infrastructure damage.

- ▶ Select species that are well-adapted to the soils in the area for restoration projects.³⁷⁷
- ▶ Select species that can provide multiple ecosystem services, such as carbon sequestration, stormwater mitigation, or pollinator habitat.
- ▶ Develop species selection decision-support tools to ensure “right tree, right place” planting.
- ▶ Add information to recommended plant lists to identify species that may provide human health, carbon mitigation, or adaptation benefits (appendix 3).
- ▶ Plant or retain fast-growing species to provide carbon sequestration, cover, shade, and food sources for wildlife during forest restoration efforts.
- ▶ Establish or promote species with higher carbon sequestration capacity near buildings and homes as part of plantings designed to reduce building energy use.
- ▶ Plant species with diverse timing of phenological events (e.g., flowering, fruiting, leaf out, and leaf drop) to provide necessary resources over a longer timeframe to forest-dependent wildlife species.³⁷⁸
- ▶ Work with nurseries, the local community, or conservation organizations to ensure the future availability of desired planting stock.



Urban forest planning can be integrated with parks planning to promote physical, mental, and social health. Courtesy photo by Guy Kramer, used with permission.

Approach 6.4: Maintain or enhance genetic diversity

Enhancing genetic diversity of urban trees can help reduce risks to a variety of stressors, including pests and pathogens. Street and park trees are often cultivars or grown from seeds that are sourced from a small number of parents to ensure that trees have predictable growth, survivorship, and tolerances. However, this can result in a lack of genetic diversity and may prove to be deleterious to long-term survivorship during changing conditions.³³³ For example, if all of the maple trees in an area are a single cultivar, they will probably react to climate change in a nearly identical manner. Increasing genetic diversity in the urban forest will ensure that some individuals are better equipped to withstand climate-induced stressors. Also, urban natural areas are heavily fragmented, which can cause reduced gene flow and lead to a decline in genetic diversity.³⁷⁹ This reduction can lead to increased vulnerability if the variants prove susceptible to prevalent climate change impacts.

EXAMPLE ADAPTATION TACTICS

Urban natural areas

- ▶ Select seeds and planting stock that originate from a variety of sites or a broader geographic area to increase genetic diversity in restoration and reclamation projects when local provenance is not a priority.
- ▶ Collect and plant seeds from individuals or populations that have survived pest outbreaks, dieback events, or extreme weather events as these plants may have resistant genotypes or have a greater tolerance to stressors.

Developed urban sites

- ▶ Plant a variety of both cultivars and wild genotypes for a given species. Use a greater number of plant varieties and sources to increase overall genetic diversity.
- ▶ Use cultivars of species that will be better suited for hotter and drier climates.
- ▶ Plant disease-resistant cultivars that had been previously lost due to pests or disease to re-establish a form of this species on the landscape, such as disease-resistant elm (*Ulmus* spp.) or chestnut (*Castanea* spp.).
- ▶ Work with growers to create new genotypes and cultivars of currently planted species that will be best adapted to anticipated climate changes.
- ▶ Use contract growing nurseries—or establish municipal nurseries—to increase supply of promising new cultivars.

Strategy 7. Alter urban ecosystems toward new and expected conditions

Urban areas often contain a mixture of planted species that come from diverse regions. Planted trees may be non-native taxa or species that are regionally native; that is, those from the same region but not currently growing at the particular location. Because these species

evolved in locations with different climates or site conditions, they may have very different tolerances to future climates and local conditions. Continuing to plant novel species can facilitate climate adaptation when species are carefully selected. Intentional consideration of the tolerances and traits of species will ideally help increase the capacity of the urban landscape to cope with change. Additionally, urban forests could help facilitate the migration of species that will be favored under future climate to new habitats at or beyond the edges of their current range.³⁸⁰

Approach 7.1: Favor or restore non-invasive species that are expected to be adapted to future conditions

Selecting native and non-invasive species already present in an urban area that are likely to do well under a range of future climate conditions can be a low-risk approach for transitioning to future climate conditions while ensuring continued or enhanced provisioning of health and mitigation benefits. Native species can provide important ecosystem services such as habitat for vertebrate and invertebrate species, and do not carry a risk of becoming invasive or lead to genetic mixing.³⁷⁰ However, there may be a limited set of native species that will be able to withstand future climate conditions and also thrive in urban environments.^{381, 382} Therefore, the use of “near-native” (e.g., from within 100 miles of the site) and non-native species that have been proven to tolerate urban conditions and lack invasive characteristics may be warranted in highly disturbed or developed areas.³⁸³ This approach complements efforts to remove problematic non-native invasive plant species (Approach 5.5).

EXAMPLE ADAPTATION TACTICS

Urban natural areas

- ▶ Plant native seedlings in restoration projects that are likely to do well, based on climate model projections and information about climatic tolerances.
- ▶ Plant “near-native” seedlings that do not have invasive properties.

Developed urban sites

- ▶ Promote native species that are near their northern range limit and future-adapted native species in tree planting lists and projects.
- ▶ Incorporate species and cultivars that are proven urban tolerant and not invasive when native species are insufficient in achieving biodiversity goals.³⁷⁷
- ▶ Select trees that are hardier to extreme storm and wind events and less likely to break up when pruned correctly, especially in wind-prone areas.
- ▶ Plant tree species that are less sensitive to flooding in low-lying areas that are expected to become wetter.
- ▶ Select species based on their physiological tolerance to drought.³⁸⁴

Approach 7.2: Establish or encourage new species mixes

Future conditions in urban areas are likely to become extreme, with higher peaks in temperature and moisture than might be seen in non-urban landscapes.³⁸⁵ However, considerable uncertainty exists in what future conditions will be, especially at the site level. Thus, encouraging new species mixtures when planting in urban forest communities could help these systems adapt, reducing tree mortality while maintaining ecological function¹² and carbon mitigation value³⁸⁶. This approach could also discourage invasion by exotic invasive species that reduce valued ecological function.³⁸⁷ Managers may need to prioritize diversity—in terms of species, as well as functional groups³⁸⁸⁻³⁹⁰ and genetic lineages—over historical species combinations to increase community resilience.^{391, 392}

EXAMPLE ADAPTATION TACTICS

Urban natural areas

- ▶ Plant a mixture of locally and regionally native species during ecosystem restoration to diversify beyond species that are adapted to current site conditions or that represent the historical plant community.
- ▶ Create heterogeneous conditions in canopy structure, ground layer, and hydrology that will allow a variety of species to become established.

Developed urban sites

- ▶ Include a diverse mix of locally and regionally native species in plantings, especially in plantings that are near natural areas.
- ▶ Intentionally design ecosystems to meet human-centered needs, such as green stormwater infrastructure or agro-ecosystems.³⁹³
- ▶ Establish mixes of tree functional groups (e.g., conifer, deciduous, and evergreen broadleaf) that can provide important ecosystem services across different seasons.^{388, 390}

Approach 7.3: Introduce species, genotypes, and cultivars that are expected to be adapted to future conditions

Urban foresters have moved species across states, continents, and even oceans for centuries, although only in recent decades has their intent included climate adaptation. This adaptation approach can include relatively low-risk actions, such as moving a species to slightly north of its current range. It also could include riskier actions, such as introducing a nonnative species from another continent. Although seeds and nursery stock from local sources may be the best adapted for an area currently, they may be maladapted to the changing climate in coming decades.³⁹⁴ Likewise, the unique climate in an urban center may already necessitate a different set of genetic material than for more pristine natural areas. Alternatively, seeds and nursery stock that are native to areas with a climate similar to the projected climate and

conditions of the targeted urban landscape may have higher survivorship than local seeds if other habitat factors (e.g., soils) are also suitable.³⁹⁵ Risks associated with introducing trees from distant sources include the potential for also introducing foreign pests and diseases, or the possibility that the introduced species may become invasive or hybridize with other local species.^{394, 396} These risks are reduced, but not eliminated, when a species is moved within its native range.³⁹⁵ Trials can help ensure that seedlings from distant areas will thrive in a new environment, but not harm ecosystem values, before large-scale plantings are undertaken.

EXAMPLE ADAPTATION TACTICS

Urban natural areas

- ▶ Incorporate regionally native species as well as non-native species into reclamation projects in degraded habitats to assess their viability and aggressiveness.
- ▶ Encourage southern species that become established in natural areas.
- ▶ Use mapping programs to track the origin of seed stocks and monitor their success to inform seed sourcing decisions in the future.
- ▶ Source seeds from a variety of areas to increase overall genetic diversity.

Developed urban sites:

- ▶ Introduce or increase regionally native or likely future native species in urban plant projects; for example, Kentucky coffeetree (*Gymnocladus dioica*) or tulip tree in upper midwestern cities.
- ▶ Plant non-native species from analogous climates.
- ▶ Use climate change projections to determine what region currently has a climate that is similar to the expected future climate in the target area, then source seeds from this area.
- ▶ Create a dialogue with nurseries and growers to ensure that seeds are being selected from healthy trees in areas that have a climate similar to the target area's expected climate.
- ▶ Plant and produce individuals collected or propagated from a variety of sites (including drought- and flood-prone areas) in consideration of the uncertainty of future conditions.
- ▶ Use cultivars of species that will be better suited for hotter and drier climates.
- ▶ Work with growers to create new genotypes and cultivars of currently planted species that will be best adapted to climate changes.
- ▶ Plant disease-resistant cultivars that had been previously lost due to pests or disease to re-establish a form of this species on the landscape, such as disease-resistant elm or chestnut.

Approach 7.4: Disfavor species that are distinctly maladapted

Urban areas are already experiencing especially rapid changes in climatic extremes, and some species at the edges of their natural ranges may more quickly become maladapted to

these conditions.³⁷⁸ With some urban environments already experiencing extreme climatic events, it is possible that an increasing number of species eventually may become more poorly adapted there.³⁹⁷ For example, species with a climate envelope that encompasses an urban area in large-scale future climate projections may not be able to tolerate conditions in more extreme urban microclimates.³⁹⁸ In some cases, species that are no longer adapted to an area may be removed from—or be considered for removal from—recommended planting lists.

EXAMPLE ADAPTATION TACTICS

- ▶ Use information from especially extreme urban sites, or areas with similar climatic extremes and fluctuations, to determine which native species are likely to decline.
- ▶ Do not replace, trees of selected species that are not drought-, heat-, or flood-tolerant. Instead, promote more tolerant native or near-native species.
- ▶ Protect healthy legacy trees that fail to regenerate, while de-emphasizing their importance in the mix of species being planted or regenerated.
- ▶ Remove species from recommended planting lists that are no longer able to tolerate current conditions.

Approach 7.5: Move at-risk species to more suitable locations

Climate may be changing more rapidly than some species can migrate, and the movement of plant and animal species may be restricted by land use or other impediments between



Research shows that exposure to the outdoors in all seasons is beneficial for human health. Courtesy photo by Eben Dente, American Forests, used with permission.

areas of suitable habitat.^{399, 400} Fragmentation of natural ecosystems and physical barriers in urban areas may make this even more challenging, especially for rare or threatened species.⁴⁰¹⁻⁴⁰³ At-risk species, such as rare or sensitive species that are constrained to a specific set of environmental conditions, are often incorporated into urban reserves (e.g., botanical gardens, arboretums, and municipal parks) and urban plantings (e.g., street trees and backyard gardens), and could, in some cases, be included in restoration or reclamation projects, such as urban river ways. Additional at-risk species—or species that provide habitat for at-risk species—could be added to urban planting lists where suitable habitat exists to increase their representation in the landscape. Assisted migration for species rescue, which focuses on avoiding extinction by physically relocating climate-threatened species, may be an option to consider in some cases. This approach is best implemented with great caution, incorporating due consideration of the uncertainties inherent in climate change, the sparse record of previous examples, and continued uncertainties of forest response.³⁹⁶ Providing new artificial habitat for at-risk species could help sustain them through increasing alterations in climate.

EXAMPLE ADAPTATION TACTICS

- ▶ Plant or seed a rare or threatened species that is at risk for extinction into a newly suitable habitat outside its current range.
- ▶ Assist the migration of wildlife around barriers by trapping and releasing into newly suitable locations.
- ▶ Plant heat-sensitive species on north slopes or in cold-air drainages.
- ▶ Collect seeds and other genetic material of at-risk species to contribute to a genetic repository.
- ▶ Plant individuals in a protected location that is expected to provide suitable habitat into the future.
- ▶ Use local conservatories, arboreta, botanical gardens, and parks to cultivate species once climate change challenges natural regeneration.
- ▶ Include at-risk species (or species that provide habitat for at-risk wildlife) in urban park, street, or campus plantings and in restoration or reclamation projects whenever possible or feasible (e.g., planting endangered southern species in parks).

Approach 7.6: Promptly revegetate and remediate sites after disturbance

Changes in climate will increase some large-scale disturbances, such as floods and windstorms. These disturbances can lead to catastrophic losses of trees and other vegetation in some areas. Proper management prior to disturbance events is critical, as an estimated 80 percent of tree damage during natural disasters can be attributed to pre-existing defects.⁴⁰⁴ The risk of damage can be minimized through proper selection of planting site and species, and appropriate maintenance, such as pruning.^{316, 405} Following a disturbance, swiftly

remediating and re-establishing vegetation on disturbed sites can help maintain the carbon sequestration capacity of urban forests. Additionally, planting vegetation may be needed to stabilize soils to prevent erosion and could help reduce the impacts of invasive species. In highly developed areas, replanting may be the only way to ensure the presence of species that provide desired ecosystem services, such as shade, aesthetics, or stormwater control. In natural areas, where a native seedbank may remain, replanting or managing natural regeneration may be beneficial to ensure the area has a species composition and structure that is aligned with management goals. In most cases, disturbances will not lead to complete loss of vegetation, but the remaining trees may have some degree of damage. After urgent severe hazards are handled, damage assessments and restoration planning are important to site recovery, including attention to amenity and legacy trees for their carbon and ecosystem benefits. The retention of these trees can help provide ecosystem services while new trees and other vegetation are becoming established.³⁶⁰ Revegetation may require appropriate site preparation. Some disturbances will leave behind woody debris or sedimentation that may need to be remediated prior to planting. Areas that are at risk to debris flows may require some amount of stabilization before restoration work can begin. In other cases, the soil organic layer may have been removed; therefore, mulch or other soil amendments will be needed. Disturbance events also provide an opportunity to strengthen public-private partnerships, as well as engaging and educating citizens and volunteers in greening efforts.¹⁸⁹


EXAMPLE ADAPTATION TACTICS

Urban natural areas

- ▶ Promptly mitigate and prepare the site for restoration to avoid further damage to soils, vegetation, and property.
- ▶ Amend the soil to restore organic matter when topsoil is lost from previous disturbance.
- ▶ Manage for natural regeneration of native species after disturbance through such practices as protecting seedlings and saplings from herbivory.
- ▶ Use the disturbance as an opportunity to eradicate or reduce the impact of invasive or undesirable species.
- ▶ Allow non-invasive species migrants that are not native to the site to remain as part of a novel mix of species, rather than eradicating these species.
- ▶ Prioritize planting of trees into recently disturbed areas.
- ▶ Include early-successional species in forest restoration projects to provide initial canopy cover.
- ▶ Retain nonhazardous dead and damaged trees for wildlife habitat and other ecosystem services.

Developed urban sites

- ▶ Replace damaged trees with those that increase taxonomic and functional diversity.
- ▶ Plant over several years to create a diverse age and size class structure in the urban canopy.
- ▶ Establish fast-growing or future-adapted tree species in locations heavily impacted by disturbance or human use.⁴⁰⁶
- ▶ Remove the remaining severely damaged trees that were not immediately removed during emergency response if they pose a hazard to people or property.
- ▶ Implement appropriate tree crown restoration pruning strategies for less severely damaged trees.
- ▶ If possible, stand fallen trees back up and use stakes or guy wires for support until the root system is structurally stable (usually applicable to newly planted trees or trees less than 4 inches in diameter).^{183, 316}
- ▶ Provide irrigation for stressed trees to encourage formation of new roots.^{183, 316}
- ▶ Develop a tree salvage plan to maximize the use of woody debris following a disturbance event.¹⁸⁹



From single trees to forested natural areas, nearby nature provides many human health benefits. Courtesy photo by Guy Kramer, used with permission.

Approach 7.7: Realign severely altered systems toward future conditions

Many urban areas may experience such significant alterations from human- and climate-induced disturbance that it will become difficult to restore systems that reflect native ecosystems of the past. The physical disturbance, altered abiotic conditions, and changed species composition arising from a major disturbance may cause an ecosystem to flip into a fundamentally different state.^{407, 408} Management of these systems may be realigned to create necessary changes in species composition and structure to better adapt forests to current and anticipated environments, rather than historical predisturbance conditions.^{12, 393, 409, 410} In urban natural areas, it may be beneficial to allow for natural regeneration while also engaging in human-assisted reforestation, with an emphasis on selecting species that are better adapted to weather extremes, such as high winds.⁴⁰⁷ In more developed areas, this could mean designing “novel ecosystems” that incorporate both natural and engineered elements and contain entirely new species compositions.⁴¹¹ In light of recent disasters, some cities are taking a hybrid approach by incorporating green and gray infrastructure in an effort to protect against future damages. This may include nature-based approaches, such as restoring marshes and beaches, and structural protection, such as levees, floodwalls, and pumps.⁴¹² Developing clear plans that establish processes for realigning significantly altered ecosystems before undertaking these actions will allow for more thoughtful discussion and better coordination with other adaptation responses.

EXAMPLE ADAPTATION TACTICS

- ▶ Allow community transition by planting future-adapted species within a site that is already declining or is expected to decline (e.g., converting a mesic maple forest to an oak savanna).
- ▶ Allow non-native, non-invasive species to remain as part of a novel mix of species, rather than eradicating them.
- ▶ Develop monitoring plans of natural regeneration to determine how the ecosystem is changing post-disturbance.⁴⁰⁷
- ▶ Assess the quality of the seedbank or regeneration post-disturbance, to determine the need for supplemental planting.
- ▶ Design “novel ecosystems” composed of a carefully selected mix of native and non-native species that align with projected future climates.
- ▶ Convert areas to green spaces that are expected to be vulnerable to future climate impacts, such as low-lying coastal areas.

Strategy 8: Promote mental and social health in response to climate change

This strategy addresses the personal and community level of interactions of people with nearby nature, and how to build capacities to cope in the face of climate change. The health and well-being of individuals, households, and neighborhoods can be negatively affected by climate change in numerous ways.^{413, 414} First, both long-term and abrupt changes in people's lives can bring on mental stress and anxiety. Also, the effects of climate change range from local to global in geography, and our awareness of such conditions can become personal and social stressors. Although many people might first think of trees as an ecological adaptation to address climate, urban forests and green spaces also serve as resources that help people and communities cope and recover.⁴¹⁵ Individuals and communities may receive a wide range of health benefits—from experiences of nearby nature,⁴¹⁶ including faster healing for hospital patients,⁴¹⁷ improved school performance,⁴¹⁸ reduced hypertension,⁴¹⁹ and increased lifespan.⁴²⁰ Climate change impacts people at different scales, from mental health of individuals⁴²¹ to how entire communities react and respond to adversity.⁴²² Community-based actions to care for and steward ecosystems can foster social connections, acknowledge cultural diversity, and help create places that are meaningful and healing. Urban forestry priorities and plans can be informed by understanding socio-psychological reactions to climate stressors and risks, as well as the healing effects of nature, so that actions can effectively promote mental and social health in any community.⁴²³

Approach 8.1: Provide nature experiences to ease stress and support mental function

Exposure to trees, forests, and urban greening ease the causes and symptoms of general mental health concerns and can be therapy for clinical conditions,^{421, 424} a critical consideration as climate change can affect mental health in multiple ways.⁴⁵ Stress disorders, anxiety, and depression are most directly influenced by acute natural disasters and disturbances as people cope with immediate and tragic changes in their lives. Additionally, chronic stress and behavioral change can arise from slowly progressing, long-term conditions, such as rising temperatures and changing precipitation. For instance, high temperatures are associated with an increased incidence of violence, aggression, and suicide, as well as higher rates of treatment for those with psychiatric conditions.⁴¹⁴ Further, living with uncertainty, coping with environmental threats, and experiencing changes in familiar places and personal routines are pathways to multiple health issues,¹¹¹ solastalgia (the loss of a sense of place), and changing attitudes about one's self and relationships with others.¹¹⁴ These effects are expressed across different population groups, depending on how directly exposed or vulnerable people are in their geographical conditions.⁴²⁵ Exposure to nature can reduce rumination (a propensity to dwell on negative thoughts).⁴²⁶⁻⁴²⁸ The presence of surrounding tree canopy and brief walks in forested areas can ease stress and

depression.⁴²⁹⁻⁴³³ Experiences in a more biodiverse outdoor settings may have provide greater benefits.^{434, 435} Nature can provide respite and restorative experiences that reduce stress, frustration, and anxiety, thus helping one to think more clearly and make better decisions in challenging situations.⁴³⁶⁻⁴³⁸ Finally, nature experiences, especially outdoor walks, can help individuals generate creative solutions to challenges in their lives, communities, and society.⁴³⁹⁻⁴⁴¹

EXAMPLE ADAPTATION TACTICS

- ▶ Plant and promote more tree canopy cover within a range of 100 meters of homes in residential areas, particularly for neighborhoods that have few trees.⁴⁴²⁻⁴⁴⁴
- ▶ Plant and conserve trees to augment environments that provide quality, short-term restorative nature experiences (e.g., forest views, forest bathing, green streetscapes).^{445, 446}
- ▶ Develop places for quiet, nature-filled walks of up to 2 miles.^{447, 448}
- ▶ Implement the principles of Attention Restoration Theory to enhance or create forest and nature spaces that offer opportunity for cognitive and attention recovery,⁴⁴⁹⁻⁴⁵² including the restorative elements of being away, compatibility, and soft fascination—meaning the outdoor elements that attract our attention without effort, such as water features, wildlife movement, or daily and seasonal changes.⁴⁵²
- ▶ Promote greater sensory, ecological, and vegetation diversity in nature-experience spaces to enhance mental health and healing.⁴⁵³⁻⁴⁵⁵
- ▶ Plant trees to buffer extreme noise sources and provide natural sounds, as chronic negative sound causes stress, reduces quality of sleep, and impacts social relationships.⁴⁵⁶⁻⁴⁵⁸
- ▶ Promote nature-based mental health and therapy for various vulnerable populations, such as locating tree projects within or near education, care, or treatment facilities for children⁴⁵⁹⁻⁴⁶³ or older adults.^{92, 464}
- ▶ Design and implement plantings for nature experiences that support creativity, a psychological resource people call on to address personal and community-level stresses and challenges (garden paths, water features, and outdoor seating).^{439, 465, 466}

Approach 8.2: Encourage community and social cohesion to support climate response

Climate change and disturbance events can shape and test social relationships within communities. Social cohesion is a shared resource that has practical implications for climate response. People in cities often gather in public spaces, including parks and green spaces, to relax, play, learn, and engage in civic events. It is around these multiple informal interactions that social community takes shape.⁴⁶⁷ Community also may form around a project or purpose as participants develop differing, yet harmonious, perceptions, skills, or shared

interests in a cooperative way to achieve desired outcomes. Social cohesion is the resulting interdependence among members of a community, experienced as shared values, loyalties, and cooperation. Social capital—meaning the networks of social relations characterized by trust and shared give-and-take for mutual benefit— can emerge. Social capital makes it possible to achieve things that cannot be accomplished solely by individuals, and makes achieving community-oriented goals possible.⁴⁶⁸ Individuals and groups within communities with strong social cohesion and social capital experience multiple health benefits.^{469, 470} Urban forestry programs and projects can foster the social interactions and dynamics that enable communities to adaptively respond to climate and improve health.^{415, 471} At one level, people often pursue intentional, purposeful contact with green space and gardens as a way to summon the capacities to recover during or after a crisis.⁴⁷² On a larger scale, sustained programs and activities, such as tree planting and stewardship, can catalyze social cohesion and social capital.⁴⁷³ The presence of urban green spaces, along with participation in urban forest activities, can encourage positive social interactions,⁴⁷¹ helping people develop the capacity to respond to ongoing risks (such as air quality) or extreme events (such as storms). In times of challenges and crises, local groups can activate their established networks and capacities to become “first responders” to address community and individual needs with local expertise. Social resilience depends upon connectedness and innovation, and daily nature-based experiences and interactions encourage locally relevant and resourceful response.^{474, 475}

EXAMPLE ADAPTATION TACTICS

- ▶ Engage residents and organizations in tree planting in ways that acknowledge cultural diversity within the community, encourage formation of social relationships and networks, build capacity for communities to address challenges or needs, and nurture place attachment.⁴⁷⁶⁻⁴⁷⁹
- ▶ Collaborate with residents and local organizations on data assessments that support urban forestry using community science, such as tree inventories, canopy assessment, tree health and risk assessments, and heat mapping or stewardship mapping.⁴⁸⁰⁻⁴⁸³
- ▶ Engage residents in tree-planting programs as stronger connections to nature support more pro-environmental behavior and is related to use of nature for psychological restoration.⁴⁸⁴
- ▶ Use trees to create spaces that serve as community gathering spaces used for celebration, festivals, feasts, and other community events, such as Arbor Day and Earth Day.
- ▶ Partner with local retail and commercial businesses, civic organizations, and faith centers to foster community-wide networks and capacity for action.
- ▶ Collaborate on community-based performance or outcome goals, such as wildlife habitat or sustainability, associated with trees and forest projects to motivate a program of shared community actions and purposes.

- ▶ Promote tree projects and green spaces as pathways to immigrant and cross-cultural interactions, understanding local heritage, and welcoming new arrivals to a community.^{485, 486} Examples include food forests, nature-based international holiday celebrations, and farm-to-table festivals.
- ▶ Engage children and youth in tree projects to nurture their sense of care and stewardship of trees and the environment, including motivating stipends, academic or service credit, or employment for sustained involvement.⁴⁸⁷
- ▶ Acknowledge and support trees and green spaces as living memorials and places for grieving following climate disasters or community change, as people often seek nature and gardens to heal from hardship.⁴⁸⁸⁻⁴⁹⁰
- ▶ Encourage the use of a green space or forests as a sacred space—a place generally shared for solace and contemplative reflection,⁴⁹¹ to connect with communities and places of worship and faith.
- ▶ Carefully plan and design green spaces to include creating public environments that discourage crime, instead encouraging perceptions of safety and security, including guiding principles of Crime Prevention Through Environmental Design.^{492, 493}

Strategy 9: Promote human health co-benefits in nature-based climate adaptation

This strategy encourages project decisions that support co-design, cultural ecosystem services, and the positive land-use aspects of urban forestry. Although forest and tree health are important to provide long-term climate benefits, additional attention to the placement and configuration of both conserved and newly planted forests can optimize human health. Considering human health benefits when prioritizing nature-based adaptation can significantly increase the influence and value proposition of the investment for human well-being and health promotion. Human health benefits are contingent on locating trees, forests, or parks near where people live or spend time. For example, if tree planting for carbon sequestration and asthma reduction are co-benefits goals, then a schoolyard adjacent to a high-speed road is a good choice for a planting project location. Health promotion includes many things: it refers to the full array of social determinants of human health, including food security, quality housing, and accessibility to nearby nature, and how these factors can support general wellness. It also refers to particular functional benefits, as nature experiences help to prevent specific illnesses, such as attention deficit disorder⁴⁹⁴ and depression.⁴³¹ Routine physical activity is one of the best ways to improve general health, reduce incidence of chronic disease, and promote healthy aging.⁴⁹⁵ As decisions are made about locations of urban forestry projects, plantings that improve streetscapes, trails, or pathways for walking would achieve a co-benefits approach. While response to a climate effect or threat may be the primary purpose of a tree or forest project, a modest amount of

additional planning and resources can extend goals to include a wide array of public health benefits.¹¹ A health co-benefits approach optimizes ecosystem services, offers opportunities to engage with nontraditional public health and community partners, and often provides a compelling case for sustaining urban forestry programs.⁴⁹⁶

Approach 9.1: Co-design large-scale green infrastructure and built systems to promote health

Nature-based and green infrastructure initiatives are increasingly used in urban climate adaptation. Engineered ecosystems integrate the underlying ecological functions of urban natural resources as remedies for multiple short- and long-term climate change effects. As cities progress from a historic emphasis on sanitary systems to more comprehensive and holistic sustainability systems, green infrastructure and urban greening can be imagined and implemented to provide co-benefits through co-design.⁴⁹⁷ As one example, green stormwater infrastructure incorporates trees with bioswales or rain gardens, and these natural precipitation management features can be co-designed as micro-parks in a neighborhood. Trees and novel ecosystems can be incorporated into built systems to directly influence healthy responses. This is an important contribution of climate adaptation to health equity because, in many places, health is determined more by where people live than any other factor, including genetics.⁴⁹⁸ While public health and wellness is appropriately addressed in city planning and policy, actual design implementation happens across multiple local government departments, such as engineering, transportation, and residential development. The urban forest can be incorporated into the functional details of urban infrastructure and systems at each level of implementation to satisfy public welfare goals at the broadest scale and enhance quality of life for neighborhoods and households. Including the community in co-design wherever possible can increase social cohesion and capital, while it ensures these systems are consistent with perceived social needs.

EXAMPLE ADAPTATION TACTICS

- ▶ Adopt a co-design for a co-benefits approach in urban forestry projects so that any tree planting for climate adaptation also optimizes systems or structures that promote public health outcomes,^{55, 499} particularly in communities having tree canopy or human health disparities.⁵⁰⁰
- ▶ Use tree projects to promote community walkability and physical activity,^{501, 502} according to public health guidelines,⁵⁰³ including routes linking homes to transit stations, schools, and workplaces.
- ▶ Plan, design, and activate facilities and programs that support and motivate physical activity, as more intentional engagement of nearby residents and users through activities such as community walking programs and yoga sessions facilitates and motivates more healthful behaviors and outcomes.⁵⁰⁴⁻⁵⁰⁶
- ▶ Use forest plantings, restoration, and conservation to enhance connectivity of existing

green spaces and align with transportation, utilities, riparian, and other urban corridors to expand functional connectivity of ecological, sociocultural, and active living systems to include walking/running paths, hiking trails, and bike routes.^{448, 507}

- ▶ Engage engineers, urban planners, and sustainability officers, along with other allied professionals, to pursue co-benefits project designs that integrate trees and expand open space in communities to promote health, such as green stormwater infrastructure, cloudburst management plans, and utilities projects.⁵⁵
- ▶ As cities implement multi-modal transportation projects, including Complete Streets,^{508, 509} provide shaded sidewalks and bike lanes so people are comfortable and safe when choosing to walk, bike, or access public transit.
- ▶ Plan projects to encourage longer nature encounters near residential areas and user facilities to provide optimal nature “dosage” opportunities of 15 to 60 minutes.^{445, 450, 510, 511}
- ▶ Promote greater biodiversity in projects—in terms of forest structure and vegetation composition—for greater psychological and physiological human health benefit.^{434, 512}

Approach 9.2: Provide micro-scale nature experiences to promote health and healing

People can gain health advantages from remarkably brief experiences while in nearby nature spaces of even limited size.^{513, 514} These small, yet frequent, micro-scale experiences may become increasingly important as climate changes challenge health.⁴⁵ Urban forest conservation, restoration, or planting policies and programs may be expansive in scope, but implementation may entail multiple small-scale projects. Small spaces, such as hospital gardens, are valuable resources for healing, therapy, and stress recovery.^{417, 515} Simply having window views of nature from homes, schools, and workplaces provide health benefits.⁵¹⁶ Tree plantings at the parcel level on private and public properties in cities can have important cumulative effects on both climate and human health, especially if adjacent to residential centers or facilities that serve vulnerable populations or imbedded within disadvantaged communities. In all these situations, additional thought can be given to how a tree project that is primarily oriented to biological or ecological functions might also be configured to promote human health. Modest additional planning can help ensure residents and users are engaged in the stewardship activities that provide healthful experiences and social cohesion benefits.

EXAMPLE ADAPTATION TACTICS

- ▶ Optimize access to nature settings of any size; plant and manage so that residents and visitors find such spaces to be accessible, secure, safe, and support experiences of fitness, adaptability, and delight.^{517, 518}
- ▶ Conserve large trees, as this has numerous benefits; multiple studies indicate human preferences for large trees over small, and both the health and ecological benefits quotient is many times greater for large trees compared to small ones.⁵¹⁹

- ▶ Optimize nature views from windows and interactions in and around the common spaces of civic service facilities, such as libraries, community centers, and government buildings, to model climate-friendly landscapes and create restorative opportunities as people engage in meetings, wait for service providers, take breaks, and arrive or depart from the building.⁵²⁰
- ▶ Optimize nature views and interactions in places where people work on focused tasks—including workplaces, schools, healthcare centers, and campuses—as brief, frequent nature encounters improve creativity and restore cognitive capacity for task attention and focus.^{460, 520-522}
- ▶ Collaborate with transportation and public works departments to promote shady corridors, create Green Streets or Complete Streets,⁵²³ and provide comfortable connecting spaces for “10 Minute Walks”—a nature and health metric being adopted by many local governments—to parks and green spaces.⁵²⁴
- ▶ Plan and reinforce access to forests, groves, and significant trees for forest therapy and bathing programs, as some of the most rigorous studies of nature and health indicate that this activity supports multiple health improvements, including stress reduction, improved immune function, and reductions in symptoms of diabetes and cardiovascular disease.^{525, 526}
- ▶ Select tree species to provide community food supplement,⁵²⁷ expressed as cultural food groves,⁵²⁸ food arboretums, or food forests,⁵²⁹ and include food trees in urban forest management plans.^{530, 531}
- ▶ Incorporate information about trees, forests, and urban greening and human health response in environmental and ecological interpretive materials, such as signs, brochures, and project outreach.
- ▶ Collaborate to support tree plantings in healthcare and clinical care settings, as nature provides therapeutic and healing benefits to patients^{417, 424, 532} and reduces stress in healthcare providers.⁵¹⁵
- ▶ Use tree planting and species selection to optimize protection from ultraviolet (UV) ray exposure by shading waiting areas or high-use outdoor spaces, such as sports fields, festival areas, and bus stops.^{135, 533, 534}

GLOSSARY

Adaptation: Adjustments, both planned and unplanned, in natural and human systems in response to climatic changes and subsequent effects. Ecosystem-based adaptation activities use a range of opportunities for sustainable management, conservation, and restoration.

Adaptive capacity: The general ability of institutions, systems, and individuals to moderate the risks of climate change, or to realize benefits, through changes in their characteristics or behavior. Adaptive capacity can be an inherent property, or it could have been developed because of previous policy, planning, or design decisions.

Adaptive management: A dynamic approach to forest management in which the effects of treatments and decisions are continually monitored and used, along with research results, to modify management on a continuing basis to ensure that objectives are being met.

Afforestation: Converting land to forest.

Age class: An interval into which the age range of trees is divided for classification or use.

Amenity trees: Trees with recreational, functional, environmental, ecological, social, health, or aesthetic value rather than for production purposes.

At-risk species: A species that has been determined to be vulnerable to observed or projected changes in climate or other stressors.

Biological legacy: Individual trees of a variety of species retained from harvest in order to maintain their presence on the landscape, provide a potential seed source for both species and genotypes that are expected to be better adapted to future conditions, and serve as future nurse logs for regeneration of some species.

Biomass: The mass of living organic matter (plant and animal) in an ecosystem. Biomass also refers to organic matter (living and dead) available on a renewable basis for use as a fuel; biomass includes trees and plants (both terrestrial and aquatic), agricultural crops and wastes, wood and wood wastes, forest and mill residues, animal wastes, livestock operation residues, and some municipal and industrial wastes.

Bioswale: A landscape feature designed to filter silt and pollution from surface runoff, consisting of a shallow drainage course with gently sloped sides and filled with vegetation.

Brownfield: Previously developed land potentially contaminated by a pollutant, hazardous substance, or other contaminant. The term may also be used to describe former industrial or commercial sites where its future expansion, reuse, or redevelopment is affected by known or perceived environmental pollution, such as soil contamination.

Carbon sequestration: The process of storing carbon in a carbon pool.

Climate: The statistical description of the weather in terms of the mean and variability of relevant quantities (usually temperature, precipitation, and wind) over periods of several decades (typically three decades). In a wider sense, the “climate” is the description of the state of the climate system.

Climate change: A change in the state of the climate that can be identified (e.g., by using statistical tests) by changes in the mean and/or the variability of its properties, and that persists for an extended period, typically decades or longer.

Climate refugia: Areas that are buffered from the impacts of climate change, and may allow for the persistence of physical, ecological, and socio-cultural resources. For example, variations in topography can create cooler areas in the landscape where sensitive species may be better able to persist into the future.

Disturbance: Stresses and destructive agents, such as invasive species, diseases, and fire; changes in climate and serious weather events, including hurricanes and ice storms; pollution of the air, water, and soil; real estate development of forest lands; and timber harvest.

Diversity: The variety and abundance of life forms, processes, functions, and structures of plants, animals, and other living organisms, including the relative complexity of species, communities, gene pools, and ecosystems at spatial scales that range from local through regional to global. There are commonly five levels of biodiversity: (1) genetic diversity—the genetic variation within a species; (2) species diversity—the variety of species in an area; (3) community or ecosystem diversity—the variety of communities or ecosystems in an area; (4) landscape diversity—the variety of ecosystems across a landscape; and (5) regional diversity—the variety of species, communities, ecosystems, or landscapes within a specific geographic region.

Ecological function: The sum of physical conditions (e.g., depth of water and soil type) and ecological processes (such as nutrient cycling and sediment movement) that make up an ecosystem and, ultimately, habitats on which species depend. A loss of ecological function is the removal or disruption of an ecological process that produces a certain physical condition or the loss of or damage to a physical condition.

Ecological processes: Processes fundamental to the functioning of a healthy and sustainable ecosystem, usually involving the transfer of energy and substances from one medium or trophic level to another (e.g., water flows and movement, nutrient cycling, sediment movement, and predator–prey relationships).

Ecosystem: A system of living organisms interacting with each other and their physical environment. The boundaries of an ecosystem are somewhat arbitrary, depending on the focus of interest or study. Thus, the extent of an ecosystem may range from very small spatial scales to, ultimately, the entire Earth.

Ecosystem resilience: The capacity of a system to absorb disturbance and reorganize while undergoing change yet retain essentially the same function, structure, identity, and feedbacks.

Ecosystem services: Ecological processes or functions having monetary or nonmonetary value to individuals or society at large. These are frequently classified as (1) supporting services, such as productivity or biodiversity maintenance; (2) provisioning services, such as food or fiber; (3) regulating services, such as climate regulation or carbon sequestration; and (4) cultural services, such as tourism or spiritual and aesthetic appreciation.

Equity: Equity is the principle of fairness in burden-sharing and is a basis for understanding how the impacts and responses to climate change, including costs and benefits, are distributed in and by society in equal ways. It is often aligned with ideas of equality, fairness, and justice, and applied with respect to equity in the responsibility for, and distribution of, climate impacts and policies across society, generations, and gender, as well as in the sense of who participates and controls the processes of decision-making.

Evapotranspiration: The process by which plants and soils release moisture into the atmosphere.

Fragmentation: A disruption of ecosystem or habitat connectivity, caused by human or natural disturbance, creating a mosaic of successional and developmental stages within or between forested tracts of varying patch size, isolation (distance between patches), and edge (cumulative length of patch edges).

Functional groups: A set of similar species that have similar traits. In trees, typical functional groups are conifers, deciduous broadleaf trees, and evergreen broadleaf trees.

Gene flow: Transfer of genetic material from one population to another, resulting in a changed composition of the gene pool of the receiving population.

Genetic diversity: Genetic variation within a species.

Gray infrastructure: Constructed structures that are often made of concrete such as dams, pipes, roads, seawalls, sewer systems, treatment facilities, storage basins, or stormwater systems.

Green infrastructure: A water management approach that mimics, restores, or protects the natural water cycle using measures such as landscaping, permeable pavement, plant or soil systems, or stormwater harvest and reuse.

Greenhouse gas: Gaseous constituents of the atmosphere, both natural and anthropogenic, that absorb and emit radiation at specific wavelengths within the spectrum of terrestrial radiation emitted by the Earth's surface, the atmosphere itself, and clouds. This property causes the greenhouse effect. Water vapor (H₂O), carbon dioxide (CO₂), nitrous oxide (N₂O), methane (CH₄), and ozone (O₃) are the primary greenhouse gases in the Earth's atmosphere. Moreover, there are a number of entirely human-made greenhouse gases present in the atmosphere, such as halocarbons and other chlorine- and bromine-containing substances

Health Impact Assessment: Widely used policy analysis tools that evaluate the potential health impacts of a project or policy and provide recommendations to increase positive health co-benefits and mitigate negative health impacts. These assessments include a broad definition of health; consideration of economic, social, or environmental health determinants; application to a broad set of policy sectors; involvement of affected stakeholders; explicit concerns about social justice; and a commitment to transparency.

Health outcomes: Changes in health that result from specific health care investments or interventions. Positive health outcomes include a general sense of well-being, as well as functioning well mentally, physically, and socially. Population-level health outcome metrics include life expectancy and self-reported level of health and function.

Hydrologic processes: The processes that occur as part of the water cycle, including evapotranspiration, condensation, infiltration, precipitation, and runoff. Earth's water circulates continuously between the oceans, atmosphere, and land.

Invasive species: A species that is not native to an area and whose introduction causes, or is likely to cause, economic or environmental harm or harm to human health.

Landscape fragmentation: A process in which larger areas of habitat or natural land cover are broken into smaller, more isolated patches. This generally occurs due to land-use change for agriculture, urban development, and road building.

Migration: The movement of genes, individuals, or species from one population or geographic location to another. Tree migration is largely influenced by dispersal ability, landscape connectivity, and climatological and other factors.

Mitigation (of climate change): A human intervention to reduce emissions or enhance the terrestrial sequestration of carbon dioxide (CO₂) and other greenhouse gases

Monitoring: The collection of information over time, generally on a sample basis, by measuring change in an indicator or variable to determine the effects of resource management treatments in the long term.

Native species: A species that has been present in a given place for a long enough period that it has adapted to the physical environment and developed complex relationships with other organisms in an ecological community

Novel ecosystem: A unique assemblage of species and environmental conditions resulting from human actions, both intentional and unintentional.

Nutrient cycling: The biological, geological, and chemical processes involved in the transfer and movement of energy and matter between living organisms and nonliving matter. Water, carbon, oxygen, and nitrogen are examples of major nutrient cycles.

Realignment: The process of tuning ecosystems or habitats to current and anticipated future conditions in such a way that they can respond adaptively to ongoing change.

Refugia: Locations and habitats that support populations of organisms that are limited to small fragments of their previous geographic range.

Regeneration: The vegetative (e.g., sprouting from clonal root structures and coppicing) or sexual regeneration of a plant species.

Reserve: Natural areas with little to no harvest activity, unless required to maintain the system, that do not exclude fire management or other natural disturbance processes.

Resistance: An adaptation option intended to improve the defenses of an ecosystem against anticipated changes or directly defend the forest against disturbance to maintain relatively unchanged conditions.

Restoration: The process of returning ecosystems or habitats to their original structure and species composition.

Riparian: Related to, living, or located in conjunction with a wetland, on the bank of a river or stream, or at the edge of a lake or tidewater. The riparian community significantly influences, and is significantly influenced by, the neighboring body of water.

Risk: The chance of something happening that will have an impact on objectives, often specified in terms of an event or circumstance and the consequences that may flow from it. Measured in terms of the consequences of an event and their likelihoods, risk may have a positive or negative impact.

Silvicultural practices: Management actions taken to guide the establishment, growth, composition, health, and quality of forests to meet the needs for timber, wildlife habitat, restoration, and recreation. Treatments may include actions such as thinning, prescribed burning, planting, pruning, and harvesting.

Social determinants of health: The social, economic, and physical conditions in the places people are born, live, work, educate, and play that affect a wide range of health and quality-of-life risks and outcomes. Health outcomes can be significantly impacted by social determinants—resources that enhance quality of life, such as affordable housing, access to nutritious food, quality education, a healthy environment, and access to nature/natural surroundings.

Soft fascination: Natural scenes can easily and almost effortlessly hold people's attention, while allowing room in the mind for other thoughts and reflection. These qualities play an important role in the restorative quality of nature. In contrast, "hard" fascination, such as watching television, fully occupies the mind, leaving little space for contemplation. Sunsets, clouds, and wind blowing through trees are examples of soft fascination.

Soil profile: The vertical cross-section of soil layers.

Stormwater infiltration: The ability of excess water during storm events to soak into the soil and increase groundwater recharge, rather than quickly running off the surface and into nearby water bodies.

Taxonomic diversity: A measure of biodiversity that includes the number and relative abundance of species in a given area.

Transition: An adaptation option intended to accommodate change and enable ecosystems to adaptively respond to changing and new conditions.

Upstream health planning: Considers the social, economic, and environmental origins of population-level health outcomes. Addressing the fundamental, root causes of poor health, rather than focusing merely on correcting symptoms, often requires governmental interventions and policy-change approaches. Housing, neighborhood conditions, and socioeconomic status are all upstream factors that play a fundamental causal role in health outcomes.

Urban resilience: The ability of an urban system—and all its constituent socioecological and sociotechnical networks across temporal and spatial scales—to maintain or rapidly return to desired functions in the face of a disturbance, adapt to change, and quickly transform systems that limit current or future adaptive capacity.

Volatile organic compounds (VOCs): Natural and human-made chemicals that are emitted as gases from various solids and liquids; plants developed a communication system to convey information based on VOCs and emit various compounds while communicating.

Vulnerability: The susceptibility of a system to the adverse effects of climate change. Vulnerability is a function of the magnitude of climatic change, the sensitivity of a system, and the ability of the system to adapt.

Wildland-urban interface (WUI): Any area where manmade improvements are built close to, or within, natural terrain and flammable vegetation; there may be a higher potential for wildland fire in these areas.

LITERATURE CITED

1. Intergovernmental Panel on Climate Change [IPCC]. 2018. **Global warming of 1.5°C. An IPCC special report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty.** In: Masson-Delmotte, V.; Zhai, P.; Pörtner, H.-O. [et al.], eds. Geneva, Switzerland: World Meteorological Organization. 32 p. <https://www.ipcc.ch/sr15/> (accessed February 28, 2021).
2. U.S. Global Change Research Program. 2018. **Impacts, risks, and adaptation in the United States: fourth national climate assessment, volume II.** In: Reidmiller, D.R.; Avery, C.W.; Easterling, D.R. [et al.], eds. Washington, DC: U.S. Global Change Research Program. 470 p. <https://doi.org/10.7930/NCA4.2018>.
3. Hunt, A.; Watkiss, P. 2011. **Climate change impacts and adaptation in cities: a review of the literature.** *Climatic Change*. 104(1): 13–49. <https://doi.org/10.1007/s10584-010-9975-6>.
4. Wilby, R.L. 2007. **A review of climate change impacts on the built environment.** *Built Environment*. 33(1): 31–45. <https://doi.org/10.2148/benv.33.1.31>.
5. Benedict, M.E.; McMahon, E.T. 2006. **Green infrastructure: linking landscapes and communities.** Washington, DC: Island Press. 299 p.
6. Rudnick, D.A.; Ryan, S.J.; Beier, P.; Cushman, S.A.; Dieffenbach, F., [et al.] 2012. **The role of landscape connectivity in planning and implementing conservation and restoration priorities.** *Issues in Ecology*. 16(Fall): 1–20.
7. Kaplan, R. 1984. **Impact of urban nature: a theoretical analysis.** *Urban Ecology*. 8(3): 189–197. [https://doi.org/10.1016/0304-4009\(84\)90034-2](https://doi.org/10.1016/0304-4009(84)90034-2).
8. Roman, L.A.; Conway, T.M.; Eisenman, T.S.; Koeser, A.K.; Ordóñez Barona, C. [et al.]. 2021. **Beyond ‘trees are good’: disservices, management costs, and tradeoffs in urban forestry.** *Ambio*. 50(3): 615–630. <https://doi.org/10.1007/s13280-020-01396-8>.
9. Jacobson, T.A.; Kler, J.S.; Hernke, M.T.; Braun, R.K.; Meyer, K.C. [et al.]. 2019. **Direct human health risks of increased atmospheric carbon dioxide.** *Nature Sustainability*. 2(8): 691–701. <https://doi.org/10.1038/s41893-019-0323-1>.
10. Shmool, J.L.; Yonas, M.A.; Newman, O.D.; Kubzansky, L.D.; Joseph, E. [et al.]. 2015. **Identifying perceived neighborhood stressors across diverse communities in New York City.** *American Journal of Community Psychology*. 56(1–2): 145–55. <https://doi.org/10.1007/s10464-015-9736-9>.
11. Wolf, K.L.; Lam, S.T.; McKeen, J.K.; Richardson, G.R.A.; van den Bosch, M. [et al.]. 2020. **Urban trees and human health: a scoping review.** *International Journal of Environmental Research and Public Health*. 17: 4371. <https://doi.org/10.3390/ijerph17124371>.
12. Millar, C.I.; Stephenson, N.L.; Stephens, S.L. 2007. **Climate change and forests of the future: managing in the face of uncertainty.** *Ecological Applications*. 17(8): 2145–2151. <https://doi.org/10.1890/06-1715.1>.
13. Peterson, D.L.; Millar, C.I.; Joyce, L.A.; Furniss, M.J.; Halofsky, J.E. [et al.]. 2011. **Responding to climate change on national forests: a guidebook for developing adaptation options.** Gen. Tech. Rep. PNW-GTR-855. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 109 p. <https://doi.org/10.2737/PNW-GTR-855>.
14. Chornesky, E.A.; Ackerly, D.D.; Beier, P.; Davis, F.W.; Flint, L.E. [et al.]. 2015. **Adapting California’s ecosystems to a changing climate.** *Bioscience*. 65(3): 247–262. <https://doi.org/10.1093/biosci/biu233>.

15. Heinz Center. 2008. **Strategies for managing the effects of climate change on wildlife and ecosystems.** Washington, DC: The H. John Heinz III Center for Science, Economics and the Environment. 43 p.
16. Heller, N.E.; Zavaleta, E.S. 2009. **Biodiversity management in the face of climate change: a review of 22 years of recommendations.** *Biological Conservation*. 142(1): 14–32. <https://doi.org/10.1016/j.biocon.2008.10.006>.
17. Ogden, A.; Innes, J. 2008. **Climate change adaptation and regional forest planning in Southern Yukon, Canada. Mitigation and Adaptation Strategies for Global Change.** 13(8): 833–861. <https://doi.org/10.1007/s11027-008-9144-7>.
18. Cross, M.S.; Zavaleta, E.S.; Bachelet, D.; Brooks, M.L.; Enquist, C.A. [et al.]. 2012. **The adaptation for conservation targets (ACT) framework: a tool for incorporating climate change into natural resource management.** *Environmental Management*. 50(3): 341–351. <https://doi.org/10.1007/s00267-012-9893-7>.
19. Morelli, T.L.; S., Y.; Smith, N.; Hennessey, M.B.; Millar, C.I. 2012. **Climate project screening tool: an aid for climate change adaptation.** Res. Pap. PSW-RP-263. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 29 p. <https://doi.org/10.2737/PSW-RP-263>.
20. Stein, B.A.; Glick, P.; Edelson, N.; Staudt, A. 2014. **Climate-smart conservation: putting adaptation principles into practice.** Washington, DC: National Wildlife Federation. 272 p. https://www.nwf.org/~media/PDFs/Global-Warming/2014/Climate-Smart-Conservation-Final_06-06-2014.pdf (accessed February 28, 2021).
21. Swanston, C.W.; Janowiak, M.K.; Brandt, L.A.; Butler, P.R.; Handler, S.D. [et al.]. 2016. **Forest adaptation resources: climate change tools and approaches for land managers, 2nd ed.** Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 161 p. <https://doi.org/10.2737/NRS-GTR-87-2>.
22. Griscom, B.W.; Adams, J.; Ellis, P.W.; Houghton, R.A.; Lomax, G. [et al.]. 2017. **Natural climate solutions. Proceedings of the National Academy of Sciences.** 114(44): 11645. <https://doi.org/10.1073/pnas.1710465114>.
23. Fargione, J.E.; Bassett, S.; Boucher, T.; Bridgham, S.D.; Conant, R.T. [et al.]. 2018. **Natural climate solutions for the United States. Science Advances.** 4(11): eaat1869. <https://doi.org/10.1126/sciadv.aat1869>.
24. McPherson, E.G.; Simpson, J.R. 2003. **Potential energy savings in buildings by an urban tree planting programme in California.** *Urban Forestry & Urban Greening*. 2(2): 73–86. <https://doi.org/10.1078/1618-8667-00025>.
25. Nowak, D.J.; Appleton, N.; Ellis, A.; Greenfield, E. 2017. **Residential building energy conservation and avoided power plant emissions by urban and community trees in the United States.** *Urban Forestry & Urban Greening*. 21: 158–165. <https://doi.org/10.1016/j.ufug.2016.12.004>.
26. Hockstad, L.; Hanel, L. 2018. **Inventory of U.S. greenhouse gas emissions and sinks.** In: Environmental System Science Data Infrastructure for a Virtual Ecosystem. <https://doi.org/10.15485/1464240>.
27. Ontl, T.A.; Janowiak, M.K.; Swanston, C.W.; Daley, J.; Handler, S. [et al.] 2020. **Forest management for carbon sequestration and climate adaptation.** *Journal of Forestry*. 118(1): 86–101. <https://doi.org/10.1093/jofore/fvz062>.
28. Klein, R.; Huq, S.; Denton, F.; Downing, T.E.; Richels, R.G. [et al.]. 2007. **Inter-relationships between adaptation and mitigation. Climate change 2007: Impacts, adaptation and vulnerability. Contribution of working group II to the fourth assessment report of the intergovernmental panel on climate change.** Cambridge, U.K.: Cambridge University Press: 745–777. <https://www.ipcc.ch/site/assets/uploads/2018/02/ar4-wg2-chapter18-1.pdf> (accessed February 28, 2021).
29. Vibrant Cities Lab. 2021. **Climate & health action guide.** <https://www.vibrantcitieslab.com/guides/climate-health-action-guide/> (accessed May 3, 2021).

30. Janowiak, M.K.; Dostie, D.D.; Wilson, M.A.; Kucera, M.J.; Skinner, R.H. [et al.]. 2016. **Adaptation resources for agriculture: responding to climate variability and change in the Midwest and Northeast**. Tech. Bull. 1944. Washington, DC: U.S. Department of Agriculture. 70 p. <https://www.nrs.fs.fed.us/pubs/54751> (accessed February 28, 2021).
31. O'Toole, D.; Brandt, L.A.; Janowiak, M.K.; Schmitt, K.M.; Shannon, P.D. [et al.]. 2019. **Climate change adaptation strategies and approaches for outdoor recreation**. Sustainability. 11(24): 7030. <https://doi.org/10.3390/su11247030>.
32. Shannon, P.D.; Swanston, C.W.; Janowiak, M.K.; Handler, S.D.; Schmitt, K.M. [et al.]. 2019. **Adaptation strategies and approaches for forested watersheds**. Climate Services. 13: 51–64. <https://doi.org/10.1016/j.cliser.2019.01.005>.
33. Staffen, A.; O'Connor, R.; Johnson, S.E.; Shannon, P.D.; Kearns, K. [et al.]. 2019. **Climate adaptation strategies and approaches for conservation and management of non-forested wetlands**. Report NFCH-4. Houghton, MI: U.S. Department of Agriculture, Northern Forests Climate Hub. 41 p. <https://www.climatehubs.usda.gov/hubs/northern-forests/topic/climate-adaptation-tools-wetland-conservation-and-management> (accessed February 28, 2021).
34. Tribal Adaptation Menu Team. 2019. **Dibaginjigaadeg anishinaabe ezhitwaad: a tribal climate adaptation menu**. Odanah, WI: Great Lakes Indian Fish and Wildlife Commission. 54 p. <https://www.fs.usda.gov/treesearch/pubs/58352> (accessed February 28, 2021).
35. City of Providence & Racial and Environmental Justice Committee of Providence. 2019. **The City of Providence's climate justice plan**. Providence, RI: The City of Providence. 88 p. <https://www.providenceri.gov/sustainability/climate-justice-action-plan-providence/> (accessed May 2, 2021).
36. Nesbitt, L.; Hotte, N.; Barron, S.; Cowan, J.; Sheppard, S.R.J. 2017. **The social and economic value of cultural ecosystem services provided by urban forests in North America: a review and suggestions for future research**. Urban Forestry and Urban Greening. (25): 103–111. <https://doi.org/10.1016/j.ufug.2017.05.005>.
37. Clark, J.R.; Matheny, N.P.; Cross, G.; Wake, V. 1997. **A model of urban forest sustainability**. Journal of Arboriculture. 23: 17–30.
38. da Silva, J.; Kernaghan, S.; Luque, A. 2012. **A systems approach to meeting the challenges of urban climate change**. International Journal of Urban Sustainable Development. 4: 125–145. <https://doi.org/10.1080/19463138.2012.718279>.
39. Tobey, M.B.; Binder, R.B.; Chang, S.; Yoshida, T.; Yamagata, Y. [et al.]. 2019. **Urban systems design: a conceptual framework for planning smart communities**. Smart Cities. 2: 522–537. <https://doi.org/10.3390/smartcities2040032>.
40. Chu, E.; Brown, A.; Michael, K.; Du, J.; Lwasa, S. [et al.]. 2019. **Unlocking the potential for transformative climate adaptation in cities**. Background Paper prepared for the Global Commission on Adaptation. Washington, DC: World Resources Institute, Ross Center. 76 p. <https://wrirosscities.org/research/publication/unlocking-potential-transformative-climate-adaptation-cities> (accessed February 28, 2021).
41. Nagendra, H.; Ostrom, E. 2012. **Polycentric governance of multifunctional forested landscapes**. International Journal of the Commons. 6(2): 104–133. <https://doi.org/10.18352/ijc.321>.
42. van den Bosch, M.; Ode Sang, Å. 2017. **Urban natural environments as nature-based solutions for improved public health – a systematic review of reviews**. Environmental Research. 158: 373–384. <https://doi.org/10.1016/j.envres.2017.05.040>.
43. Fox, H.; Cundill, G. 2018. **Towards increased community-engaged ecological restoration: a review of current practice and future directions**. Ecological Restoration. 36: 208–218. <https://doi.org/10.3368/er.36.3.208>.

44. National Academies of Sciences, Engineering and Medicine. 2018. **Protecting the health and well-being of communities in a changing climate: proceedings of a workshop.** Washington, DC: The National Academies Press. 154 p. <https://doi.org/10.17226/24846>.
45. Rudolph, L.; Harrison, C.; Buckley, L.; North, S. 2018. **Climate change, health and equity: a guide for local health departments.** Oakland, CA: Public Health Institute; and Washington, DC: American Public Health Association. 369 p. https://www.apha.org/-/media/files/pdf/topics/climate/climate_health_equity_ashx?la=en&hash=14D2F64530F1505EAE7AB16A-9F9827250EAD6C79 (accessed February 28, 2021).
46. Wolf, K.L. 2020. **Human health co-benefits of green stormwater infrastructure: residential green space opportunities in Seattle & King County.** Seattle, WA: University of Washington. 23 p.
47. Williams, D.R.; Costa, M.V.; Odunlami, A.O.; Mohammed, S.A. 2008. **Moving upstream: how interventions that address the social determinants of health can improve health and reduce disparities.** *Journal of Public Health Management and Practice*. 14(6): S8–S17. <https://doi.org/10.1097/01.PHH.0000338382.36695.42>.
48. Maller, C.; Townsend, M.; Pryor, A.; Brown, P.; St Leger, L. 2006. **Healthy nature healthy people: 'contact with nature' as an upstream health promotion intervention for populations.** *Health Promotion International*. 21: 45–54. <https://doi.org/10.1093/heapro/dai032>.
49. Ricklin, A.; Shah, S. 2017. **Metrics for planning healthy communities.** Chicago, IL: American Planning Association. 26 p. <https://planning-org-uploaded-media.s3.amazonaws.com/document/Metrics-Planning-Healthy-Communities.pdf> (accessed February 28, 2021).
50. Jennings, V.; Floyd, M.F.; Shanahan, D.; Coutts, C.; Sinykin, A. 2017. **Emerging issues in urban ecology: implications for research, social justice, human health, and well-being.** *Population and Environment*. 39: 69–86. <https://doi.org/10.1007/s11111-017-0276-0>.
51. Schwarz, K.; Fragkias, M.; Boone, C.G.; Zhou, W.; McHale, M. [et al.]. 2015. **Trees grow on money: urban tree canopy cover and environmental justice.** *PloS One*. 10: e0122051. <https://doi.org/10.1371/journal.pone.0122051>.
52. Hunter, R.F.; Cleland, C.; Cleary, A.; Droomers, M.; Wheeler, B.W. [et al.]. 2019. **Environmental, health, wellbeing, social and equity effects of urban green space interventions: a meta-narrative evidence synthesis.** *Environment International*. 130: 104923. <https://doi.org/10.1016/j.envint.2019.104923>.
53. Carmichael, C.E.; McDonough, M.H. 2019. **Community stories: explaining resistance to street tree-planting programs in Detroit, Michigan, USA.** *Society & Natural Resources*. 12: 588–605. <https://doi.org/10.1080/08941920.2018.1550229>.
54. Gonzalez, R.; James, T.; Ross, J. 2017. **Community-driven climate resilience planning: a framework.** [Place of publication unknown]: National Association of Climate Resilience Planners. 59 p. <https://www.adaptationclearinghouse.org/resources/community-driven-climate-resilience-planning-a-framework.html> (accessed February 28, 2021).
55. Willamette, P. 2018. **Green infrastructure & health guide.** Portland, OR: Willamette Partnership, Oregon Public Health Institute, Green Infrastructure Leadership Exchange. 66 p. http://willamettepartnership.org/wp-content/uploads/2018/07/Green-Infrastructure-final_7_12_18_sm.pdf (accessed February 28, 2021).
56. Gill, S.; Handley, J.; Ennos, A.; Pauleit, S. 2007. **Adapting cities for climate change: The role of the green infrastructure.** *Built Environment*. 33(1): 115–133. <https://doi.org/10.2148/benv.33.1.115>.
57. Norton, B.A.; Coutts, A.M.; Livesley, S.J.; Harris, R.J.; Hunter, A.M. [et al.]. 2015. **Planning for cooler cities: a framework to prioritise green infrastructure to mitigate high temperatures in urban landscapes.** *Landscape and Urban Planning*. (134): 127–138. <https://doi.org/10.1016/j.landurbplan.2014.10.018>.

58. Beckett-Camarata, J. 2003. **An examination of the relationship between the municipal strategic plan and the capital budget and its effect on financial performance.** *Journal of Public Budgeting, Accounting & Financial Management.* 15(1): 23–40. <https://doi.org/10.1108/JPBAFM-15-01-2003-B002>.
59. Michels, A.; De Graaf, L. 2010. **Examining citizen participation: local participatory policy making and democracy.** *Local Government Studies.* 36(4): 477–491. <https://doi.org/10.1080/03003930.2010.494101>.
60. American Forests. 2013. **Urban forest assessments resource guide.** <https://www.americanforests.org/wp-content/uploads/2013/06/Click-here-to-download-the-Urban-Forest-Assessments-Resource-Guide-as-a-PDF-3.pdf> (accessed March 10, 2021).
61. Garzón, C.; Cooley, H.; Heberger, M.; Moore, E.; Allen, L. [et al.]. 2012. **Community based climate adaptation planning: case study of Oakland, California. CEC-500-2012-038.** Oakland, CA: Pacific Institute. 86 p. <https://pacinst.org/publication/community-based-climate-adaptation-planning-oakland-case-study/> (accessed February 28, 2021).
62. Watkins, S.L.; Gerrish, E. 2018. **The relationship between urban forests and race: a meta-analysis.** *Journal of Environmental Management.* 209: 152–168. <https://doi.org/10.1016/j.jenvman.2017.12.021>.
63. U.S. Centers for Disease Control and Prevention. 2021. **Parks & trails health impact assessment toolkit.** https://www.cdc.gov/healthyplaces/parks_trails/ (accessed February 28, 2021).
64. Mate, K.; Wyatt, R. 2017. **Health equity must be a strategic priority.** *NEJM Catalyst.* <https://catalyst.nejm.org/doi/full/10.1056/CAT.17.0556> (accessed February 28, 2021).
65. Jennings, V.; Baptiste, A.K.; Osborne Jelks, N.T.; Skeete, R. 2017. **Urban green space and the pursuit of health equity in parts of the United States.** *International Journal of Environmental Research and Public Health.* 14: 1432. <https://doi.org/10.3390/ijerph14111432>.
66. California Urban Forest Advisory Committee. 2018. **CAL FIRE U&CF program strategic plan 2019-2024.** Sacramento, CA: The CAL FIRE Urban and Community Forestry Program. 2 p. https://www.fire.ca.gov/media/10515/cufac_strategic_plan_summary_1620.pdf (accessed May 1, 2021).
67. City of Ann Arbor. 2014. **Expanding urban tree canopy as a community health climate adaptation strategy.** Ann Arbor, MI: Michigan Department of Community Health, Division of Environmental Health. 72 p. https://www.michigan.gov/documents/mdhhs/A_Health_Impact_Assessment_of_the_Ann_Arbor_Urban_Forest_Community_Management_Plan_514426_7.pdf (accessed February 28, 2021).
68. Head, C. 2017. **Community forest storm mitigation planning: a guide for North Carolina communities.** Raleigh, NC: North Carolina Forest Service. 62 p. https://www.ncforestservice.gov/urban/pdf/CFSMP_W2017.pdf (accessed February 28, 2021).
69. Gould, S.; Dervin, K.; Watson, T. 2012. **Climate action for health: integrating public health into climate action planning.** Sacramento, CA: California Department of Public Health. 52 p. https://www.cdph.ca.gov/Programs/OHE/CDPH%20Document%20Library/CCHEP-General/CDPH-2012-Climate-Action-for-Health_accessible.pdf (accessed February 28, 2021).
70. Jennings, V.; Browning, M.H.E.M.; Rigolon, A. 2019. **Planning urban green spaces in their communities: intersectional approaches for health equity and sustainability.** In: Jennings, V.; Browning, M.H.E.M.; Rigolon, A., eds. *Urban green spaces.* Cham, Switzerland: Springer: 71–99. https://doi.org/10.1007/978-3-030-10469-6_5.
71. Local Government Commission. 2021. **Budgeting for climate resilience.** <https://www.lgc.org/newsletter/budgeting-for-climate-resilience/> (accessed May 3, 2021).

72. U.S. Indian Health Service. 2017. **Cultural highlights for Indian health professionals: a reference guide to American Indian and Alaska Native Cultures**. Rockville, MD: U.S. Department of Health and Human Services, Indian Health Service, Division of Health Professions Support. 2 p. <https://www.ihs.gov/careerops/indianhealthcareers/culturalinsights/> (accessed February 28, 2021).
73. Metrics for Healthy Communities. 2021. **Physical activity logic model**. St, Paul, MN: A.H. Wilder Foundation; and Minneapolis, MN: Federal Reserve Bank of Minneapolis. <http://metrics-forhealthycommunities.org/logic-models/physical-activity> (accessed February 21, 2021).
74. Houck, M. 2016. **Regional parks and green-spaces planning in Portland, Oregon: The politics and science of providing for nature in cities**. The George Wright Forum. 33: 295–307. (accessed February 28, 2021).
75. The Intertwine Alliance. 2019. **Strategic plan 2019**. https://www.theintertwine.org/sites/default/files/Intertwine-Alliance-Strategic-Plan-2019_0.pdf (accessed February 14, 2021).
76. Nardone, A.; Chiang, J.; Corburn, J. 2020. **Historic redlining and urban health today in U.S. cities**. Environmental Justice. 13(4): 109–119. <https://doi.org/10.1089/env.2020.0011>.
77. Nardone, A.; Rudolph, K.E.; Morello-Frosch, R.; Casey, J.A. 2021. **Redlines and greenspace: the relationship between historical redlining and 2010 greenspace across the United States**. Environmental Health Perspectives. 129(1): 017006. <https://doi.org/10.1289/EHP7495>.
78. Cutter, S.L.; Barnes, L.; Berry, M.; Burton, C.; Evans, E. [et al.]. 2008. **Community and regional resilience: perspectives from hazards, disasters, and emergency management**. Columbia, SC: University of South Carolina, Community & Regional Resilience Initiative. 19 p. <https://merid.org/case-study/community-and-regional-resilience-institute/> (accessed February 28, 2021).
79. Candib, L.M. 2007. **Obesity and diabetes in vulnerable populations: reflection on proximal and distal causes**. Annals of Family Medicine. 5: 547–556. <https://doi.org/10.1370/afm.754>.
80. Hattis, D.; Ogneva-Himmelberger, Y.; Ratick, S. 2012. **The spatial variability of heat-related mortality in Massachusetts**. Applied Geography. 33: 45–52. <https://doi.org/10.1016/j.apgeog.2011.07.008>.
81. Greene, C.S.; Robinson, P.J.; Millward, A.A. 2018. **Canopy of advantage: who benefits most from city trees?** Journal of Environmental Management. 208: 24–35. <https://doi.org/10.1016/j.jenvman.2017.12.015>.
82. Gerrish, E.; Watkins, S.L. 2018. **The relationship between urban forests and income: a meta-analysis**. Landscape and Urban Planning. 170: 293–308. <https://doi.org/10.1016/j.landurbplan.2017.09.005>.
83. Browning, M.H.E.M.; Rigolon, A. 2018. **Do income, race and ethnicity, and sprawl influence the greenspace-human health link in city-level analyses? Findings from 496 cities in the United States**. International Journal of Environmental Research and Public Health. 15: 1541. <https://doi.org/10.3390/ijerph15071541>.
84. Chuang, W.-C.; Boone, C.G.; Locke, D.H.; Grove, J.M.; Whitmer, A. [et al.]. 2017. **Tree canopy change and neighborhood stability: a comparative analysis of Washington, D.C. and Baltimore, MD**. Urban Forestry & Urban Greening. 27: 363–372. <https://doi.org/10.1016/j.ufug.2017.03.030>.
85. Shokry, G.; Connolly, J.J.T.; Anguelovski, I. 2020. **Understanding climate gentrification and shifting landscapes of protection and vulnerability in green resilient Philadelphia**. Urban Climate. 31: 100539. <https://doi.org/10.1016/j.uclim.2019.100539>.
86. Nathan, A. 2019. **Climate is the newest gentrifying force, and its effects are already re-shaping cities**. In: Harvard University Science in the News blog. <https://sitn.hms.harvard.edu/flash/2019/climate-newest-gentrifying-force-effects-already-re-shaping-cities> (accessed February 28, 2021).

87. Mummert, J. 2019. **Creating equitable access to parks and recreation: community engagement resource guide**. In: Open space [blog]. Ashburn, VA: National Recreation and Park Association. <https://www.nrpa.org/blog/creating-equitable-access-to-parks-and-recreation-nrpa-community-engagement-resource-guide/> (accessed February 28, 2021).
88. Manangan, A.P.; Uejio, C.K.; Saha, S.; Schramm, P.J.; Marinucci, G.D. [et al.]. 2016. **Assessing health vulnerability to climate change: a guide for health departments**. Atlanta, GA: U.S. Centers for Disease Control and Prevention. 23 p. <https://www.cdc.gov/climateandhealth/pubs/assessinghealthvulnerabilitytoclimatechange.pdf> (accessed February 28, 2021).
89. McNamara, K.E.; Buggy, L. 2017. **Community-based climate change adaptation: a review of academic literature**. *Local Environment*. 22: 443–460. <https://doi.org/10.1080/13549839.2016.1216954>.
90. U.S. Environmental Protection Agency. 2015. **Community-based adaptation to a changing climate**. EPA-230-F-15-001. Washington, DC: U.S. Environmental Protection Agency, Office of Policy. https://www.epa.gov/sites/production/files/2016-09/documents/community-based-adaptation_handout.pdf (accessed February 28, 2021).
91. Stevenson, K.T.; Moore, R.; Cosco, N.; Floyd, M.F.; Sullivan, W. [et al.]. 2020. **A national research agenda supporting green schoolyard development and equitable access to nature**. *Elementa: Science of the Anthropocene*. 8: 10. <https://doi.org/10.1525/elementa.406>.
92. Browning, M.H.E.M.; Lee, K.; Wolf, K.L. 2019. **Tree cover shows an inverse relationship with depressive symptoms in elderly residents living in U.S. nursing homes**. *Urban Forestry & Urban Greening*. 41: 23–32. <https://doi.org/10.1016/j.ufug.2019.03.002>.
93. Gehl Institute. 2018. **Inclusive healthy places - a guide to inclusion & health in public space: learning globally to transform locally**. New York, NY: Gehl Institute. 75 p. <https://gehlpeople.com/shopfront/inclusive-healthy-places/> (accessed February 28, 2021).
94. Johnston, M.; Shimada, L.D. 2004. **Urban forestry in a multicultural society**. *Journal of Arboriculture*. 30: 185–192.
95. Jack-Scott, E.; Piana, M.R.; Troxel, B.; Murphy-Dunning, C.; Ashton, M. 2013. **Stewardship success: how community group dynamics affect urban street tree survival and growth**. *Arboriculture & Urban Forestry*. 39(4): 189–196.
96. Cox, D.; Streeter, R. 2019. **The importance of place: neighborhood amenities as a source of social connection and trust**. Washington, DC: American Enterprise Institute. 12 p. <https://www.aei.org/research-products/report/the-importance-of-place-neighborhood-amenities-as-a-source-of-social-connection-and-trust/> (accessed February 28, 2021).
97. McMillen, H.L.; Campbell, L.K.; Svendsen, E.S.; Kealiikanakaoleohaililani, K.; Francisco, K.S. [et al.]. 2020. **Biocultural stewardship, indigenous and local ecological knowledge, and the urban crucible**. *Ecology and Society*. 25: 9. <https://doi.org/10.5751/ES-11386-250209>.
98. Tidball, K.G. 2014. **Seeing the forest for the trees: hybridity and social-ecological symbols, rituals and resilience in postdisaster contexts**. *Ecology and Society*. 19(4): 25. <https://doi.org/10.5751/es-06903-190425>.
99. Falconer, J. 2021. **The cultural and symbolic importance of forest resources**. Food and Agriculture Organization of the United Nations. <http://www.fao.org/3/t9450e/t9450e06.htm> (accessed May 3, 2021).
100. Wolch, J.R.; Byrne, J.; Newell, J.P. 2014. **Urban green space, public health, and environmental justice: the challenge of making cities 'just green enough'**. *Landscape and Urban Planning*. 125: 234–244. <https://doi.org/10.1016/j.landurbplan.2014.01.017>.
101. Rigolon, A.; Christensen, J. 2019. **Greening without gentrification: learning from parks-related anti-displacement strategies nationwide**. Los Angeles, CA: University of California, Los Angeles. 5 p. <https://www.ioes.ucla.edu/wp-content/uploads/Greening-without-Gentrification-report-2019.pdf> (accessed February 28, 2021).

102. Anguelovski, I. 2016. **From toxic sites to parks as (green) lulus? New challenges of inequity, privilege, gentrification, and exclusion for urban environmental justice.** *Journal of Planning Literature*. 31: 23–36. <https://doi.org/10.1177/0885412215610491>.
103. Anguelovski, I.; Cole, H.; Connolly, J.; Triguero-Mas, M. 2018. **Do green neighbourhoods promote urban health justice?** *The Lancet Public Health*. 3: e270. [https://doi.org/10.1016/S2468-2667\(18\)30096-3](https://doi.org/10.1016/S2468-2667(18)30096-3).
104. Cole, H.V.S.; Garcia Lamarca, M.; Connolly, J.J.T.; Anguelovski, I. 2017. **Are green cities healthy and equitable? Unpacking the relationship between health, green space and gentrification.** *Journal of Epidemiology and Community Health*. 71: 1118–1121. <https://doi.org/10.1136/jech-2017-209201>.
105. Enelow, N.; Schildt, C.; Abbott, B.; Sharma, S.; Cawley, A. [et al.]. 2017. **Jobs & equity in the urban forest.** Portland, OR: Ecotrust & PolicyLink. 110 p. <https://ecotrust.org/about-us/publications/#jobs-and-equity-in-the-urban-forest> (accessed February 28, 2021).
106. Falxa-Raymond, N.; Svendsen, E.; Campbell, L.K. 2013. **From job training to green jobs: a case study of a young adult employment program centered on environmental restoration in New York City, USA.** *Urban Forestry & Urban Greening*. 12: 287–295. <https://doi.org/10.1016/j.ufug.2013.04.003>.
107. VanWynsberghe, R. 2016. **Green jobs for the disadvantaged in British Columbia: the perspectives of non-governmental organisations and social entrepreneurs.** *Local Environment*. 21: 504–526. <https://doi.org/10.1080/13549839.2014.974151>.
108. Meltzer, R.; Ghorbani, P. 2017. **Does gentrification increase employment opportunities in low-income neighborhoods?** *Regional Science and Urban Economics*. 66: 52–73. <https://doi.org/10.1016/j.regsciurbeco.2017.06.002>.
109. Cole, H.V.S.; Triguero-Mas, M.; Connolly, J.J.T.; Anguelovski, I. 2019. **Determining the health benefits of green space: does gentrification matter?** *Health & Place*. 57: 1–11. <https://doi.org/10.1016/j.healthplace.2019.02.001>.
110. Anderson, S.; Hall, E.; Leahy, I. 2021. **Career pathways action guide.** Washington, DC: Vibrant Cities Lab. <https://www.vibrantcitieslab.com/guides/career-pathways-action-guide/> (accessed February 23, 2021).
111. Frumkin, H.; Hess, J.; Luber, G.; Malilay, J.; McGeehin, M. 2008. **Climate change: the public health response.** *American Journal of Public Health*. 98: 435–445. <https://doi.org/10.2105/AJPH.2007.119362>.
112. Dahl, K.; Licker, R.; Abatzoglou, J.T.; Delet-Barreto, J. 2019. **Increased frequency of and population exposure to extreme heat index days in the United States during the 21st century.** *Environmental Research Communications*. 1: 075002. <https://doi.org/10.1088/2515-7620/ab27cf>.
113. Hassan, N.A.; Hashim, Z.; Hashim, J.H. 2016. **Impact of climate change on air quality and public health in urban areas.** *Asia-Pacific Journal of Public Health*. 28(2 Suppl): 38s–48s. <https://doi.org/10.1177/1010539515592951>.
114. Galway, L.P.; Beery, T.; Jones-Casey, K.; Tasala, K. 2019. **Mapping the solastalgia literature: a scoping review study.** *International Journal of Environmental Research and Public Health*. 16. <https://doi.org/10.3390/ijerph16152662>.
115. Fagliano, J.A.; Diez Roux, A.V. 2018. **Climate change, urban health, and the promotion of health equity.** *PLoS Medicine*. 15: e1002621. <https://doi.org/10.1371/journal.pmed.1002621>.
116. U.S. Centers for Disease Control and Prevention. 2013. **Climate change and extreme heat events.** Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Environmental Health. 16 p. <https://www.cdc.gov/climateandhealth/pubs/climatechangeandextremeheatevents.pdf> (accessed February 28, 2021).
117. Bowler, D.E.; Buyung-Ali, L.; Knight, T.M.; Pullin, A.S. 2010. **Urban greening to cool towns and cities: a systematic review of the empirical evidence.** *Landscape and Urban Planning*. 97: 147–155. <https://doi.org/10.1016/j.landurbplan.2010.05.006>.

118. Morakinyo, T.E.; Kong, L.; Lau, K.K.-L.; Yuan, C.; Ng, E. 2017. **A study on the impact of shadow-cast and tree species on in-canyon and neighborhood's thermal comfort.** *Building and Environment*. 115: 1–17. <https://doi.org/10.1016/j.buildenv.2017.01.005>.
119. Kilbourne, E.M.; Choi, K.; Jones, S. 1982. **Risk factors for heatstroke.** A case-control study. *Journal of the American Medical Association*. 247: 3332–3336. <https://doi.org/10.1001/jama.1982.03320490030031>.
120. Graham, D.A.; Vanos, J.K.; Kenny, N.A.; Brown, R.D. 2016. **The relationship between neighbourhood tree canopy cover and heat-related ambulance calls during extreme heat events in Toronto, Canada.** *Urban Forestry & Urban Greening*. 20: 180–186. <https://doi.org/10.1016/j.ufug.2016.08.005>.
121. Oke, T.R. 1989. **The micrometeorology of the urban forest.** *Philosophical Transactions of the Royal Society of London B, Biological Sciences*. 324: 335–349. <https://doi.org/10.1098/rstb.1989.0051>.
122. Akbari, H.; Kurn, D.M.; Bretz, S.E.; Hanford, J.W. 1997. **Peak power and cooling energy savings of shade trees.** *Energy and Buildings*. 25: 139–148. [https://doi.org/10.1016/S0378-7788\(96\)01003-1](https://doi.org/10.1016/S0378-7788(96)01003-1).
123. Ziter, C.D.; Pedersen, E.J.; Kucharik, C.J.; Turner, M.G. 2019. **Scale-dependent interactions between tree canopy cover and impervious surfaces reduce daytime urban heat during summer.** *Proceedings of the National Academy of Sciences of the United States of America*. 116: 7575–7580. <https://doi.org/10.1073/pnas.1817561116>.
124. Lin, Y.-H.; Tsai, K.-T. 2017. **Screening of tree species for improving outdoor human thermal comfort in a Taiwanese city.** *Sustainability*. 9: 340. <https://doi.org/10.3390/su9030340>.
125. Zhao, Q.; Sailor, D.J.; Wentz, E.A. 2018. **Impact of tree locations and arrangements on outdoor microclimates and human thermal comfort in an urban residential environment.** *Urban Forestry & Urban Greening*. 32: 81–91. <https://doi.org/10.1016/j.ufug.2018.03.022>.
126. Yoshida, A.; Hisabayashi, T.; Kashihara, K.; Kinoshita, S.; Hashida, S. 2015. **Evaluation of effect of tree canopy on thermal environment, thermal sensation, and mental state.** *Urban Climate*. 14: 240–250. <https://doi.org/10.1016/j.uclim.2015.09.004>.
127. Jiao, M.; Zhou, W.; Zheng, Z.; Wang, J.; Qian, Y. 2017. **Patch size of trees affects its cooling effectiveness: A perspective from shading and transpiration processes.** *Agricultural and Forest Meteorology*. 247: 293–299. <https://doi.org/10.1016/j.agrformet.2017.08.013>.
128. Mohajerani, A.; Bakaric, J.; Jeffrey-Bailey, T. 2017. **The urban heat island effect, its causes, and mitigation, with reference to the thermal properties of asphalt concrete.** *Journal of Environmental Management*. 197: 522–538. <https://doi.org/10.1016/j.jenvman.2017.03.095>.
129. Aminipouri, M.; Knudby, A.J.; Krayenhoff, E.S.; Zickfeld, K.; Middel, A. 2019. **Modelling the impact of increased street tree cover on mean radiant temperature across Vancouver's local climate zones.** *Urban Forestry & Urban Greening*. 39: 9–17. <https://doi.org/10.1016/j.ufug.2019.01.016>.
130. Wang, Y.; Akbari, H. 2016. **The effects of street tree planting on urban heat island mitigation in Montreal.** *Sustainable Cities and Society*. 27: 122–128. <https://doi.org/10.1016/j.scs.2016.04.013>.
131. Rahman, M.A.; Moser, A.; Rötzer, T.; Pauleit, S. 2017. **Microclimatic differences and their influence on transpirational cooling of *Tilia cordata* in two contrasting street canyons in Munich, Germany.** *Agricultural and Forest Meteorology*. 232: 443–456. <https://doi.org/10.1016/j.agrformet.2016.10.006>.
132. Coutts, A.; Tapper, N. 2017. **Trees for a cool city: guidelines for optimised tree placement.** Melbourne, Australia: Cooperative Research Centre for Water Sensitive Cities. 24 p. <https://watersensitivecities.org.au/content/trees-cool-city-guidelines-optimised-tree-placement/> (accessed February 28, 2021).

133. Hondula, D.M.; Davis, R.E.; Leisten, M.J.; Saha, M.V.; Veazey, L.M. [et al.]. 2012. **Fine-scale spatial variability of heat-related mortality in Philadelphia County, USA, from 1983-2008: a case-series analysis.** *Environmental Health*. 11: 16. <https://doi.org/10.1186/1476-069X-11-16>.
134. Ng, E.; Chen, L.; Wang, Y.; Yuan, C. 2012. **A study on the cooling effects of greening in a high-density city: an experience from Hong Kong.** *Building and Environment*. 47: 256–271. <https://doi.org/10.1016/j.buildenv.2011.07.014>.
135. Olsen, H.; Kennedy, E.; Vanos, J. 2019. **Shade provision in public playgrounds for thermal safety and sun protection: A case study across 100 play spaces in the United States.** *Landscape and Urban Planning*. 189: 200–211. <https://doi.org/10.1016/j.landurbplan.2019.04.003>.
136. Park, J.; Kim, J.-H.; Lee, D.K.; Park, C.Y.; Jeong, S.G. 2017. **The influence of small green space type and structure at the street level on urban heat island mitigation.** *Urban Forestry & Urban Greening*. 21: 203–212. <https://doi.org/10.1016/j.ufug.2016.12.005>.
137. Voelkel, J.; Hellman, D.; Sakuma, R.; Shandas, V. 2018. **Assessing vulnerability to urban heat: a study of disproportionate heat exposure and access to refuge by socio-demographic status in Portland, Oregon.** *International Journal of Environmental Research and Public Health*. 15: 640. <https://doi.org/10.3390/ijerph15040640>.
138. Heaviside, C.; Macintyre, H.; Vardoulakis, S. 2017. **The urban heat island: implications for health in a changing environment.** *Current Environmental Health Reports*. 4: 296–305. <https://doi.org/10.1007/s40572-017-0150-3>.
139. Kenney, W.L.; Craighead, D.H.; Alexander, L.M. 2014. **Heat waves, aging, and human cardiovascular health.** *Medicine & Science in Sports and Exercise*. 46: 1891–1899. <https://doi.org/10.1249/MSS.0000000000000325>.
140. Mitchell, B.C.; Chakraborty, J. 2018. **Exploring the relationship between residential segregation and thermal inequity in 20 U.S. cities.** *Local Environment*. 23: 796–813. <https://doi.org/10.1080/13549839.2018.1474861>.
141. Fann, N.; Brennan, T.; Dolwick, P.; Gamble, J.L.; Ilacqua, V. [et al.]. 2016. **Ch. 3: Air quality impacts. The impacts of climate change on human health in the United States: a scientific assessment.** In: Crimmins, A.; Balbus, J.; Gamble, J.L. [et al.], eds. Washington, DC: U.S. Global Change Research Program: 69–98. <https://doi.org/10.7930/J0GQ6VP6>.
142. Calfapietra, C.; Fares, S.; Manes, F.; Morani, A.; Sgrigna, G. [et al.]. 2013. **Role of biogenic volatile organic compounds (BVOC) emitted by urban trees on ozone concentration in cities: a review.** *Environmental Pollution*. 183: 71–80. <https://doi.org/10.1016/j.envpol.2013.03.012>.
143. U.S. Environmental Protection Agency. 2009. **Integrated science assessment for particulate matter.** EPA/600/R-08/139F. Washington, DC: U.S. Environmental Protection Agency. 1071 p. http://ofmpub.epa.gov/eims/eimscomm.getfile?p_download_id=494959 (accessed February 28, 2021).
144. Fann, N.; Lamson, A.D.; Anenberg, S.C.; Wesson, K.; Risley, D. [et al.]. 2012. **Estimating the national public health burden associated with exposure to ambient PM_{2.5} and ozone.** *Risk Analysis*. 32(1): 81–95. <https://doi.org/10.1111/j.1539-6924.2011.01630.x>.
145. Zhang, Y.; Bielory, L.; Mi, Z.; Cai, T.; Robock, A. [et al.]. 2015. **Allergenic pollen season variations in the past two decades under changing climate in the United States.** *Global Change Biology*. 21: 1581–1589. <https://doi.org/10.1111/gcb.12755>.
146. Osborne, N.J.; Alcock, I.; Wheeler, B.W.; Hajat, S.; Sarran, C. [et al.]. 2017. **Pollen exposure and hospitalization due to asthma exacerbations: daily time series in a European city.** *International Journal of Biometeorology*. 61: 1837–1848. <https://doi.org/10.1007/s00484-017-1369-2>.
147. Yamazaki, S.; Shima, M.; Nakadate, T.; Ohara, T.; Omori, T. [et al.]. 2015. **Patterns of sensitization to inhalant allergens in Japanese lower-grade schoolchildren and related factors.** *International Archives of Allergy and Immunology*. 167: 253–263. <https://doi.org/10.1159/000439534>.

148. Kim, D.H.; Park, Y.-S.; Jang, H.J.; Kim, J.H.; Lim, D.H. 2016. **Prevalence and allergen of allergic rhinitis in Korean children.** *American Journal of Rhinology & Allergy*. 30: 72–78. <https://doi.org/10.2500/ajra.2013.27.4317>.
149. Ziska, L.; Knowlton, K.; Rogers, C.; Dalan, D.; Tierney, N. [et al.]. 2011. **Recent warming by latitude associated with increased length of ragweed pollen season in central North America.** *Proceedings of the National Academy of Sciences*. 108: 4248–4251. <https://doi.org/10.1073/pnas.1014107108>.
150. Makri, A.; Stilianakis, N.I. 2008. **Vulnerability to air pollution health effects.** *International Journal of Hygiene and Environmental Health*. 211(3–4): 326–336. <https://doi.org/10.1016/j.ijheh.2007.06.005>.
151. Donovan, G.H.; Gatzliolis, D.; Longley, I.; Douwes, J. 2018. **Vegetation diversity protects against childhood asthma: results from a large New Zealand birth cohort.** *Nature Plants*. 4: 358–364. <https://doi.org/10.1038/s41477-018-0151-8>.
152. Hartley, K.; Ryan, P.; Brokamp, C.; Gillespie, G.L. 2020. **Effect of greenness on asthma in children: a systematic review.** *Public Health Nursing*. 37: 453–460. <https://doi.org/10.1111/phn.12701>.
153. McDonald, R.; Kroeger, T.; Boucher, T.; Longzhu, W.; Salem, R. 2016. **Planting healthy air: a global analysis of the role of urban trees in addressing particulate matter pollution and extreme heat.** Arlington, VA: The Nature Conservancy. 130 p. https://www.nature.org/content/dam/tnc/nature/en/documents/20160825_PHA_Report_Final.pdf (accessed February 28, 2021).
154. Nowak, D.J.; Hirabayashi, S.; Bodine, A.; Greenfield, E. 2014. **Tree and forest effects on air quality and human health in the United States.** *Environmental Pollution*. 193: 119–129. <https://doi.org/10.1016/j.envpol.2014.05.028>.
155. Nowak, D.J.; Stein, S.M.; Randler, P.B.; Greenfield, E.J.; Comas, S.J. [et al.]. 2010. **Sustaining America's urban trees and forests: a forests on the edge report.** Gen. Tech. Rep. NRS-62. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 27 p. <https://doi.org/10.2737/NRS-GTR-62>.
156. Eisenman, T.S.; Churkina, G.; Jariwala, S.P.; Kumar, P.; Lovasi, G.S. [et al.]. 2019. **Urban trees, air quality, and asthma: an interdisciplinary review.** *Landscape and Urban Planning*. 187: 47–59. <https://doi.org/10.1016/j.landurbplan.2019.02.010>.
157. Kumar, P.; Druckman, A.; Gallagher, J.; Gatersleben, B.; Allison, S. [et al.] 2019. **The nexus between air pollution, green infrastructure and human health.** *Environment International*. 133: 105181. <https://doi.org/10.1016/j.envint.2019.105181>.
158. Nowak, D.J.; Civerolo, K.L.; Rao, S.T.; Sistla, G.; Luley, C.J. [et al.]. 2000. **A modeling study of the impact of urban trees on ozone.** *Atmospheric Environment*. 34: 1601–1613. <https://doi.org/10.2737/NRS-GTR-62>.
159. Calfapietra, C.; Morani, A.; Sgrigna, G.; Di Giovanni, S.; Muzzini, V. [et al.]. 2016. **Removal of ozone by urban and peri-urban forests: evidence from laboratory, field, and modeling approaches.** *Journal of Environmental Quality*. 45: 224–33. <https://doi.org/10.2134/jeq2015.01.0061>.
160. Han, D.; Shen, H.; Duan, W.; Chen, L. 2020. **A review on particulate matter removal capacity by urban forests at different scales.** *Urban Forestry & Urban Greening*. 48: 126565. <https://doi.org/10.1016/j.ufug.2019.126565>.
161. Yang, J.; Chang, Y.; Yan, P. 2015. **Ranking the suitability of common urban tree species for controlling PM_{2.5} pollution.** *Atmospheric Pollution Research*. 6: 267–277. <https://doi.org/10.5094/APR.2015.031>.
162. Beckett, K.P.; Freer-Smith, P.H.; Taylor, G. 1998. **Urban woodlands: their role in reducing the effects of particulate pollution.** *Environmental Pollution*. 99: 347–360. [https://doi.org/10.1016/S0269-7491\(98\)00016-5](https://doi.org/10.1016/S0269-7491(98)00016-5).

163. Beckett, K.P.; Freer Smith, P.H.; Taylor, G. 2000. **Effective tree species for local air quality management.** *Journal of Arboriculture*. 26: 12–19.
164. Beckett, K.P.; Freer-Smith, P.H.; Taylor, G. 2000. **Particulate pollution capture by urban trees: effect of species and windspeed.** *Global Change Biology*. 6: 995–1003. <https://doi.org/10.1046/j.1365-2486.2000.00376.x>.
165. Beckett, K.P.; Freer-Smith, P.H.; Taylor, G. 2000. **The capture of particulate pollution by trees at five contrasting urban sites.** *Arboricultural Journal*. 24: 209–230. <https://doi.org/10.1080/03071375.2000.9747273>.
166. Freer-Smith, P.H.; El-Khatib, A.A.; Taylor, G. 2004. **Capture of particulate pollution by trees: a comparison of species typical of semi-arid areas (*Ficus nitida* and *Eucalyptus globulus*) with European and North American species.** *Water, Air, & Soil Pollution*. 155: 173–187. <https://doi.org/10.1023/B:WATE.0000026521.99552.fd>.
167. Mikati, I.; Benson, A.F.; Luben, T.J.; Sacks, J.D.; Richmond-Bryant, J. 2018. **Disparities in distribution of particulate matter emission sources by race and poverty status.** *American Journal of Public Health*. 108: 480–485. <https://doi.org/10.2105/AJPH.2017.304297>.
168. Baldauf, R. 2016. **Recommendations for constructing roadside vegetation barriers to improve near-road air quality.** EPA600/R-16/072. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development. 13 p. <https://www.epa.gov/air-research/recommendations-constructing-roadside-vegetation-barriers-improve-near-road-air-quality> (accessed February 21, 2021).
169. Barwise, Y.; Kumar, P. 2020. **Designing vegetation barriers for urban air pollution abatement: a practical review for appropriate plant species selection.** *Climate and Atmospheric Science*. 3: 12. <https://doi.org/10.1038/s41612-020-0115-3>.
170. Vos, P.E.J.; Maiheu, B.; Vankerkom, J.; Janssen, S. 2013. **Improving local air quality in cities: to tree or not to tree?** *Environmental Pollution*. 183: 113–22. <https://doi.org/10.1016/j.envpol.2012.10.021>.
171. Pugh, T.A.M.; Mackenzie, A.R.; Whyatt, J.D.; Hewitt, C.N. 2012. **Effectiveness of green infrastructure for improvement of air quality in urban street canyons.** *Environmental Science & Technology*. 46: 7692–7699. <https://doi.org/10.1021/es300826w>.
172. Cariñanos, P.; Casares-Porcel, M. 2011. **Urban green zones and related pollen allergy: a review. Some guidelines for designing spaces with low allergy impact.** *Landscape and Urban Planning*. 101: 205–214. <https://doi.org/10.1016/j.landurbplan.2011.03.006>.
173. Cariñanos, P.; Casares-Porcel, M.; Díaz de la Guardia, C.; Aira, M.J.; Belmonte, J. [et al.]. 2017. **Assessing allergenicity in urban parks: a nature-based solution to reduce the impact on public health.** *Environmental Research*. 155: 219–227. <https://doi.org/10.1016/j.envres.2017.02.015>.
174. Kasprzyk, I.; Ćwik, A.; Kluska, K.; Wójcik, T.; Cariñanos, P. 2019. **Allergenic pollen concentrations in the air of urban parks in relation to their vegetation.** *Urban Forestry & Urban Greening*. 46: 126486. <https://doi.org/10.1016/j.ufug.2019.126486>.
175. Ren, Y.; Qu, Z.; Du, Y.; Xu, R.; Ma, D., et al. 2017. **Air quality and health effects of biogenic volatile organic compounds emissions from urban green spaces and the mitigation strategies.** *Environmental Pollution*. 230: 849–861. <https://doi.org/10.1016/j.envpol.2017.06.049>.
176. Churkina, G.; Grote, R.; Butler, T.M.; Lawrence, M. 2015. **Natural selection? Picking the right trees for urban greening.** *Environmental Science & Policy*. 47: 12–17. <https://doi.org/10.1016/j.envsci.2014.10.014>.
177. Fitzky, A.C.; Sandén, H.; Karl, T.; Fares, S.; Calfapietra, C. [et al.]. 2019. **The interplay between ozone and urban vegetation-bvoc emissions, ozone deposition, and tree ecophysiology.** *Frontiers in Forests and Global Change*. 2: 50. <https://doi.org/10.3389/ffgc.2019.00050>.

178. Lane, K.; Charles-Guzman, K.; Wheeler, K.; Abid, Z.; Graber, N. [et al.]. 2013. **Health effects of coastal storms and flooding in urban areas: a review and vulnerability assessment.** *Journal of Environmental and Public Health*. 2013: 913064. <https://doi.org/10.1155/2013/913064>.
179. Shoaf, K.; Rottman, S.J. 2000. **Public health impact of disasters.** *Australian Journal of Emergency Management*. 15: 58–63.
180. Giorgadze, T.; Maisuradze, I.; Japaridze, A.; Utiashvili, Z.; Abesadze, G. 2011. **Disasters and their consequences for public health.** *Georgian Medical News*. May (194): 59–63.
181. Ashley, W.S. 2007. **Spatial and temporal analysis of tornado fatalities in the United States: 1880–2005.** *Weather and Forecasting*. 22: 1214–1228. <https://doi.org/10.1175/2007WAF2007004.1>.
182. Smart Trees Pacific. 2013. **Urban forestry emergency operations planning guide for storm response.** Honolulu, HI: Smart Trees Pacific. <https://smarttreespacific.org/wp-content/uploads/UrbanForestry-EOP-Guide-printable-11-2013.pdf> (accessed February 28, 2021).
183. Gilman, E.; Duryea, M.; Kampf, E.; Delgado, A.; Escobedo, F. [et al.]. **Urban forest hurricane recovery program.** Gainesville, FL: University of Florida. <https://hort.ifas.ufl.edu/treesandhurricanes/> (accessed February 28, 2021).
184. Hauer, R.J.; Dawson, J.O.; Werner, L.P. 2006 **Trees and ice storms: the development of ice storm-resistant urban tree populations, 2nd edition.** Stevens Point, WI: College of Natural Resources, University of Wisconsin-Stevens Point ; and Urbana-Champaign, IL: Department of Natural Resources and Environmental Sciences and the Office of Continuing Education, University of Illinois at Urbana-Champaign. 20 p. <https://digitalcommons.unl.edu/usdafsfac-pub/295/> (accessed February 28, 2021).
185. Sutton-Grier, A.; Wowk, K.; Bamford, H. 2015. **Future of our coasts: the potential for natural and hybrid infrastructure to enhance the resilience of our coastal communities, economies and ecosystems.** *Environmental Science and Policy*. 51: 137–148. <https://doi.org/10.1016/j.envsci.2015.04.006>.
186. Cunniff, S.; Schwartz, A. 2015. **Performance of natural infrastructure and nature-based measures as coastal risk reduction features.** Washington DC: Environmental Defense Fund. 35 p. <http://www.biofund.org/mz/wp-content/uploads/2018/11/F1284.2015-Summary-Ni-Literature-Compilation-0-1.pdf> (accessed February 28, 2021).
187. Carmichael, C.; Danks, C.; Vatovec, C. 2019. **Green infrastructure solutions to health impacts of climate change: perspectives of affected residents in Detroit, Michigan, USA.** *Sustainability*. 11: 5688. <https://doi.org/10.3390/su11205688>.
188. Stanturf, J.A.; Goodrick, S.L.; Outcalt, K.W. 2007. **Disturbance and coastal forests: a strategic approach to forest management in hurricane impact zones.** *Forest Ecology and Management*. 250: 119–135. <https://doi.org/10.1016/j.foreco.2007.03.015>.
189. Burban, L.L.; Andresen, J. 1994. **Storms over the urban forest: planning, responding, and greening - a community guide to natural disaster relief.** Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northeastern Area State and Private Forestry. 94 p.
190. U.S. Centers for Disease Control and Prevention. 2020. **Neighborhood preparedness.** https://www.cdc.gov/cpr/infographics/00_docs/pfe-neighborhood-preparedness.pdf (accessed February 23, 2021).
191. Johnson, G.; Johnson, B. [N.d.]. **Storm damage to landscape trees.** St. Paul, MN: University of Minnesota Extension. <https://extension.umn.edu/planting-and-growing-guides/storm-damage-landscape-trees> (accessed February 28, 2021).

192. Duryea, M.; Kampf, E. 2007. **Wind and trees: lessons learned from hurricanes.** Gainesville, FL: University of Florida IFAS Extension. 17 p. <https://edis.ifas.ufl.edu/publication/fr173> (accessed February 28, 2021).
193. Parent, J.R.; Meyer, T.H.; Volin, J.C.; Fahey, R.T.; Witharana, C. 2019. **An analysis of enhanced tree trimming effectiveness on reducing power outages.** *Journal of Environmental Management*. 241: 397–406. <https://doi.org/10.1016/j.jenvman.2019.04.027>.
194. University of Connecticut. 2021. **Roadside forest management - stormwise.** <https://stormwise.uconn.edu/management/> (accessed February 23, 2021)
195. National Fire Protection Association. 2021. **Firewise USA.** <https://www.nfpa.org/Public-Education/Fire-causes-and-risks/Wildfire/Firewise-USA> (accessed February 23, 2021)
196. Chapman, S.; Watson, J.E.M.; Salazar, A.; Thatcher, M.; McAlpine, C.A. 2017. **The impact of urbanization and climate change on urban temperatures: a systematic review.** *Landscape Ecology*. 32(10): 1921–1935. <https://doi.org/10.1007/s10980-017-0561-4>.
197. Livesley, S.J.; McPherson, E.G.; Calfapietra, C. 2016. **The urban forest and ecosystem services: impacts on urban water, heat, and pollution cycles at the tree, street, and city scale.** *Journal of Environmental Quality*. (45): 119–124. <https://doi.org/10.2134/jeq2015.11.0567>.
198. Nowak, D.J.; Greenfield, E.J.; Hoehn, R.E.; Lapoint, E. 2013. **Carbon storage and sequestration by trees in urban and community areas of the United States.** *Environmental Pollution*. 178(Supplement C): 229–236. <https://doi.org/10.1016/j.envpol.2013.03.019>.
199. Nowak, D.J.; Greenfield, E.J. 2018. **Declining urban and community tree cover in the United States.** *Urban Forestry & Urban Greening*. (32): 32–55. <https://doi.org/10.1016/j.ufug.2018.03.006>.
200. State of Maryland, Sustainable Forestry Council. 2011. **No-net-loss of forest policy recommendations to the Maryland Department of Natural Resources.** https://dnr.maryland.gov/forests/Documents/sfc/SFC_NNL_110811.pdf (accessed February 28, 2021).
201. State of New Jersey. 2021. **No net loss - New Jersey state entities are required to replant trees when trees are removed during development projects involving one-half acre or more.** https://www.state.nj.us/dep/parksandforests/forest/community/No_Net_Loss.htm (accessed February 23, 2021).
202. American Public Works Association. 2021. **Urban forestry best management practices for public works managers - ordinances, regulations, & public policies.** <https://www2.apwa.net/Documents/About/CoopAgreements/UrbanForestry/UrbanForestry-3.pdf> (accessed February 23, 2021).
203. McKinney, M.L. 2002. **Urbanization, biodiversity, and conservation: the impacts of urbanization on native species are poorly studied, but educating a highly urbanized human population about these impacts can greatly improve species conservation in all ecosystems.** *Bioscience*. 52(10): 883–890. [https://doi.org/10.1641/0006-3568\(2002\)052\[0883:UBAC\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2002)052[0883:UBAC]2.0.CO;2).
204. Clark, C.; Ordóñez, C.; Livesley, S.J. 2020. **Private tree removal, public loss: valuing and enforcing existing tree protection mechanisms is the key to retaining urban trees on private land.** *Landscape and Urban Planning*. 203: 103899. <https://doi.org/10.1016/j.landurbplan.2020.103899>.
205. Elmendorf, W.; Gerhold, H.; Kuhns, L. 2021. **A guide to preserving trees in development projects.** University Park, PA: Penn State Extension. <https://extension.psu.edu/a-guide-to-preserving-trees-in-development-projects> (accessed May 5, 2021).
206. Hill, E.; Dorfman, J.; Kramer, E. 2010. **Evaluating the impact of government land use policies on tree canopy coverage.** *Land Use Policy*. 27: 407–414. <https://doi.org/10.1016/j.landusepol.2009.05.007>.

207. Salbitano, F.; Borelli, S.; Conigliaro, M.; Chen, Y. 2016. **Guidelines on urban and peri-urban forestry**. FAO Forestry Paper No. 178. Rome, Italy: Food and Agriculture Organization of the United Nations. 172 p. <http://www.fao.org/3/i6210e/i6210e.pdf> (accessed February 28, 2021).
208. Vogt, J.; Hauer, R.; Fischer, B. 2015. **The cost of not maintaining the urban forest**. *Arborist News*. 24: 12–17.
209. McPherson, E.; Nowak, D.; Heisler, G.; Grimmond, C.; Souch, C. [et al.]. 1994. **Chicago's urban forest ecosystem: results of the Chicago urban forest climate project**. Gen. Tech. Rep. NE-186. Radnor, PA: U. S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station. 201 p. <https://doi.org/10.2737/NE-GTR-186>.
210. Sadof, C.; Hughes, G.P.; Witte, A.R.; Peterson, D.; Ginzler, M.D. 2017. **Tools for staging and managing emerald ash borer in the urban forest**. *Arboriculture and Urban Forestry*. 43: 15–26. <https://doi.org/10.1603/ICE.2016.111745>.
211. Vannatta, A.; Hauer, R.; Schuettpelz, N. 2012. **Economic analysis of emerald ash borer (Coleoptera: Buprestidae) management options**. *Journal of Economic Entomology*. 105: 196–206. <https://doi.org/10.1603/EC11130>.
212. Roman, L.A.; Scatena, F.N. 2011. **Street tree survival rates: meta-analysis of previous studies and application to a field survey in Philadelphia, PA, USA**. *Urban Forestry & Urban Greening*. 10(4): 269–274. <https://doi.org/10.1016/j.ufug.2011.05.008>.
213. Costello, L. 2013. **Urban trees and water: an overview of studies on irrigation needs in the western United States and a discussion regarding future research**. *Arboriculture & Urban Forestry*. 39(3): 132–135.
214. Hammes, M.C.; Brandt, L.; Nagel, L.; Peterson, C.; Windmuller-Campione, M. [et al.]. 2020. **Adaptive silviculture for climate change in the Mississippi National River and Recreation Area, an urban national park in the Twin Cities area, Minnesota**. *Cities and the Environment*. 13(1): 11. <https://doi.org/10.15365/cate.2020.130111>.
215. Swanson, D.C. 2021. **Guidelines for planting trees and shrubs**. University of Massachusetts-Amherst: The Center for Agriculture, Food, and the Environment. <https://ag.umass.edu/landscape/fact-sheets/guidelines-for-planting-trees-shrubs> (accessed May 5, 2021)
216. Mulligan, J.; Rudee, A.; Lebling, K.; Levin, K.; Anderson, J. [et al.]. 2020. **Carbonshot: federal policy options for carbon removal in the United States**. Washington, DC: World Resources Institute. 96 p. <https://www.wri.org/research/carbonshot-federal-policy-options-carbon-removal-united-states> (accessed February 28, 2021).
217. Cook-Patton, S.C.; Gopalakrishna, T.; Daigneault, A.; Leavitt, S.M.; Platt, J. [et al.]. 2020. **Lower cost and more feasible options to restore forest cover in the contiguous United States for climate mitigation**. *One Earth*. 3: 739–752. <https://doi.org/10.1016/j.oneear.2020.11.013>.
218. Jo, H.-K.; McPherson, G.E. 1995. **Carbon storage and flux in urban residential greenspace**. *Journal of Environmental Management*. 45(2): 109–133. <https://doi.org/10.1006/jema.1995.0062>.
219. Pretzsch, H.; Biber, P.; Uhl, E.; Dahlhausen, J.; Schütze, G. [et al.]. 2017. **Climate change accelerates growth of urban trees in metropolises worldwide**. *Scientific Reports*. 7(1): 15403–15410. <https://doi.org/10.1038/s41598-017-14831-w>.
220. O'Brien, A.M.; Ettinger, A.K.; HilleRisLambers, J. 2012. **Conifer growth and reproduction in urban forest fragments: predictors of future responses to global change?** *Urban Ecosystems*. 15(4): 879–891. <https://doi.org/10.1007/s11252-012-0250-7>.
221. Hession, W.; Johnson, T.E.; Charles, D.; Hart, D.; Horwitz, R. [et al.] 2000. **Ecological benefits of riparian reforestation in urban watersheds: study design and preliminary results**. *Environmental Monitoring and Assessment*. 63: 211–222. <https://doi.org/10.1023/A:1006495805300>.

222. Siikamaki, J.; Wernstedt, K. 2008. **Turning brownfields into greenspaces: examining incentives and barriers to revitalization.** *Journal of Health Politics, Policy and Law.* 33(3): 559–593. <https://doi.org/10.1215/03616878-2008-008>.
223. De Sousa, C.A. 2003. **Turning brownfields into green space in the city of Toronto.** *Landscape and Urban Planning.* 62(4): 181–198. [https://doi.org/10.1016/S0169-2046\(02\)00149-4](https://doi.org/10.1016/S0169-2046(02)00149-4).
224. Scott, L.; Miller, K.A.; McCarthy, J.; Freer, M.; Darling, L. [et al.]. 2020. **Prioritization and planning to improve urban tree health in the Chicago region. Cities and the Environment.** 13(1): art. 6. <https://doi.org/10.15365/cate.2020.130106>.
225. Berland, A.; Shiflett, S.A.; Shuster, W.D.; Garmestani, A.S.; Goddard, H.C. [et al.]. 2017. **The role of trees in urban stormwater management. Landscape and Urban Planning.** (162): 167–177. <https://doi.org/10.1016/j.landurbplan.2017.02.017>.
226. Wenger, S.J.; Roy, A.H.; Jackson, C.R.; Bernhardt, E.S.; Carter, T.L. [et al.]. 2009. **Twenty-six key research questions in urban stream ecology: an assessment of the state of the science. Journal of the North American Benthological Society.** 28(4): 1080–1098. <https://doi.org/10.1899/08-186.1>.
227. University of Minnesota. 2021. **Gravel beds.** <https://trees.umn.edu/learn-more/gravel-beds> (accessed February 23, 2021).
228. Battin, J. 2004. **When good animals love bad habitats: ecological traps and the conservation of animal populations.** *Conservation Biology.* 18: 1482–1491. <https://doi.org/10.1111/j.1523-1739.2004.00417.x>.
229. Alvey, A.A. 2006. **Promoting and preserving biodiversity in the urban forest.** *Urban Forestry & Urban Greening.* 5(4): 195–201. <https://doi.org/10.1016/j.ufug.2006.09.003>.
230. Keppel, G.; Van Niel, K.P.; Wardell-Johnson, G.W.; Yates, C.J.; Byrne, M. [et al.]. 2012. **Refugia: identifying and understanding safe havens for biodiversity under climate change.** *Global Ecology and Biogeography.* 21(4): 393–404. <https://doi.org/10.1111/j.1466-8238.2011.00686.x>.
231. Anderson, M.G.; Clark, M.; Sheldon, A.O. 2012. **Resilient sites for terrestrial conservation in the Northeast and Mid-Atlantic region.** Boston, MA: The Nature Conservancy, Eastern Conservation Science. 168 p. https://eastern-division.s3.amazonaws.com/GP_Resilience/Great_Plains_Resilience.pdf (accessed February 21, 2021).
232. Yesilonis, I.; Pouyat, R. 2012. **Carbon stocks in urban forest remnants: Atlanta and Baltimore as case studies.** In: Lal, R.; Augustin, B., eds. *Carbon sequestration in urban ecosystems.* Dordrecht, The Netherlands: Springer: 103–120. https://doi.org/10.1007/978-94-007-2366-5_5.
233. Dybala, K.E.; Steger, K.; Walsh, R.G.; Smart, D.R.; Gardali, T. [et al.]. 2019. **Optimizing carbon storage and biodiversity co-benefits in reforested riparian zones.** *Journal of Applied Ecology.* 56(2): 343–353. <https://doi.org/10.1111/1365-2664.13272>.
234. Morelli, T.L.; Daly, C.; Dobrowski, S.Z.; Dulen, D.M.; Ebersole, J.L. [et al.]. 2016. **Managing climate change refugia for climate adaptation. PloS One.** 11(8): e0159909. <https://doi.org/10.1371/journal.pone.0159909>.
235. Ossola, A.; Locke, D.; Lin, B.; Minor, E. 2019. **Yards increase forest connectivity in urban landscapes. Landscape Ecology.** 34: 2935–2948. <https://doi.org/10.1007/s10980-019-00923-7>.
236. Groffman, P.M.; Bain, D.J.; Band, L.E.; Belt, K.T.; Brush, G.S. [et al.]. 2003. **Down by the riverside: urban riparian ecology.** 1: 315–321. [https://doi.org/10.1890/1540-9295\(2003\)001\[0315:DBTRUR\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2003)001[0315:DBTRUR]2.0.CO;2).
237. Fischenich, J.C. 2001. **Technologies for urban stream restoration and watershed management. EMRRP Bulletin No. 01-1. Vicksburg, MS: U.S. Army Corps of Engineers, Ecosystem Management and Restoration Research Program.** 11 p. <https://apps.dtic.mil/sti/citations/ADA391702> (accessed February 28, 2021).

238. Lipton, D.; Rubenstein, M.A.; Weiskopf, S.R.; Carter, S.; Peterson, J. [et al.]. 2018. **Ecosystems, ecosystem services, and biodiversity.** In: Reidmiller, D.R.; Avery C.W.; Easterling, D.R. [et al.], eds. *Impacts, risks, and adaptation in the United States: fourth national climate assessment, volume II.* Washington, DC: U.S. Global Change Research Program: 268–321. <https://doi.org/10.7930/NCA4.2018.CH7>.
239. Maxwell, K.; Julius, S.; Grambsch, A.; Kosmal, A.; Larson, L. [et al.]. 2018. **Built environment, urban systems, and cities.** In: Reidmiller, D.R.; C.W. Avery, C.W.; Easterling, D.R. [et al.], eds. *Impacts, risks, and adaptation in the United States: fourth national climate assessment, volume II.* Washington, DC: U.S. Global Change Research Program: 438–478. <https://doi.org/10.7930/NCA4.2018.CH11>.
240. Patterson, J.D.; Mader, D.L. 1982. **Soil compaction: causes and control in urban forest soils: a reference workbook, chapter 3.** Syracuse, NY: State University of New York, College of Environmental Science and Forestry.
241. Pouyat, R.; Szlávecz, K.; Yesilonis, I. 2019. **Human influences on urban soil development.** In: Pickett, S.; Cadenasso, M.; Grove, J. [et al.], eds. *Science for the sustainable city: empirical insights from the Baltimore school of urban ecology.* New Haven, CT: Yale University Press: 132–154. <https://doi.org/10.2307/j.ctvqc6gb7.13>.
242. Pouyat, R.; Szlávecz, K.; Yesilonis, I.D.; Groffman, P.; Schwarz, K. 2010. **Chemical, physical, and biological characteristics of urban soils.** In: Aitkenhead-Peterson, J.; Volder, A., eds. *Urban ecosystem ecology.* Agronomy Monograph 55. Madison, WI: American Society of Agronomy, Crop Science Society of America, Soil Science Society of America: 119–152.243.
243. Craul, P.J. 1985. **A description of urban soils and their desired characteristics.** *Journal of Arboriculture.* 11(11): 330–339.
244. Pouyat, R.; Yesilonis, I.; Nowak, D. 2006. **Carbon storage by urban soils in the United States.** *Journal of Environmental Quality.* 35: 1566–1575. <https://doi.org/10.2134/jeq2005.0215>.
245. Yesilonis, I.; Szlávecz, K.; Pouyat, R.; Whigham, D.; Xia, L. 2016. **Historical land use and stand age effects on forest soil properties in the Mid-Atlantic US.** *Forest Ecology and Management.* 370: 83–92. <https://doi.org/10.1016/j.foreco.2016.03.046>.
246. Pouyat, R.; Groffman, P.; Yesilonis, I.; Hernandez, L. 2002. **Soil carbon pools and fluxes in urban ecosystems.** *Environmental Pollution.* 116: S107–S118. [https://doi.org/10.1016/S0269-7491\(01\)00263-9](https://doi.org/10.1016/S0269-7491(01)00263-9).
247. Somerville, P.D.; Farrell, C.; May, P.B.; Livesley, S.J. 2020. **Biochar and compost equally improve urban soil physical and biological properties and tree growth, with no added benefit in combination.** *Science of the Total Environment.* 706: 135736. <https://doi.org/10.1016/j.scitotenv.2019.135736>.
248. Kumar, K.; Hundal, L.S. 2016. **Soil in the city: sustainably improving urban soils.** *Journal of Environmental Quality.* 45(1): 2–8. <https://doi.org/10.2134/jeq2015.11.0589>.
249. Rivenshield, A.; Bassuk, N.L. 2007. **Using organic amendments to decrease bulk density and increase macroporosity in compacted soils.** *Arboriculture and Urban Forestry.* 33(2): 140–146..
250. Heneghan, L.; Fatemi, F.; Umek, L.; Grady, K.; Fagen, K. [et al.]. 2006. **The invasive shrub european buckthorn (*Rhamnus cathartica*, L.) alters soil properties in midwestern US woodlands.** *Applied Soil Ecology.* 32(1): 142–148. <https://doi.org/10.1016/j.apsoil.2005.03.009>.
251. Asmelash, F.; Bekele, T.; Birhane, E. 2016. **The potential role of arbuscular mycorrhizal fungi in the restoration of degraded lands.** *Frontiers in Microbiology.* 7: art. 1095. <https://doi.org/10.3389/fmicb.2016.01095>.
252. Scharenbroch, B.C.; Carter, D.; Bialecki, M.; Fahey, R.; Scheberl, L. [et al.]. 2017. **A rapid urban site index for assessing the quality of street tree planting sites.** *Urban Forestry & Urban Greening.* 27: 279–286. <https://doi.org/10.1016/j.ufug.2017.08.017>.

253. Head, C.P.; Fisher, R.; O'Brien, M. 2001. **Best management practices for community trees: a technical guide to tree conservation in Athens-Clarke County, Georgia**. Athens, GA: Athens-Clarke County, GA Landscape Management Division Office. 134 p. https://www.accgov.com/DocumentCenter/View/280/Tree_BMPs?bidId= (accessed February 28, 2021).
254. Scharenbroch, B.C.; Catania, M. 2012. **Soil quality attributes as indicators of urban tree performance**. *Arboriculture & Urban Forestry*. 38(5): 214–228.
255. Watson, G.W.; Himelick, E.B. 2013. **The practical science of planting trees**. Champaign, IL: International Society of Arboriculture. 250 p.
256. Zhang, Y.; Qiao, L.; Chen, C.; Tian, L.; Zheng, X. 2020. **Effects of organic ground covers on soil moisture content of urban green spaces in semi-humid areas of China**. *Alexandria Engineering Journal*. 60(1): 251–259. <https://doi.org/10.1016/j.aej.2020.08.001>.
257. Layman, R.M.; Day, S.D.; Mitchell, D.K.; Chen, Y.; Harris, J.R. [et al.]. 2016. **Below ground matters: urban soil rehabilitation increases tree canopy and speeds establishment**. *Urban Forestry & Urban Greening*. 16: 25–35. <https://doi.org/10.1016/j.ufug.2016.01.004>.
258. Virginia Polytechnic Institute and State University. 2021. **Soil profile rebuilding**. <https://www.urbanforestry.frec.vt.edu/SRES/> (accessed May 5, 2021).
259. Nowak, D.J.; Bodine, A.R.; Hoehn, R.E.; Edgar, C.B.; Hartel, D.R. [et al.]. 2016. **Austin's urban forest, 2014**. *Resour. Bull. NRS-100*. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 55 p. <https://doi.org/10.2737/NRS-RB-100>.
260. Ellison, D.; Morris, C.E.; Locatelli, B.; Sheil, D.; Cohen, J. [et al.]. 2017. **Trees, forests and water: cool insights for a hot world**. *Global Environmental Change*. 43: 51–61. <https://doi.org/10.1016/j.gloenvcha.2017.01.002>.
261. Barling, R.D.; Moore, I.D. 1994. **Role of buffer strips in management of waterway pollution: a review**. *Environmental Management*. 18(4): 543–558. <https://doi.org/10.1007/bf02400858>.
262. Castelle, A.J.; Johnson, A.W.; Conolly, C. 1994. **Wetland and stream buffer size requirements-a review**. *Journal of Environmental Quality*. 23(5): 878–882. <https://doi.org/10.2134/jeq1994.00472425002300050004x>.
263. Haase, D. 2017. **Urban wetlands and riparian forests as a nature-based solution for climate change adaptation in cities and their surroundings**. In: Kabisch, N.; Korn, H.; Stadler, J. [et al.], eds. *Nature-based solutions to climate change adaptation in urban areas: linkages between science, policy and practice*. Cham, Switzerland: Springer International Publishing: 111–121. https://doi.org/10.1007/978-3-319-56091-5_7.
264. Palta, M.M.; Grimm, N.B.; Groffman, P.M. 2017. **"Accidental" urban wetlands: ecosystem functions in unexpected places**. *Frontiers in Ecology and the Environment*. 15(5): 248–256. <https://doi.org/10.1002/fee.1494>.
265. Richardson, D.M.; Holmes, P.M.; Esler, K.J.; Galatowitsch, S.M.; Stromberg, J.C. [et al.]. 2007. **Riparian vegetation: degradation, alien plant invasions, and restoration prospects**. *Diversity and Distributions*. 13(1): 126–139. <https://doi.org/10.1111/j.1366-9516.2006.00314.x>.
266. Groffman, P.M.; Crawford, M.K. 2003. **Denitrification potential in urban riparian zones**. *Journal of Environmental Quality*. 32(3): 1144–1149. <https://doi.org/10.2134/jeq2003.1144>.
267. Olive, N.D.; Marion, J.L. 2009. **The influence of use-related, environmental, and managerial factors on soil loss from recreational trails**. *Journal of Environmental Management*. 90(3): 1483–1493. <https://doi.org/10.1016/j.jenvman.2008.10.004>.
268. Marion, J.L.; Leung, Y.-F. 2004. **Environmentally sustainable trail management**. In: Buckley, R.C., ed. *Environmental impact of tourism*. Cambridge, MA: CABI Publishing: 229–244.
269. Fulé, P.Z. 2008. **Does it make sense to restore wildland fire in changing climate?** *Restoration Ecology*. 16: 526–531. <https://doi.org/10.1111/j.1526-100X.2008.00489.x>.

270. Hamilton, M.; Fischer, A.P.; Guikema, S.D.; Keppel-Aleks, G. 2018. **Behavioral adaptation to climate change in wildfire-prone forests.** WIREs Climate Change. 9: e553. <https://doi.org/10.1002/wcc.553>.
271. Gobster, P.H.; Floress, K.; Westphal, L.M.; Watkins, C.A.; Vining, J. [et al.]. 2016. **Resident and user support for urban natural areas restoration practices.** *Biological Conservation*. 203: 216–225. <https://doi.org/10.1016/j.biocon.2016.09.025>.
272. Kaval, P.; Loomis, J.; Seidl, A. 2007. **Willingness-to-pay for prescribed fire in the Colorado (USA) wildland urban interface.** *Forest Policy and Economics*. 9(8): 928–937. <https://doi.org/10.1016/j.forpol.2006.08.003>.
273. Blanchard, B.; Ryan, R.L. 2007. **Managing the wildland–urban interface in the Northeast: perceptions of fire risk and hazard reduction strategies.** *Northern Journal of Applied Forestry*. 24(3): 203–208. <https://doi.org/10.1093/njaf/24.3.203>.
274. Heuberger, K.A.; Putz, F.E. 2003. **Fire in the suburbs: ecological impacts of prescribed fire in small remnants of longleaf pine (*Pinus palustris*) sandhill.** *Restoration Ecology*. 11(1): 72–81. <https://doi.org/10.1046/j.1526-100X.2003.09982.x>.
275. Walker, R.B.; Coop, J.D.; Parks, S.A.; Trader, L. 2018. **Fire regimes approaching historic norms reduce wildfire-facilitated conversion from forest to non-forest.** *Ecosphere*. 9(4): e02182. <https://doi.org/10.1002/ecs2.2182>.
276. Tian, D.; Wang, Y.; Bergin, M.; Hu, Y.; Liu, Y. [et al.]. 2008. **Air quality impacts from prescribed forest fires under different management practices.** *Environmental Science & Technology*. 42(8): 2767–2772. <https://doi.org/10.1021/es0711213>.
277. Long, J.W.; Tarnay, L.W.; North, M.P. 2018. **Aligning smoke management with ecological and public health goals.** *Journal of Forestry*. 116(1): 76–86. <https://doi.org/10.5849/jof.16-042>.
278. Blades, J.J.; Shook, S.R.; Hall, T.E. 2014. **Smoke management of wildland and prescribed fire: understanding public preferences and trade-offs.** *Canadian Journal of Forest Research*. 44(11): 1344–1355. <https://doi.org/10.1139/cjfr-2014-0110>.
279. Menges, E.; Smith, S.; McAllister, S. 2012. **Post-fire cohort demography of Florida rosemary, a foundation shrub in Florida scrub.** In: 97th ESA Annual Convention 2012; 2012 August 5-10; Portland, OR. Washington, DC: Ecological Society of America. <https://eco.confex.com/eco/2012/webprogram/Paper34649.html> (accessed May 10, 2021).
280. McIver, J.D.; Stephens, S.L.; Agee, J.K.; Barbour, J.; Boerner, R.E.J. [et al.]. 2013. **Ecological effects of alternative fuel-reduction treatments: highlights of the national fire and fire surrogate study (FFS).** *International Journal of Wildland Fire*. 22(1): 63. <https://doi.org/10.1071/wf11130>.
281. Steenberg, J.W.N.; Duinker, P.N.; Nitoslawski, S.A. 2019. **Ecosystem-based management revisited: updating the concepts for urban forests.** *Landscape and Urban Planning*. 186: 24–35. <https://doi.org/10.1016/j.landurbplan.2019.02.006>.
282. Winter, G.J.; Vogt, C.; Fried, J.S. 2002. **Fuel treatments at the wildland-urban interface: common concerns in diverse regions.** *Journal of Forestry*. 100(1): 15–21. <https://doi.org/10.1093/jof/100.1.15>.
283. Schoennagel, T.; Nelson, C.R.; Theobald, D.M.; Carnwath, G.C.; Chapman, T.B. 2009. **Implementation of national fire plan treatments near the wildland–urban interface in the western United States.** *Proceedings of the National Academy of Sciences*. 106(26): 10706–10711. <https://doi.org/10.1073/pnas.0900991106>.
284. Schwilk, D.W.; Keeley, J.E.; Knapp, E.E.; McIver, J.; Bailey, J.D. [et al.]. 2009. **The national fire and fire surrogate study: effects of fuel reduction methods on forest vegetation structure and fuels.** *Ecological Applications*. 19(2): 285–304. <https://doi.org/10.1890/07-1747.1>.

285. Brandt, L.A.; Lewis, A.D.; Scott, L.; Darling, L.; Fahey, R.T. [et al.]. 2017. **Chicago wilderness region urban forest vulnerability assessment and synthesis: a report from the urban forestry climate change response framework Chicago wilderness pilot project**. Gen. Tech. Rep. NRS-168. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 142 p. <https://doi.org/10.2737/NRS-GTR-168>.
286. Liu, Z.; Wimberly, M.C.; Lamsal, A.; Sohl, T.L.; Hawbaker, T.J. 2015. **Climate change and wildfire risk in an expanding wildland-urban interface: a case study from the Colorado Front Range corridor**. *Landscape Ecology*. 30(10): 1943–1957. <https://doi.org/10.1007/s10980-015-0222-4>.
287. Hallegatte, S.; Green, C.; Nicholls, R.J.; Corfee-Morlot, J. 2013. **Future flood losses in major coastal cities**. *Nature Climate Change*. 3(9): 802–806. <https://doi.org/10.1038/nclimate1979>.
288. Jactel, H.; Koricheva, J.; Castagneyrol, B. 2019. **Responses of forest insect pests to climate change: not so simple**. *Current Opinion in Insect Science*. 35: 103–108. <https://doi.org/10.1016/j.cois.2019.07.010>.
289. Ramsfield, T.D.; Bentz, B.J.; Faccoli, M.; Jactel, H.; Brockerhoff, E.G. 2016. **Forest health in a changing world: effects of globalization and climate change on forest insect and pathogen impacts**. *Forestry*. 89(3): 245–252. <https://doi.org/10.1093/forestry/cpw018>.
290. Weed, A.S.; Ayres, M.P.; Hicke, J.A. 2013. **Consequences of climate change for biotic disturbances in North American forests**. *Ecological Monographs*. 83(4): 441–470. <https://doi.org/10.1890/13-0160.1>.
291. Hellmann, J.J.; Byers, J.E.; Bierwagen, B.G.; Dukes, J.S. 2008. **Five potential consequences of climate change for invasive species**. *Conservation Biology*. 22(3): 534–543. <https://doi.org/10.1111/j.1523-1739.2008.00951.x>.
292. Wallingford, P.D.; Morelli, T.L.; Allen, J.M.; Beaury, E.M.; Blumenthal, D.M. [et al.]. 2020. **Adjusting the lens of invasion biology to focus on the impacts of climate-driven range shifts**. *Nature Climate Change*. 10(5): 398–405. <https://doi.org/10.1038/s41558-020-0768-2>.
293. Dale, A.G.; Frank, S.D. 2017. **Warming and drought combine to increase pest insect fitness on urban trees**. *PloS One*. 12(3): e0173844. <https://doi.org/10.1371/journal.pone.0173844>.
294. Dietz, M. 2007. **Low impact development practices: a review of current research and recommendation for future directions**. *Water, Air, and Soil Pollution*. 186: 351–363. <https://doi.org/10.1007/s11270-007-9484-z>.
295. Pyke, C.; Warren, M.; Johnson, T.; LaGro, J.J.; Scharfenberg, J. [et al.]. 2011. **Assessment of low-impact development for managing storm water with changing precipitation due to climate change**. *Landscape and Urban Planning*. 103: 166–173. <https://doi.org/10.1016/j.landurbplan.2011.07.006>.
296. Ahiablame, L.; Engel, B.; Venort, T. 2012. **Improving water supply systems for domestic uses in urban Togo: the case of a suburb in Lomé**. *Water*. 4(1): 123–134. <https://doi.org/10.3390/w4010123>.
297. Kirshen, P.; Caputo, L.; Vogel Richard, M.; Mathisen, P.; Rosner, A. [et al.]. 2015. **Adapting urban infrastructure to climate change: a drainage case study**. *Journal of Water Resources Planning and Management*. 141(4): 04014064. [https://doi.org/10.1061/\(ASCE\)WR.1943-5452.0000443](https://doi.org/10.1061/(ASCE)WR.1943-5452.0000443).
298. Augustyn, F.; Chou, B. 2013. **Getting climate smart: a water preparedness guide for state action**. Washington, DC: American Rivers; Natural Resources Defense Council. 151 p. https://www.americanrivers.org/wp-content/uploads/2017/03/GettingSmart_ClimateChange.pdf (accessed February 21, 2021).
299. Hermansen, A. 2020. **Urban forest systems and green stormwater infrastructure**. SFS-1146. Washington, DC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 23 p. <https://www.fs.usda.gov/treesearch/pubs/61010> (accessed May 10, 2021).

300. Kavehei, E.; Jenkins, G.; Adame, F.; Lemckert, C. 2018. **Carbon sequestration potential for mitigating the carbon footprint of green stormwater infrastructure.** *Renewable and Sustainable Energy Reviews.* 94: 1179–1191. <https://doi.org/10.1016/j.rser.2018.07.002>.
301. Xiao, Q.; McPherson, E.G. 2016. **Surface water storage capacity of twenty tree species in Davis, California.** *Journal of Environmental Quality.* 45(1): 188–198. <https://doi.org/10.2134/jeq2015.02.0092>.
302. Shuster, W.D.; Gehring, R.; Gerken, J. 2007. **Prospects for enhanced groundwater recharge via infiltration of urban storm water runoff: a case study.** *Journal of Soil and Water Conservation.* 62(3): 129–137.
303. Smiley, E.T.; Calfee, L.; Fraedrich, B.R.; Smiley, E.J. 2006. **Comparison of structural and noncompacted soils for trees surrounded by pavement.** *Arboriculture and Urban Forestry.* 32(4): 164.
304. Deeproot. 2021. **Deeproot Silva Cell - integrated tree, soil and stormwater system.** https://www.deeproot.com/silvapdfs/resources/supporting/silva_cell_brochure.pdf (accessed February 23, 2021).
305. Wuebbles, D.J.; Fahey, D.W.; Hibbard, K.A.; DeAngelo, B.; Doherty, S. [et al.]. 2017. **Executive summary. In: Climate science special report: fourth national climate assessment.** Washington, DC: U.S. Global Change Research Program: 12–34. <https://doi.org/10.7930/J0DJ5CTG>.
306. Simpson, P.; van Bossuyt, R. 1996. **Tree-caused electric outages.** *Journal of Arboriculture.* 22: 117–121.
307. Schmidlin, T.W. 2009. **Human fatalities from wind-related tree failures in the United States, 1995–2007.** *Natural Hazards.* 50: 131–142. <https://doi.org/10.1007/s11069-008-9314-7>.
308. Koeser, A.K.; Smiley, E.T. 2017. **Impact of assessor on tree risk assessment ratings and prescribed mitigation measures.** *Urban Forestry & Urban Greening.* 24: 109–115. <https://doi.org/10.1016/j.ufug.2017.03.027>.
309. Duryea, M.L.; Kampf, E.; Littell, R.C. 2007. **Hurricanes and the urban forest: I. Effects on southeastern United States coastal plain tree species.** *Arboriculture and Urban Forestry.* 33(2): 83.
310. Francis, J.K. 2000. **Comparison of hurricane damage to several species of urban trees in San Juan, Puerto Rico.** *Journal of Arboriculture.* 26: 189–196.
311. King, D.A. 1986. **Tree form, height growth, and susceptibility to wind damage in *Acer saccharum*.** *Ecology.* 67: 980–990. <https://doi.org/10.2307/1939821>.
312. Putz, F.E.; P.D. Coley; K. Lu; A. Montalvo; Aiello, A. 1983. **Uprooting and snapping of trees: structural determinants and ecological consequences.** *Canadian Journal of Forest Research.* 13: 1011–1020. <https://doi.org/10.1139/x83-133>.
313. Miller, R.W. 1997. **Urban forestry: planning and managing urban greenspaces.** Upper Saddle River, NJ: Prentice Hall. 502 p.
314. Purcell, L. 2012. **Tree risk management.** FNR-475-W. West Lafayette, IN: Purdue University Extension. 3 p. <https://mdc.itap.purdue.edu/item.asp?itemID=20925> (accessed February 28, 2021).
315. Gardiner, B.A.; Stacey, G.R. 1996. **Designing forest edges to improve wind stability.** Tech. Pap. 16. Edinburgh, Scotland: U.K. Forestry Commission. 18 p. <https://www.forestresearch.gov.uk/research/archive-designing-forest-edges-to-improve-wind-stability/> (accessed February 28, 2021).
316. Gilman, E.F.; Partin, T. 2007. **Urban design for a wind resistant urban forest.** ENH 1056. Gainesville, FL: University of Florida IFAS extension. 8 p. <https://hort.ifas.ufl.edu/woody/documents/EP309.pdf> (accessed February 28, 2021).
317. Tree Care Industry Association. 2021. **Trends in tree care incidents (2015).** http://www.tcia.org/tcia/Blog_Items/2016/Trends_in_Tree_Care_Incidents_2015.aspx (accessed February 28, 2021).

318. Purcell, L. 2015. **Trees and storms**. FNR-FAQ-12-W. West Lafayette, IN: Purdue University Extension. 5 p. https://www.edustore.purdue.edu/item.asp?Item_Number=FNR-FAQ-12-W (accessed February 28, 2021).
319. Galvin, M.; Grove, J.M.; Hines, S.; Marshall, L. 2020. **The urban wood workbook: a framework for the Baltimore wood project**. NRS-INF-37-20. Madison, WI: U.S. Department of Agriculture, Forest Service, Northern Research Station. 36 p. https://www.fs.fed.us/research/docs/urban/nrs_inf_37_20_final_2020-05-05.pdf (accessed February 28, 2021).
320. Syphard, A.D.; Radeloff, V.C.; Keeley, J.E.; Hawbaker, T.J.; Clayton, M.K. [et al.]. 2007. **Human influence on California fire regimes**. *Ecological Applications*. 17(5): 1388–1402. <https://doi.org/10.1890/06-1128.1>.
321. Abatzoglou, J.T.; Kolden, C.A. 2013. **Relationships between climate and macroscale area burned in the western United States**. *International Journal of Wildland Fire*. 22(7): 1003–1020. <https://doi.org/10.1071/WF13019>.
322. Schoennagel, T.; Balch, J.K.; Brenkert-Smith, H.; Dennison, P.E.; Harvey, B.J. [et al.]. 2017. **Adapt to more wildfire in western North American forests as climate changes**. *Proceedings of the National Academy of Sciences*. 114(18): 4582–4590. <https://doi.org/10.1073/pnas.1617464114>.
323. Ager, A.A.; Palaiologou, P.; Evers, C.R.; Day, M.A.; Ringo, C. [et al.]. 2019. **Wildfire exposure to the wildland urban interface in the western US**. *Applied Geography*. 111: 102059. <https://doi.org/10.1016/j.apgeog.2019.102059>.
324. Halofsky, J.E.; Peterson, D.L.; Harvey, B.J. 2020. **Changing wildfire, changing forests: the effects of climate change on fire regimes and vegetation in the Pacific Northwest, USA**. *Fire Ecology*. 16(1): 4. <https://doi.org/10.1186/s42408-019-0062-8>.
325. Radeloff, V.C.; Helmers, D.P.; Kramer, H.A.; Mockrin, M.H.; Alexandre, P.M. [et al.]. 2018. **Rapid growth of the US wildland-urban interface raises wildfire risk**. *Proceedings of the National Academy of Sciences*. 115(13): 3314. <https://doi.org/10.1073/pnas.1718850115>.
326. Jakes, P.J.; Esposito, C.; Nelson, K.C.; Sturtevant, V.E.; Williams, D.R. [et al.]. 2012. **Best management practices for creating a community wildfire protection plan**. Gen. Tech. Rep. NRS-89. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 27 p. <https://doi.org/10.2737/NRS-GTR-89>. <https://doi.org/10.2737/NRS-GTR-89>.
327. Mockrin, M.H.; Fishler, H.K.; Stewart, S.I. 2020. **After the fire: perceptions of land use planning to reduce wildfire risk in eight communities across the United States**. *International Journal of Disaster Risk Reduction*. 45: 101444. <https://doi.org/10.1016/j.ijdrr.2019.101444>.
328. Boyd, I.L.; Freer-Smith, P.H.; Gilligan, C.A.; Godfray, H.C.J. 2013. **The consequence of tree pests and diseases for ecosystem services**. *Science*. 342(6160): 1235773. <https://doi.org/10.1126/science.1235773>.
329. Jactel, H.; Bauhus, J.; Boberg, J.; Bonal, D.; Castagneyrol, B. [et al.]. 2017. **Tree diversity drives forest stand resistance to natural disturbances**. *Current Forestry Reports*. 3(3): 223–243. <https://doi.org/10.1007/s40725-017-0064-1>.
330. Urban, J. 2008. **Up by roots: healthy soils and trees in the built environment**. Atlanta, GA: International Society of Arboriculture. 479 p.
331. Näsi, R.; Honkavaara, E.; Blomqvist, M.; Lyytikäinen-Saarenmaa, P.; Hakala, T. [et al.]. 2018. **Remote sensing of bark beetle damage in urban forests at individual tree level using a novel hyperspectral camera from UAV and aircraft**. *Urban Forestry & Urban Greening*. 30: 72–83. <https://doi.org/10.1016/j.ufug.2018.01.010>.
332. Zhang, J.; Huang, Y.; Pu, R.; Gonzalez-Moreno, P.; Yuan, L. [et al.]. 2019. **Monitoring plant diseases and pests through remote sensing technology: a review**. *Computers and Electronics in Agriculture*. 165: 104943. <https://doi.org/10.1016/j.compag.2019.104943>.
333. Raupp, M.J.; Cumming, A.B.; Raupp, E.C. 2006. **Street tree diversity in eastern North America and its potential for tree loss to exotic borers**. *Arboriculture and Urban Forestry*. 32(6): 297–304.

334. Li, H.P.; Wickham, J.D.; Bushley, K.; Wang, Z.G.; Zhang, B. [et al.]. 2020. **New approaches in urban forestry to minimize invasive species impacts: The case of Xiongan new area in China.** *Insects*. 11(5): 300. <https://doi.org/10.3390/insects11050300>.
335. The Nature Conservancy. 2021. **Healthy trees, healthy cities tree health initiative.** www.healthytreeshealthycities.org (accessed February 28, 2021).
336. Chornesky, E.A.; Bartuska, A.M.; Aplet, G.H.; Britton, K.O.; Cummings-Carlson, J. [et al.]. 2005. **Science priorities for reducing the threat of invasive species to sustainable forestry.** *Bioscience*. 55(4): 335–348. [https://doi.org/10.1641/10006-3568\(2005\)055\[0335:SPFRTT\]2.0.CO;2](https://doi.org/10.1641/10006-3568(2005)055[0335:SPFRTT]2.0.CO;2).
337. Ward, J.; Williams, S.; Worthley, T. 2013. **Japanese barberry control methods: reference guide for foresters and professional woodland managers.** Special Bulletin. Storrs, CT: The Connecticut Agricultural Experiment Station. 13 p. <https://doi.org/10.13140/RG.2.1.1880.3287>.
338. Tu, M. 2021. **Waipuna™ hot foam system.** <https://www.invasive.org/gist/tools/hotfoam.html> (accessed February 28, 2021).
339. Curtis, P.D. 2020. **After decades of suburban deer research and management in the eastern United States: where do we go from here?** *Human Wildlife Interactions*. 14(1): 16. <https://doi.org/10.26077/K7YE-K912>.
340. Hunt, V.M.; Magle, S.B.; Vargas, C.; Brown, A.W.; Lonsdorf, E.V. [et al.]. 2014. **Survival, abundance, and capture rate of eastern cottontail rabbits in an urban park.** *Urban Ecosystems*. 17(2): 547–560. <https://doi.org/10.1007/s11252-013-0334-z>.
341. Soulsbury, C.; White, P. 2015. **Human-wildlife interactions in urban areas: a review of conflicts, benefits and opportunities.** *Wildlife Research*. 42: 541–553. <https://doi.org/10.1071/WR14229>.
342. Côté, S.D.; Rooney, T.P.; Tremblay, J.-P.; Dussault, C.; Waller, D.M. 2004. **Ecological impacts of deer overabundance.** *Annual Review of Ecology, Evolution, and Systematics*. 35: 113–147. <https://doi.org/10.1146/annurev.ecolsys.35.021103.105725>.
343. Dale, V.H.; Joyce, L.A.; McNulty, S.; Neilson, R.P.; Ayres, M.P. [et al.]. 2001. **Climate change and forest disturbances.** *Bioscience*. 51(9): 723–734. [https://doi.org/10.1641/0006-3568\(2001\)051\[0723:CCAF-D\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0723:CCAF-D]2.0.CO;2).
344. Mohan, J.; Cox, R.; Iverson, L. 2009. **Composition and carbon dynamics of forests in northeastern North America in a future, warmer world.** *Canadian Journal of Forest Research*. 39(2): 213–230. <https://doi.org/10.1139/X08-185>.
345. Rodenhouse, N.; Christenson, L.; Parry, D.; Green, L.E. 2009. **Climate change effects on native fauna of northeastern forests.** *Canadian Journal of Forest Research*. 39: 249–263. <https://doi.org/10.1139/X08-160>.
346. Weiskopf, S.R.; Ledee, O.E.; Thompson, L.M. 2019. **Climate change effects on deer and moose in the Midwest.** *The Journal of Wildlife Management*. 83(4): 769–781. <https://doi.org/10.1002/jwmg.21649>.
347. Wolff, C.L.; Demarais, S.; Brooks, C.P.; Barton, B.T. 2020. **Behavioral plasticity mitigates the effect of warming on white-tailed deer.** *Ecology and Evolution*. 10(5): 2579–2587. <https://doi.org/10.1002/ece3.6087>.
348. Cornell University. 2021. **Community deer advisor - decision support for communities managing deer.** <https://deeradvisor.dnr.cornell.edu/> (accessed February 28, 2021).
349. Wiggers, E.P. 2011. **The evolution of an urban deer-management program through 15 years.** *Wildlife Society Bulletin*. 35(3): 137–141. <https://doi.org/10.1002/wsb.49>.
350. U.S. Department of Interior, National Park Service. 2020. **White-tailed deer management.** <https://www.nps.gov/rocr/learn/management/white-tailed-deer-management.htm> (accessed February 28, 2021).

351. Staudinger, M.D.; Carter, S.L.; Cross, M.S.; Dubois, N.S.; Duffy, J.E. [et al.]. 2013. **Biodiversity in a changing climate: a synthesis of current and projected trends in the US.** *Frontiers in Ecology and the Environment*. 11(9): 465–473. <https://doi.org/10.1890/120272>.
352. Espeland, E.K.; Kettenring, K.M. 2018. **Strategic plant choices can alleviate climate change impacts: a review.** *Journal of Environmental Management*. 222: 316–324. <https://doi.org/10.1016/j.jenvman.2018.05.042>.
353. Dreistadt, S.H.; Dahlsten, D.K.; Frankie, G.W. 1990. **Urban forests and insect ecology: complex interactions among trees, insects, and people.** *Bioscience*. 40(3): 192–198. <https://doi.org/10.2307/1311364>.
354. Poland, T.M.; McCullough, D.G. 2006. **Emerald ash borer: invasion of the urban forest and the threat to North America’s ash resource.** *Journal of Forestry*. 104(3): 118–124. <https://doi.org/10.1093/jof/104.3.118>.
355. Santamour, F.S.J. 1999. **Trees for urban planting: diversity uniformity, and common sense.** In: 7th conference of the Metropolitan Tree Planting Alliance (METRA): 57–65, <http://new.www.tree-care.info/mhattachments/pdficoI0kyRZI.pdf> (accessed February 28, 2021).
356. Whitlow, T.H.; Bassuk, N.L. 1987. **Trees in difficult sites.** *Journal of Arboriculture*. 13(1): 10–17. <http://joa.isa-arbor.com/request.asp?JournalID=1&ArticleID=2128&Type=2> (accessed February 28, 2021).
357. Kirshen, P.; Ruth, M.; Anderson, W. 2008. **Interdependencies of urban climate change impacts and adaptation strategies: a case study of metropolitan Boston USA.** *Climatic Change*. 86: 105–122. <https://doi.org/10.1007/s10584-007-9252-5>.
358. Curtis, P.S.; Gough, C.M. 2018. **Forest aging, disturbance and the carbon cycle.** *New Phytologist*. 219(4): 1188–1193. <https://doi.org/10.1111/nph.15227>.
359. Gunn, J.S.; Ducey, M.J.; Whitman, A.A. 2014. **Late-successional and old-growth forest carbon temporal dynamics in the northern forest (northeastern USA).** *Forest Ecology and Management*. 312: 40–46. <https://doi.org/10.1016/j.foreco.2013.10.023>.
360. Fahey, R.T.; Bowles, M.L.; McBride, J.L.; McPherson, E.; Scharenbroch, B. 2012. **Origins of the Chicago urban forest: composition and structure in relation to pre-settlement vegetation and modern land-use.** *Arboriculture & Urban Forestry*. 38: 115–133.
361. D’Amato, A.W.; Bradford, J.B.; Fraver, S.; Palik, B.J. 2011. **Forest management for mitigation and adaptation to climate change: insights from long-term silviculture experiments.** *Forest Ecology and Management*. 262(5): 803–816. <https://doi.org/10.1016/j.foreco.2011.05.014>.
362. Ford, S.E.; Keeton, W.S. 2017. **Enhanced carbon storage through management for old-growth characteristics in northern hardwood-conifer forests.** *Ecosphere*. 8(4): e01721. <https://doi.org/10.1002/ecs2.1721>.
363. Gunn, J.S.; Hagan, J.M.; Whitman, A.A. 2009. **Forestry adaptation and mitigation in a changing climate: a forest resource manager’s guide for the northeastern United States.** Ecosystem Services Resource Center Report NCI-2009-1. Brunswick, ME: Manomet Center for Conservation Sciences. 16 p. https://www.manomet.org/wp-content/uploads/old-files/Manomet_ForestryAdaptationtoCCReport%205-09.pdf (accessed February 28, 2021).
364. Piana, M.R.; Pregitzer, C.; Hallett, R.A. In press. **Advancing management in urban-forested natural areas: toward an urban silviculture?** *Frontiers in Ecology and the Environment*.
365. Piana, M.R.; Hallett, R.A.; Aronson, M.F.J.; Conway, E.; Handel, S.N. 2020. **Natural regeneration in urban forests is limited by early-establishment dynamics: implications for management.** *Ecological Applications*. 31(2): e02255. <https://doi.org/10.1002/eap.2255>.

366. Pastick, J.; Maurer, D.; Fahey, R.T. 2020. **Testing the effect of restoration-focused silviculture on oak regeneration and groundlayer plant communities in urban-exurban oak woodlands.** *Restoration Ecology*. 29(1): e13307. <https://doi.org/10.1111/rec.13307>.
367. Swanston, C.; Brandt, L.A.; Janowiak, M.K.; Handler, S.D.; Butler-Leopold, P. [et al.]. 2018. **Vulnerability of forests of the Midwest and Northeast United States to climate change.** *Climatic Change*. 146(1): 103–116. <https://doi.org/10.1007/s10584-017-2065-2>.
368. Vose, J.M.; Peterson, D.L.; Domke, G.M.; Fettig, C.J.; Joyce, L.A. [et al.]. 2018. **Forests.** In: Reidmiller, D.R.; Avery, C.W.; Easterling, D.R. [et al.], eds. *Impacts, risks, and adaptation in the United States: fourth national climate assessment, volume II.* Washington, DC: U.S. Global Change Research Program: 232–267. <https://doi.org/10.7930/NCA4.2018.CH6>.
369. Iverson, L.R.; Peters, M.P.; Prasad, A.M.; Matthews, S.N. 2019. **Analysis of climate change impacts on tree species of the eastern US: results of DISTRIB-II modeling.** *Forests*. 10(4): 302. <https://doi.org/10.3390/f10040302>.
370. Tallamy, D.W. 2009. **Bringing nature home: how you can sustain wildlife with native plants, updated and expanded.** Portland, OR: Timber Press. 360 p.
371. Brandt, L.; Lewis, A.; Fahey, R.; Scott, L.; Darling, L. [et al.]. 2016. **A framework for adapting urban forests to climate change.** *Environmental Science & Policy*. 66: 393–402. <https://doi.org/10.1016/j.envsci.2016.06.005>.
372. Marzluff, J.M.; Bowman, R.; Donnelly, R., eds. 2001. **Avian ecology and conservation in an urbanizing world.** Boston, MA: Kluwer Academic Publishers. 598 p.
373. Nowak, D.J.; Dwyer, J.F. 2007. **Understanding the benefits and costs of urban forest ecosystems.** In: Kuser J.E., ed. *Urban and Community Forestry in the Northeast.* Dordrecht, The Netherlands: Springer: 25–46. https://doi.org/10.1007/978-1-4020-4289-8_2.
374. Savard, J.-P.L.; , P.; G. Mennechez, G. 2000. **Biodiversity concepts and urban ecosystems.** *Landscape and Urban Planning*. 48: 131–142. [https://doi.org/10.1016/S0169-2046\(00\)00037-2](https://doi.org/10.1016/S0169-2046(00)00037-2).
375. Piana, M.R.; Aronson, M.F.J.; Pickett, S.T.A.; Handel, S.N. 2019. **Plants in the city: understanding recruitment dynamics in urban landscapes.** *Frontiers in Ecology and the Environment*. 17(8): 455–463. <https://doi.org/10.1002/fee.2098>.
376. Huber, K.; Gutierrez, D.; Poon, W.S.; Robinson, B. 2015. **Enhancing resiliency in Baltimore's urban forest.** Ann Arbor, MI: University of Michigan, SNRE. 94 p. <https://deepblue.lib.umich.edu/handle/2027.42/111049> (accessed February 28, 2021).
377. Purcell, L.; Daniel, K. 2016. **Tree selection for the "un-natural" environment.** FNR-531-W. West Lafayette, IN: Purdue University Extension. 9 p., https://www.edustore.purdue.edu/item.asp?Item_Number=FNR-531-W (accessed February 28, 2021).
378. Walther, G.-R.; Post, E.; Convey, P.; Menzel, A.; Parmesan, C. [et al.]. 2002. **Ecological responses to recent climate change.** *Nature*. 416: 389–395. <https://doi.org/10.1038/416389a>.
379. Young, A.G.; Merriam, H.G.; Warwick, S.I. 1993. **The effects of forest fragmentation on genetic variation in *Acer saccharum* Marsh. (sugar maple) populations.** *Heredity*. 71: 277–289. <https://www.nature.com/articles/hdy1993136.pdf> (accessed February 28, 2021).
380. Woodall, C.W.; Nowak, D.J.; Liknes, G.C.; Westfall, J.A. 2010. **Assessing the potential for urban trees to facilitate forest tree migration in the eastern United States.** *Forest Ecology and Management*. 259(8): 1447–1454. <https://doi.org/10.1016/j.foreco.2010.01.018>.
381. Almas, A.D.; Conway, T.M. 2016. **The role of native species in urban forest planning and practice: A case study of Carolinian Canada.** *Urban Forestry & Urban Greening*. 17: 54–62. <https://doi.org/10.1016/j.ufug.2016.01.015>.

382. Sjöman, H.; Morgenroth, J.; Sjöman, J.D.; Sæbø, A.; Kowarik, I. 2016. **Diversification of the urban forest—can we afford to exclude exotic tree species?** *Urban Forestry & Urban Greening*. 18: 237–241. <https://doi.org/10.1016/j.ufug.2016.06.011>.
383. McPherson, E.G.; Berry, A.M.; van Doorn, N.S. 2018. **Performance testing to identify climate-ready trees.** *Urban Forestry & Urban Greening*. 29: 28–39. <https://doi.org/10.1016/j.ufug.2017.09.003>.
384. Sjöman, H.; Hirons, A.D.; Bassuk, N.L. 2018. **Improving confidence in tree species selection for challenging urban sites: a role for leaf turgor loss.** *Urban Ecosystems*. 21(6): 1171–1188. <https://doi.org/10.1007/s11252-018-0791-5>.
385. McCarthy, M.P.; Best, M.J.; Betts, R.A. 2010. **Climate change in cities due to global warming and urban effects.** *Geophysical Research Letters*. 37(9). <https://doi.org/10.1029/2010GL042845>.
386. Thom, D.; Rammer, W.; Seidl, R. 2017. **Disturbances catalyze the adaptation of forest ecosystems to changing climate conditions.** *Global Change Biology*. 23(1): 269–282. <https://doi.org/10.1111/gcb.13506>.
387. Naeem, S.; Knops, J. M.; Tilman, D.; Howe, K. M.; Kennedy, T. [et al.]. 2000. **Plant diversity increases resistance to invasion in the absence of covarying extrinsic factors.** *Oikos*. 91: 97–108. <https://doi.org/10.1034/j.1600-0706.2000.910108.x>.
388. Peters, E.B.; Hiller, R.V.; McFadden, J.P. 2011. **Seasonal contributions of vegetation types to suburban evapotranspiration.** *Journal of Geophysical Research: Biogeosciences*. 116(G1). <https://doi.org/10.1029/2010JG001463>.
389. Peters, E.B.; McFadden, J.P. 2010. **Influence of seasonality and vegetation type on suburban microclimates.** *Urban Ecosystems*. 13(4): 443–460. <https://doi.org/10.1007/s11252-010-0128-5>.
390. Peters, E.B.; McFadden, J.P.; Montgomery, R.A. 2010. **Biological and environmental controls on tree transpiration in a suburban landscape.** *Journal of Geophysical Research: Biogeosciences*. 115(G4). <https://doi.org/10.1029/2009JG001266>.
391. Morse, N.B.; Pellissier, P.A.; Cianciola, E.N.; Brereton, R.L.; Sullivan, M.M. [et al.]. 2014. **Novel ecosystems in the anthropocene: a revision of the novel ecosystem concept for pragmatic applications.** *Ecology and Society*. 19(2): 12. <https://doi.org/10.5751/es-06192-190212>.
392. Thomas, E.; Jalonen, R.; Loo, J.; Boshier, D.; Gallo, L. [et al.]. 2014. **Genetic considerations in ecosystem restoration using native tree species.** *Forest Ecology and Management*. 333: 66–75. <https://doi.org/10.1016/j.foreco.2014.07.015>.
393. Higgs, E. 2017. **Novel and designed ecosystems.** *Restoration Ecology*. 25(1): 8–13. <https://doi.org/10.1111/rec.12410>.
394. Vitt, P.; Havens, K.; Kramer, A.T.; Sollenberger, D.; Yates, E. 2010. **Assisted migration of plants: changes in latitudes, changes in attitudes.** *Biological Conservation*. 143(1): 18–27. <https://doi.org/10.1016/j.biocon.2009.08.015>.
395. Pedlar, J.H.; McKenney, D.W.; Aubin, I.; Beardmore, T.; Beaulieu, J. [et al.]. 2012. **Placing forestry in the assisted migration debate.** *Bioscience*. 62(9): 835–842. <https://doi.org/10.1525/bio.2012.62.9.10>.
396. Ricciardi, A.; Simberloff, D. 2009. **Assisted colonization is not a viable conservation strategy.** *Trends in Ecology & Evolution*. 24(5): 248–253. <https://doi.org/10.1016/j.tree.2008.12.006>.
397. Ordóñez, C.; Duinker, P.N. 2014. **Assessing the vulnerability of urban forests to climate change.** *Environmental Reviews*. 22(3): 311–321. <https://doi.org/10.1139/er-2013-0078>.
398. Ordóñez, C.; Duinker, P.N. 2015. **Climate change vulnerability assessment of the urban forest in three Canadian cities.** *Climatic Change*. 131(4): 531–543. <https://doi.org/10.1007/s10584-015-1394-2>.
399. Davis, M.B.; Shaw, R.G. 2001. **Range shifts and adaptive responses to quaternary climate change.** *Science*. 292(5517): 673–679. <https://doi.org/10.1126/science.292.5517.673>.

400. Iverson, L.R.; Schwartz, M.W.; Prasad, A.M. 2004. **How fast and far might tree species migrate in the eastern United States due to climate change?** *Global Ecology and Biogeography*. 13(3): 209–219. <https://doi.org/10.1111/j.1466-822X.2004.00093.x>.
401. Pompe, S.; Hanspach, J.; Badeck, F.-W.; Klotz, S.; Bruelheide, H. [et al.]. 2010. **Investigating habitat-specific plant species pools under climate change.** *Basic and Applied Ecology*. 11(7): 603–611. <https://doi.org/10.1016/j.baae.2010.08.007>.
402. Ballantyne, M.; Gudes, O.; Pickering, C.M. 2014. **Recreational trails are an important cause of fragmentation in endangered urban forests: a case-study from Australia.** *Landscape and Urban Planning*. 130: 112–124. <https://doi.org/10.1016/j.landurbplan.2014.07.004>.
403. Collinge, S.K. 1996. **Ecological consequences of habitat fragmentation: implications for landscape architecture and planning.** *Landscape and Urban Planning*. 36(1): 59–77. [https://doi.org/10.1016/s0169-2046\(96\)00341-6](https://doi.org/10.1016/s0169-2046(96)00341-6).
404. Hartel, D.R. 2019. **Disaster recovery steps to maintain and improve urban forest resilience.** In: Campbell, L.K.; Svendsen, E.; Sonti, N.F. [et al.], eds. *Gen. Tech. Rep. NRS-P-185*. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station: 220–233 p. <https://doi.org/10.2737/NRS-GTR-P-185-paper15>.
405. Beach, R.H.; Sills, E.O.; Liu, T.-M.; Pattanayak, S. 2010. **The influence of forest management on vulnerability of forests to severe weather.** In: Pye, J.M.; Rauscher, H.M.; Sands, Y. [et al.], eds. *Gen. Tech. Rep. PNW-GTR-802*. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Stations: 185–206. <https://www.fs.usda.gov/treesearch/pubs/37042> (accessed February 21, 2021).
406. USDA Forest Service. 2021. **Phytoremediation of soils using fast-growing trees in vacant lots and landfills.** Washington, DC: U.S. Department of Agriculture, Forest Service. <https://www.fs.fed.us/research/urban-webinars/phytoremediation-of-soils.php> (accessed February 21, 2021).
407. Marshman, K. 2018. **The eye of the storm: extreme weather events and sustainable urban forest management.** *Dalhousie Journal of Interdisciplinary Management*. 14. <https://doi.org/10.5931/djim.v14i0.7876>.
408. Holling, C.S. 1973. **Resilience and stability of ecological systems.** *Annual Review of Ecology and Systematics*. 4: 1–23. <https://doi.org/10.1146/annurev.es.04.110173.000245>.
409. Spittlehouse, D.L.; Stewart, R.B. 2003. **Adaptation to climate change in forest management.** *BC Journal of Ecosystems and Management*. 4(1): art1. <https://www.cakex.org/sites/default/files/documents/Adaptation%20to%20climate%20change%20in%20forest%20management.pdf> (accessed May 10, 2021).
410. Frelich, L.E.; Jöggiste, K.; Stanturf, J.; Jansons, A.; Vodde, F. 2020. **Are secondary forests ready for climate change? It depends on magnitude of climate change, landscape diversity and ecosystem legacies.** *Forests*. 11(9): 965. <https://doi.org/10.3390/f11090965>.
411. Hobbs, R.J.; Arico, S.; Aronson, J.; Baron, J.S.; Bridgewater, P. [et al.]. 2006. **Novel ecosystems: theoretical and management aspects of the new ecological world order.** *Global Ecology and Biogeography*. 15: 1–7. <https://doi.org/10.1111/j.1466-822X.2006.00212.x>.
412. Rouse, D.C.; Bunsier-Ossa, I.F. 2013. **Green infrastructure: a landscape approach.** Chicago, IL: American Planning Association. 164 p. https://caeau.com.ar/wp-content/uploads/2018/11/46_GREEN-INFRAESTRUCTURE.pdf (accessed February 28, 2021).
413. Clayton, S.; Silka, L.; Trott, C.; Chapman, D.; Mancoll, S. 2016. **Building resilient communities in the face of climate change: a resource for local communities.** Washington, DC: Society for the Psychological Study of Social Issues. 10 p. <https://www.spssi.org/index.cfm?fuseaction=Page.ViewPage&PageID=2098> (accessed February 28, 2021).

414. Dodgen, D.; Donato, D.; Kelly, N.; La Greca, A.; Morganstein, J. [et al.]. 2016. **Ch. 8 - mental health and well-being: The impacts of climate change on human health in the United States: a scientific assessment.** In: Crimmins, A.; Balbus, J.; Gamble, J.L. [et al.], eds. Washington DC: U.S. Global Change Research Program: 217–246. <https://doi.org/10.7930/J0R49NQX>.
415. Huff, E.S.; Johnson, M.L.; Roman, L.A.; Sonti, N.F.; Pregitzer, C.C. [et al.]. 2020. **A literature review of resilience in urban forestry.** *Arboriculture & Urban Forestry*. 46: 185–196. <https://doi.org/10.48044/jauf.2020.014>.
416. USDA Forest Service. 2018. **Urban nature for human health and well-being: a research summary for communicating the health benefits of urban trees and green space.** FS-1096. Washington DC: U.S. Department of Agriculture, Forest Service. 24 p. <https://www.fs.fed.us/research/urban/humanhealth.php> (accessed May 3, 2021).
417. Ulrich, R.S. 1984. **View through a window may influence recovery from surgery.** *Science*. 224: 420–421. <https://doi.org/10.1126/science.6143402>.
418. Browning, M.H.E.M.; Rigolon, A. 2019. **School green space and its impact on academic performance: a systematic literature review.** *International Journal of Environmental Research and Public Health*. 16(3). <https://doi.org/10.3390/ijerph16030429>.
419. Yang, B.Y.; Markevych, I.; Bloom, M.S.; Heinrich, J.; Guo, Y. [et al.]. 2019. **Community greenness, blood pressure, and hypertension in urban dwellers: the 33 communities Chinese health study.** *Environment International*. 126: 727–734. <https://doi.org/10.1016/j.envint.2019.02.068>.
420. Takano, T.; Nakamura, K.; Watanabe, M. 2002. **Urban residential environments and senior citizens' longevity in megacity areas: the importance of walkable green spaces.** *Journal of Epidemiology and Community Health*. 56(12): 913–918. <https://doi.org/10.1136/jech.56.12.913>.
421. Bratman, G.N.; Anderson, C.B.; Berman, M.G.; Cochran, B.; de Vries, S. [et al.]. 2019. **Nature and mental health: an ecosystem service perspective.** *Science Advances*. 5: eaax0903. <https://doi.org/10.1126/sciadv.aax0903>.
422. Svendsen, E.S.; Campbell, L.K.; Sonti, N.F.; Baine, G. 2015. **Urban stewardship as a catalyst for recovery and change.** In: Brandt, D.H.; Nordenson, C.S., eds. *Waterproofing New York*. New York, NY: Terreform: 104–111. https://www.fs.fed.us/nrs/pubs/jrnl/2015/nrs_2015_svendsen_001.pdf (accessed February 28, 2021).
423. Padhy, S.K.; Sarkar, S.; Panigrahi, M.; Paul, S. 2015. **Mental health effects of climate change.** *Indian Journal of Occupational and Environmental Medicine*. 19: 3–7. <https://doi.org/10.4103%2F0019-5278.156997>.
424. Bielinis, E.; Jaroszewska, A.; Łukowski, A.; Takayama, N. 2020. **The effects of a forest therapy programme on mental hospital patients with affective and psychotic disorders.** *International Journal of Environmental Research and Public Health*. 17: 118. <https://doi.org/10.3390/ijerph17010118>.
425. Cianconi, P.; Betrò, S.; Janiri, L. 2020. **The impact of climate change on mental health: a systematic descriptive review.** *Front Psychiatry*. 11: 74. <https://doi.org/10.3389/fpsy.2020.00074>.
426. Bratman, G.N.; Hamilton, J.P.; Hahn, K.S.; Daily, G.C.; Gross, J.J. 2015. **Nature experience reduces rumination and subgenual prefrontal cortex activation.** *Proceedings of the National Academy of Sciences of the United States of America*. 112: 8567–8572. <https://doi.org/10.1073/pnas.1510459112>.
427. Nolen-Hoeksema, S. 2000. **The role of rumination in depressive disorders and mixed anxiety/depressive symptoms.** *Journal of Abnormal Psychology*. 109: 504–511. <https://doi.org/10.1037/0021-843x.109.3.504>.
428. Lopes, S.; Lima, M.; Silva, K. 2020. **Nature can get it out of your mind: the rumination reducing effects of contact with nature and the mediating role of awe and mood.** *Journal of Environmental Psychology*. 71: 101489. <https://doi.org/10.1016/j.jenvp.2020.101489>.

429. Song, H.; Lane, K.J.; Kim, H.; Kim, H.; Byun, G. [et al.]. 2019. **Association between urban greenness and depressive symptoms: evaluation of greenness using various indicators.** *International Journal of Environmental Research and Public Health*. 16: 173. <https://doi.org/10.3390/ijerph16020173>.
430. Dzhambov, A.M.; Hartig, T.; Tilov, B.; Atanasova, V.; Makakova, D.R. [et al.]. 2019. **Residential greenspace is associated with mental health via intertwined capacity-building and capacity-restoring pathways.** *Environmental Research*. 178: 10878. <https://doi.org/10.1016/j.envres.2019.108708>.
431. Berman, M.G.; Kross, E.; Krpan, K.M.; Askren, M.K.; Burson, A. [et al.]. 2012. **Interacting with nature improves cognition and affect for individuals with depression.** *Journal of Affective Disorders*. 140: 300–305. <https://doi.org/10.1016/j.jad.2012.03.012>.
432. Antonelli, M.; Barbieri, G.; Donelli, D. 2019. **Effects of forest bathing (shinrin-yoku) on levels of cortisol as a stress biomarker: a systematic review and meta-analysis.** *International Journal of Biometeorology*. 63: 1117–1134. <https://doi.org/10.1007/s00484-019-01717-x>.
433. Morita, E.; Fukuda, S.; Nagano, J.; Hamajima, N.; Yamamoto, H. [et al.]. 2007. **Psychological effects of forest environments on healthy adults: shinrin-yoku (forest-air bathing, walking) as a possible method of stress reduction.** *Public Health*. 121: 54–63. <https://doi.org/10.1016/j.puhe.2006.05.024>.
434. Wood, E.; Harsant, A.; Dallimer, M.; Cronin de Chavez, A.; McEachan, R.R.C. [et al.]. 2018. **Not all green space is created equal: biodiversity predicts psychological restorative benefits from urban green space.** *Frontiers in Psychology*. 9: 2320. <https://doi.org/10.3389/fpsyg.2018.02320>.
435. Wolf, L.J.; Zu Ermgassen, S.; Balmford, A.; White, M.; Weinstein, N. 2017. **Is variety the spice of life? An experimental investigation into the effects of species richness on self-reported mental well-being.** *PloS One*. 12: e0170225. <https://doi.org/10.1371/journal.pone.0170225>.
436. Kaplan, S.; Peterson, C. 1993. **Health and environment: a psychological analysis.** *Landscape and Urban Planning*. 26: 17–23. [https://doi.org/10.1016/0169-2046\(93\)90004-W](https://doi.org/10.1016/0169-2046(93)90004-W).
437. Kuo, F.E. 2001. **Coping with poverty: impacts of environment and attention in the inner city.** *Environment and Behavior*. 33: 5–34. <https://doi.org/10.1177/00139160121972846>.
438. Kaplan, S.; Berman, M.G. 2010. **Directed attention as a common resource for executive functioning and self-regulation.** *Perspectives on Psychological Science*. 5: 43–57. <https://doi.org/10.1177/1745691609356784>.
439. Plambech, T.; Konijnendijk van den Bosch, C.C. 2015. **The impact of nature on creativity – a study among Danish creative professionals.** *Urban Forestry & Urban Greening*. 14: 255–263. <https://doi.org/10.1016/j.ufug.2015.02.006>.
440. Leong, L.Y.C.; Fischer, R.; McClure, J. 2014. **Are nature lovers more innovative? The relationship between connectedness with nature and cognitive styles.** *Journal of Environmental Psychology*. 40: 57–63. <https://doi.org/10.1016/j.jenvp.2014.03.007>.
441. Oppezzo, M.; Schwartz, D.L. 2014. **Give your ideas some legs: the positive effect of walking on creative thinking.** *Journal of Experimental Psychology: Learning, Memory, and Cognition*. 40: 1142–1152. <https://doi.org/10.1037/a0036577>.
442. Egorov, A.I.; Griffin, S.M.; Converse, R.R.; Styles, J.N.; Sams, E.A. [et al.]. 2017. **Vegetated land cover near residence is associated with reduced allostatic load and improved biomarkers of neuroendocrine, metabolic and immune functions.** *Environmental Research*. 158: 508–521. <https://doi.org/10.1016/j.envres.2017.07.009>.
443. Mitchell, R.; Popham, F. 2008. **Effect of exposure to natural environment on health inequalities: an observational population study.** *The Lancet*. 372: 1655–1660. [https://doi.org/10.1016/S0140-6736\(08\)61689-X](https://doi.org/10.1016/S0140-6736(08)61689-X).

444. McEachan, R.R.C.; Prady, S.L.; Smith, G.; Fairley, L.; Cabieses, B. [et al.]. 2016. **The association between green space and depressive symptoms in pregnant women: moderating roles of socioeconomic status and physical activity.** *Journal of Epidemiology and Community Health.* 70: 253–259. <https://doi.org/10.1136/jech-2015-205954>.
445. Hunter, M.R.; Gillespie, B.W.; Chen, S.Y.-P. 2019. **Urban nature experiences reduce stress in the context of daily life based on salivary biomarkers.** *Frontiers in Psychology.* 10: 722. <https://doi.org/10.3389/fpsyg.2019.00722>.
446. Korpela, K.M.; Hartig, T.; Kaiser, F.G.; Fuhrer, U. 2001. **Restorative experience and self-regulation in favorite places.** *Environment and Behavior.* 33: 572–589. <https://doi.org/10.1177/00139160121973133>.
447. Koselka, E.P.D.; Weidner, L.C.; Minasov, A.; Berman, M.G.; Leonard, W.R. [et al.]. 2019. **Walking green: developing an evidence base for nature prescriptions.** *International Journal of Environmental Research and Public Health.* 16: 4338. <https://doi.org/10.3390%2Fijerph16224338>.
448. Watts, G. 2018. **Tranquillity trails for urban areas.** *Urban Forestry & Urban Greening.* 29: 154–161. <https://doi.org/10.1016/j.ufug.2017.11.015>.
449. Sullivan, W.C.; Kaplan, R. 2016. **Nature! Small steps that can make a big difference.** *Health Environments Research & Design Journal.* 9: 6–10. <https://doi.org/10.1177%2F1937586715623664>.
450. Berman, M.G.; Jonides, J.; Kaplan, S. 2008. **The cognitive benefits of interacting with nature.** *Psychological Science.* 19: 1207–1212. <https://doi.org/10.1111/j.1467-9280.2008.02225.x>.
451. Berto, R. 2014. **The role of nature in coping with psycho-physiological stress: a literature review on restorativeness.** *Behavioral Science.* 4: 394–409. <https://doi.org/10.3390/bs4040394>.
452. Kaplan, R.; Kaplan, S.; Ryan, R.L. 1998. **With people in mind: design and management of everyday nature.** Washington DC: Island Press: 225 p.
453. Hoyle, H.; Hitchmough, J.; Jorgensen, A. 2017. **All about the 'wow factor'? The relationships between aesthetics, restorative effect and perceived biodiversity in designed urban planting.** *Landscape and Urban Planning.* 164: 109–123. <https://doi.org/10.1016/j.landurbplan.2017.03.011>.
454. Tasnim, N.; Abulizi, N.; Pither, J.; Hart, M.M.; Gibson, D.L. 2017. **Linking the gut microbial ecosystem with the environment: does gut health depend on where we live?** *Frontiers in Microbiology.* 8: 1935. <https://doi.org/10.3389/fmicb.2017.01935>.
455. Fischer, L.K.; Brinkmeyer, D.; Karle, S.J.; Cremer, K.; Huttner, E. [et al.]. 2019. **Biodiverse edible schools: linking healthy food, school gardens and local urban biodiversity.** *Urban Forestry and Urban Greening.* 40: 35–43. <https://doi.org/10.1016/j.ufug.2018.02.015>.
456. Basner, M.; Babisch, W.; Davis, A.; Brink, M.; Clark, C. [et al.]. 2014. **Auditory and non-auditory effects of noise on health.** *The Lancet.* 383: 1325–1332. [https://doi.org/10.1016/S0140-6736\(13\)61613-x](https://doi.org/10.1016/S0140-6736(13)61613-x).
457. Van Hedger, S.C.; Nusbaum, H.C.; Clohisy, L.; Jaeggi, S.M.; Buschkuhl, M. [et al.] 2019. **Of cricket chirps and car horns: the effect of nature sounds on cognitive performance.** *Psychonomic Bulletin & Review.* 26: 522–530. <https://doi.org/10.3758/s13423-018-1539-1>.
458. Rey Gozalo, G.; Barrigón Morillas, J.M.; Montes González, D.; Atanasio Moraga, P. 2018. **Relationships among satisfaction, noise perception, and use of urban green spaces.** *Science of the Total Environment.* 624: 438–450. <https://doi.org/10.1016/j.scitotenv.2017.12.148>.
459. Kuo, M.; Barnes, M.; Jordan, C. 2019. **Do experiences with nature promote learning? Converging evidence of a cause-and-effect relationship.** *Frontiers in Psychology.* 10: 305. <https://doi.org/10.3389/fpsyg.2019.00305>.

460. Kuo, M.; Browning, M.H.E.M.; Sachdeva, S.; Lee, K.; Westphal, L. 2018. **Might school performance grow on trees? Examining the link between "greenness" and academic achievement in urban, high-poverty schools.** *Frontiers in Psychology*. 9: 1669. <https://doi.org/10.3389/fpsyg.2018.01669>.
461. Sivarajah, S.; Smith, S.M.; Thomas, S.C. 2018. **Tree cover and species composition effects on academic performance of primary school students.** *PloS One*. 13: e0193254. <https://doi.org/10.1371/journal.pone.0193254>.
462. Wu, J.; Jackson, L. 2017. **Inverse relationship between urban green space and childhood autism in California elementary school districts.** *Environment International*. 107: 140–146. <https://doi.org/10.1016/j.envint.2017.07.010>.
463. Engemann, K.; Pedersen, C.B.; Arge, L.; Tsirogiannis, C.; Mortensen, P.B. [et al.]. 2019. **Residential green space in childhood is associated with lower risk of psychiatric disorders from adolescence into adulthood.** *Proceedings of the National Academy of Sciences of the United States of America*. 116: 5188–5193. <https://doi.org/10.1073/pnas.1807504116>.
464. Wolf, K.L.; Housley, E. 2016. **The benefits of nearby nature in cities for older adults.** Annapolis, MD: TKF Foundation. 16 p. <https://naturesacred.org/research/the-benefits-of-nearby-nature-in-cities-for-older-adults/> (accessed February 28, 2021).
465. Yu, C.-P.; Hsieh, H. 2020. **Beyond restorative benefits: evaluating the effect of forest therapy on creativity.** *Urban Forestry & Urban Greening*. 51: 126670. <https://doi.org/10.1016/j.ufug.2020.126670>.
466. Williams, K.J.H.; Lee, K.E.; Hartig, T.; Sargent, L.D.; Williams, N.S.G. [et al.]. 2018. **Conceptualising creativity benefits of nature experience: attention restoration and mind wandering as complementary processes.** *Journal of Environmental Psychology*. 59: 36–45. <https://doi.org/10.1016/j.jenvp.2018.08.005>.
467. Wolf, K.L. 2017. **Social aspects of urban forestry and metro nature.** In: Ferrini, F.; Konijnendijk van den Bosch, C.C.; Fini, A., eds. *Routledge handbook of urban forestry*. New York, NY: Routledge: 65–81.
468. Putnam, R. 2000. **Bowling alone: the collapse and revival of American community.** New York, NY: Simon and Schuster. 544 p.
469. Elands, B.H.M.; Peters, K.B.M.; de Vries, S. 2018. **Promoting social cohesion and social capital – increasing wellbeing.** In: van den Bosch, M.; Bird, W., eds. *Nature and public health*. Oxford, UK: Oxford University Press: 116–121. <https://doi.org/10.1093/med/9780198725916.003.0044>.
470. Morina, N.; Kip, A.; Hoppen, T.H.; Priebe, S.; Meyer, T. 2021. **A potential impact of physical distancing on physical and mental health. A rapid narrative umbrella review of meta-analyses on the link between social isolation and health.** *BJM Open*. <https://bmjopen.bmj.com/content/bmjopen/11/3/e042335.full.pdf> (accessed May 10, 2021).
471. Jennings, V. 2019. **Social cohesion and city green space: revisiting the power of volunteering.** *Challenges*. 10: 36. <https://doi.org/10.3390/challe10020036>.
472. Tidball, K.G. 2012. **Urgent biophilia: human-nature interactions and biological attractions in disaster resilience.** *Ecology and Society*. 17: 5. <https://doi.org/10.5751/ES-04596-170205>.
473. Campbell, L.K.; Svendsen, E.; Sonti, N.F.; Hines, S.J.; Maddox, D. 2019. **Green readiness, response, and recovery: a collaborative synthesis.** Gen. Tech. Rep. NRS-P-185. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 358 p. <https://doi.org/10.2737/NRS-GTR-P-185>.
474. Meerow, S.; Newell, J.P.; Stults, M. 2016. **Defining urban resilience: a review.** *Landscape and Urban Planning*. 147: 38–49. <https://doi.org/10.1016/j.landurbplan.2015.11.011>.

475. Kwok, A.H.; Doyle, E.E.H.; Becker, J.; Johnston, D.; Paton, D. 2016. **What is 'social resilience?' Perspectives of disaster researchers, emergency management practitioners, and policy-makers in New Zealand.** *International Journal of Disaster Risk Reduction*. 19: 197–211. <https://doi.org/10.1016/j.ijdrr.2016.08.013>.
476. Pradhananga, A.K.; Davenport, M.A. 2017. **Community attachment, beliefs and residents' civic engagement in stormwater management.** *Landscape and Urban Planning*. 168: 1–8. <https://doi.org/10.1016/j.landurbplan.2017.10.001>.
477. Chan, T.; Kihlsinger, R.; Hare-Grogg, J.; Jensen, M.; Conn, R. 2018. **Step-by-step guide to integrating community input into green infrastructure projects.** Washington, DC: Environmental Law Institute; and Taos, NM: Amigos Bravos. 19 p. <https://www.eli.org/research-report/step-step-guide-integrating-community-input-green-infrastructure-projects> (accessed February 28, 2021).
478. Ordóñez-Barona, C. 2017. **How different ethno-cultural groups value urban forests and its implications for managing urban nature in a multicultural landscape: a systematic review of the literature.** *Urban Forestry and Urban Greening*. 26: 65–77. <https://doi.org/10.1016/j.ufug.2017.06.006>.
479. McMillen, H.; Campbell, L.K.; Svendsen, E.S.; Reynolds, R. 2016. **Recognizing stewardship practices as indicators of social resilience: in living memorials and in a community garden.** *Sustainability*. 8: 775. <https://doi.org/10.3390/su8080775>.
480. Holtan, M.T.; Dieterlen, S.L.; Sullivan, W.C. 2015. **Social life under cover tree canopy and social capital in Baltimore, Maryland.** *Environment and Behavior*. 47: 502–525. <https://doi.org/10.1177%2F0013916513518064>.
481. Toomey, A.H.; Stehlau-Howay, L.; Manzolilo, B.; Thomas, C. 2020. **The place-making potential of citizen science: creating social-ecological connections in an urbanized world.** *Landscape and Urban Planning*. 200: 103824. <https://doi.org/10.1016/j.landurbplan.2020.103824>.
482. Pandya, R.E. 2012. **A framework for engaging diverse communities in citizen science in the US.** *Frontiers in Ecology and the Environment*. 10: 314–317. <https://doi.org/10.1890/120007>.
483. Svendsen, E.S.; Campbell, L.K.; Fisher, D.R.; Connolly, J.J.T.; Johnson, M.L. [et al.]. 2016. **Stewardship mapping and assessment project: a framework for understanding community-based environmental stewardship.** Gen. Tech. Rep. NRS-156. Newton Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 134 p. <https://doi.org/10.2737/NRS-GTR-156>.
484. Whitburn, J.; Linklater, W.L.; Milfont, T.L. 2018. **Exposure to urban nature and tree planting are related to pro-environmental behavior via connection to nature, the use of nature for psychological restoration, and environmental attitudes.** *Environment and Behavior*. 51(7): 787–810. <https://doi.org/10.1177/0013916517751009>.
485. Power, A.; Smyth, K. 2016. **Heritage, health and place: the legacies of local community-based heritage conservation on social wellbeing.** *Health & Place*. 39: 160–167. <https://doi.org/10.1016/j.healthplace.2016.04.005>.
486. Peters, K.; Stodolska, M.; Horolets, A. 2016. **The role of natural environments in developing a sense of belonging: a comparative study of immigrants in the U.S., Poland, the Netherlands and Germany.** *Urban Forestry & Urban Greening*. 17: 63–70. <https://doi.org/10.1016/j.ufug.2016.04.001>.
487. Ekenga, C.C.; Sprague, N.; Shobiye, D.M. 2019. **Promoting health-related quality of life in minority youth through environmental education and nature contact.** *Sustainability*. 11: 3544. <https://doi.org/10.3390/su11133544>.
488. McMillen, H.L.; Campbell, L.K.; Svendsen, E.S. 2017. **Co-creators of memory, metaphors for resilience, and mechanisms for recovery: flora in living memorials to 9/11.** *Journal of Ethnobiology*. 37: 1–20. <https://doi.org/10.2993/0278-0771-37.1.1>.

489. Svendsen, E.S.; Campbell, L.K.; McMillen, H.L. 2016. **Stories, shrines, and symbols: recognizing psycho-social-spiritual benefits of urban parks and natural areas.** *Journal of Ethnobiology*. 36: 881–907. <https://doi.org/10.2993/0278-0771-36.4.881>.
490. Tidball, K.G.; Krasny, M., eds. 2014. **Greening in the red zone: disaster, resilience and community greening.** New York, NY: Springer: 534 p.
491. Wolf, K.L.; Housley, E. 2016. **The sacred & nearby nature in cities.** Annapolis, MD: TKF Foundation. 64 p. <https://naturesacred.org/research/the-sacred-and-nearby-in-nature-and-cities> (accessed February 28, 2021).
492. Schusler, T.; Weiss, L.; Treering, D.; Balderama, E. 2018. **Examining the association between tree canopy, parks and crime in Chicago.** *Landscape and Urban Planning*. 170: 309–313. <https://doi.org/10.1016/j.landurbplan.2017.07.012>.
493. Pfeiffer, C.; Vogelheim, C.; Wolf, K.; Ciecko, L.; Yadrick, M. [et al.]. 2019. **Best management practices for crime prevention through environmental design in natural landscapes.** Seattle, WA: Forterra-Green City Partnerships. 36 p. https://www.greenseattle.org/wp-content/uploads/2019/02/CPTED-in-Natural-Areas-Final-Draft-Feb-2018_web.pdf (accessed February 28, 2021).
494. Kuo, F.E.; Taylor, A.F. 2004. **A potential natural treatment for attention-deficit/hyperactivity disorder: evidence from a national study.** *American Journal of Public Health*. 94: 1580–1586. <https://doi.org/10.2105%2Fajph.94.9.1580>.
495. U.S. Centers for Disease Control and Prevention. 2021. **About physical activity.** <https://www.cdc.gov/physicalactivity/about-physical-activity/index.html> (accessed February 28, 2021).
496. Donovan, G.H. 2017. **Including public-health benefits of trees in urban-forestry decision making.** *Urban Forestry & Urban Greening*. 22: 120–123. <https://doi.org/10.1016/j.ufug.2017.02.010>.
497. Wolf, K.L. 2017. **The sanitary to sustainable city: place, health, and trees.** *Arborist News*. 40–43.
498. Graham, G.N. 2016. **Why your zip code matters more than your genetic code: promoting healthy outcomes from mother to child.** *Breastfeeding Medicine*. 11: 396–397. <https://doi.org/10.1089/bfm.2016.0113>.
499. Wolf, K.L.; The Nature Conservancy. 2018. **Cascading benefits: designing green stormwater infrastructure for human wellness.** Seattle, WA: The Nature Conservancy. 24 p. http://www.cityhabitats.org/wp-content/uploads/2018/05/tnc_gsi-report_2018_final-r1_digital.pdf (accessed February 28, 2021).
500. Jennings, V.; Yun, J.; Larson, L. 2016. **Finding common ground: environmental ethics, social justice, and a sustainable path for nature-based health promotion.** *Healthcare*. 4: 9. <https://doi.org/10.3390/healthcare4030061>.
501. Zuniga-Teran, A.A.; Orr, B.J.; Gimblett, R.H.; Chalfoun, N.V.; Guertin, D.P. [et al.]. 2017. **Neighborhood design, physical activity, and wellbeing: applying the walkability model.** *International Journal of Environmental Research and Public Health*. 14: 76. <https://doi.org/10.3390/ijerph14010076>.
502. Nehme, E.K.; Oluyomi, A.O.; Calise, T.V.; Kohl, H.W. 2016. **Environmental correlates of recreational walking in the neighborhood.** *American Journal of Health Promotion*. 30: 139–148. <https://doi.org/10.4278/ajhp.130531-QUAN-281>.
503. U.S. Centers for Disease Control and Prevention. 2021. **Physical activity recommendations for different age groups.** <https://www.cdc.gov/physicalactivity/basics/age-chart.html> (accessed February 28, 2021).
504. Evenson, K.R.; Williamson, S.; Han, B.; McKenzie, T.L.; Cohen, D.A. 2019. **United States' neighborhood park use and physical activity over two years: the national study of neighborhood parks.** *Preventive Medicine*. 123: 117–122. <https://doi.org/10.1016/j.ypmed.2019.03.027>.
505. Cohen, D.A.; Han, B.; Nagel, C.J.; Harnik, P.; McKenzie, T.L. [et al.]. 2016. **The first national study of neighborhood parks: implications for physical activity.** *American Journal of Preventive Medicine*. 51: 419–426. <https://doi.org/10.1016/j.amepre.2016.03.021>.

506. Marselle, M.R.; Warber, S.L.; Irvine, K.N. 2019. **Growing resilience through interaction with nature: can group walks in nature buffer the effects of stressful life events on mental health?** *International Journal of Environmental Research and Public Health*. 16: 986. <https://doi.org/10.3390/ijerph16060986>.
507. Tilt, J.H. 2010. **Walking trips to parks: exploring demographic, environmental factors, and preferences for adults with children in the household.** *Preventive Medicine*. 50 (Suppl 1): S69–73. <https://doi.org/10.1016/j.ypmed.2009.07.026>.
508. U.S. Department of Transportation. Complete streets. <https://www.transportation.gov/mission/health/complete-streets> (accessed March 10, 2021).
509. Kweon, B.; Naderi, J.R.; Maghelal, P.; Shin, W. 2004. **Correlates of environmental constructs and perceived safety enhancements in pedestrian corridors adjacent to urban street.** College Station, TX: Texas A&M University. 91 p.
510. Shanahan, D.F.; Bush, R.; Gaston, K.J.; Lin, B.B.; Dean, J. [et al.]. 2016. **Health benefits from nature experiences depend on dose.** *Scientific Reports*. 6: 28551. <https://doi.org/10.1038/srep28551>.
511. White, M.P.; Alcock, I.; Grellier, J.; Wheeler, B.W.; Hartig, T. [et al.]. 2019. **Spending at least 120 minutes a week in nature is associated with good health and wellbeing.** *Scientific Reports*. 9: 7730. <https://doi.org/10.1038/s41598-019-44097-3>.
512. Liddicoat, C.; Bi, P.; Waycott, M.; Glover, J.; Lowe, A.J. [et al.]. 2018. **Landscape biodiversity correlates with respiratory health in Australia.** *Journal of Environmental Management*. 206: 113–122. <https://doi.org/10.1016/j.jenvman.2017.10.007>.
513. Lederbogen, F.; Kirsch, P.; Haddad, L.; Streit, F.; Tost, H. [et al.]. 2011. **City living and urban upbringing affect neural social stress processing in humans.** *Nature*. 474: 498–501. <https://doi.org/10.1038/nature10190>.
514. Ewert, A.; Chang, Y. 2018. **Levels of nature and stress response.** *Behavioral Sciences*. 8: 49. <https://doi.org/10.3390/bs8050049>.
515. Cordoza, M.; Ulrich, R.S.; Manulik, B.J.; Gardiner, S.K.; Fitzpatrick, P.S. [et al.]. 2018. **Impact of nurses taking daily work breaks in a hospital garden on burnout.** *American Journal of Critical Care*. 27: 508–512. <https://doi.org/10.4037/ajcc2018131>.
516. Chang, C.-c.; Oh, R.R.Y.; Nghiem, T.P.L.; Zhang, Y.; Tan, C.L.Y. [et al.]. 2020. **Life satisfaction linked to the diversity of nature experiences and nature views from the window.** *Landscape and Urban Planning*. 202: 103874. <https://doi.org/10.1016/j.landurbplan.2020.103874>.
517. Currie, M.A. 2017. **A design framework for small parks in ultra-urban, metropolitan, suburban and small town settings.** *Journal of Urban Design*. 22: 76–95. <https://doi.org/10.1080/13574809.2016.1234334>.
518. Hadavi, S.; Kaplan, R.; Hunter, M.C.R. 2015. **Environmental affordances: A practical approach for design of nearby outdoor settings in urban residential areas.** *Landscape and Urban Planning*. 134: 19–32. <https://doi.org/10.1016/j.landurbplan.2014.10.001>.
519. Ribe, R. 1989. **The aesthetics of forestry: what has empirical preference research taught us?** *Environmental Management*. 13: 55–74. <https://doi.org/10.1007/BF01867587>.
520. Barron, S.; Nitoslawski, S.; Wolf, K.L.; Woo, A.; Desautels, E. [et al.]. 2019. **Greening blocks: a conceptual typology of practical design interventions to integrate health and climate resilience co-benefits.** *International Journal of Environmental Research and Public Health*. 16: 4241. <https://doi.org/10.3390/ijerph16214241>.
521. Felsten, G. 2009. **Where to take a study break on the college campus: an attention restoration theory perspective.** *Journal of Environmental Psychology*. 29: 160–167. <https://doi.org/10.1016/j.jenvp.2008.11.006>.
522. Kaplan, R. 1993. **The role of nature in the context of the workplace.** *Landscape and urban planning*. 26: 193–201. [https://doi.org/10.1016/0169-2046\(93\)90016-7](https://doi.org/10.1016/0169-2046(93)90016-7).

523. U.S. Environmental Protection Agency. 2017. **Learn about green streets.** <https://www.epa.gov/G3/learn-about-green-streets> (accessed February 28, 2021).
524. Vibrant Cities Lab. 2021. **Complete streets.** <https://www.vibrantcitieslab.com/research/city-planning-complete-streets/> (accessed February 28, 2021).
525. Wen, Y.; Yan, Q.; Pan, Y.; Gu, X.; Liu, Y. 2019. **Medical empirical research on forest bathing (shinrin-yoku): a systematic review.** *Environmental Health and Preventive Medicine.* 24: 1–21. <https://doi.org/10.1186%2Fs12199-019-0822-8>.
526. Hansen, M.M.; Jones, R.; Tocchini, K. 2017. **Shinrin-yoku (forest bathing) and nature therapy: a state-of-the-art review.** *International Journal of Environmental Research and Public Health.* 14: 851. <https://doi.org/10.3390/ijerph14080851>.
527. Colinas, J.; Bush, P.; Manaugh, K. 2019. **The socio-environmental impacts of public urban fruit trees: a Montreal case-study.** *Urban Forestry & Urban Greening.* 45: 126–132. <https://doi.org/10.1016/j.ufug.2018.05.002>.
528. Viladrich, A.; Tagliaferro, B. 2016. **Picking fruit from our backyard's trees: the meaning of nostalgia in shaping Latinas' eating practices in the United States.** *Appetite.* 97: 101–110. <https://doi.org/10.1016/j.appet.2015.11.017>.
529. Bukowski, C.; Munsell, J. 2018. **The community food forest handbook: how to plan, organize, and nurture edible gathering places.** Hartford, VT: Chelsea Green Publishing. 272 p.
530. Kowalski, J.M.; Conway, T.M. 2019. **Branching out: the inclusion of urban food trees in Canadian urban forest management plans.** *Urban Forestry & Urban Greening.* 45: 126142. <https://doi.org/10.1016/j.ufug.2018.05.012>.
531. McLain, R.; Poe, M.; Hurley, P.T.; Lecompte-Mastenbrook, J.; Emery, M.R. 2012. **Producing edible landscapes in Seattle's urban forest.** *Urban Forestry & Urban Greening.* 11: 187–194. <https://doi.org/10.1016/j.ufug.2011.12.002>.
532. Ulrich, R.S.; Bogren, L.; Gardiner, S.K.; Lundin, S. 2018. **Psychiatric ward design can reduce aggressive behavior.** *Journal of Environmental Psychology.* 57: 53–66. <https://doi.org/10.1016/j.jenvp.2018.05.002>.
533. Yoshimura, H.; Zhu, H.; Wu, Y.; Ma, R. 2010. **Spectral properties of plant leaves pertaining to urban landscape design of broad-spectrum solar ultraviolet radiation reduction.** *International Journal of Biometeorology.* 54: 179–191. <https://doi.org/10.1007/s00484-009-0267-7>.
534. Grant, R.H.; Heisler, G.M.; Gao, W. 2002. **Estimation of pedestrian level UV exposure under trees.** *Photochemistry and Photobiology.* 75: 369–376. [https://doi.org/10.1562/0031-8655\(2002\)075<0369:EOPLUE>2.0.CO;2](https://doi.org/10.1562/0031-8655(2002)075<0369:EOPLUE>2.0.CO;2).
535. Janowiak, M.K.; D'Amato, A.W.; Swanston, C.; Iverson, L.; Thompson, F., III [et al.]. 2018. **New England and New York forest ecosystem vulnerability assessment and synthesis: a report from the New England climate change response framework.** Gen. Tech. Rep. NRS-173. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 234 p. <https://doi.org/10.2737/nrs-gtr-173>.
536. USDA Forest Service. 2018. **Climate change pressures in the 21st century.** Washington, DC: U.S. Department of Agriculture, Forest Service, Office of Sustainability and Climate. https://usfs.maps.arcgis.com/apps/MapSeries/index.html?appid=96088b1c086a4b39b3a75d0fd97a4c40_story_map (accessed June 14, 2021).
537. Iverson, L.R.; Prasad, A.M.; Peters, M.P.; Matthews, S.N. 2019. **Facilitating adaptive forest management under climate change: A spatially specific synthesis of 125 species for habitat changes and assisted migration over the eastern United States.** *Forests.* 10(11): 989. <https://doi.org/10.3390/f10110989>.

538. Peters, M.P.; Iverson, L.R.; Prasad, A.M.; Matthews, S.N. 2019. **Utilizing the density of inventory samples to define a hybrid lattice for species distribution models: DISTRIB-II for 135 eastern U.S. trees.** *Ecology and Evolution*. 9(15): 8876–8899. <https://doi.org/10.1002/ece3.5445>.
539. Prasad, AM; Iverson, LR; Peters, MP; Matthews, SN. 2020. **Climate change tree atlas.** Madison, WI: U.S. Department of Agriculture, Forest Service, Northern Research Station. <https://doi.org/10.2737/climate-change-tree-atlas-v4>.
540. Brandt, L.A.; Derby Lewis, A.; Scott, L.; Darling, L.; Fahey, R.T. [et al.]. 2017. **Chicago Wilderness region urban forest vulnerability assessment and synthesis: a report from the Urban Forestry Climate Change Response Framework Chicago Wilderness pilot project.** Gen. Tech. Rep. NRS-168. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, Northern Research Station. 142 p. <https://doi.org/10.2737/NRS-GTR-168>.
541. i-Tree Team. N.d. **i-Tree species.** Washington, DC: U.S. Department of Agriculture, Forest Service; Kent, OH: Davey Tree Expert Co.; and other cooperators. <https://species.itreetools.org/> (accessed June 14, 2021).
542. Ogren, T.L. 2000. **Allergy-free gardening: the revolutionary guide to healthy landscaping.** Berkeley, CA: Ten Speed Press.

APPENDIX 1

Adaptation Workbook Steps in Brief

This is a brief outline of the Adaptation Workbook (Fig. 5) process. Find the full process in the “Forest Adaptation Resources: Climate Change Tools and Approaches for Land Managers, 2nd edition”²¹ and as an online tool at www.adaptationworkbook.org.




Figure 5.—The Adaptation Workbook²¹ describes an assessment and decision process that is used in conjunction with vulnerability assessments, local knowledge, and adaptation strategies menus. The results are site-specific actions that address explicit management and conservation objectives under a range of potential future climates.

Step 1: DEFINE location, project, and time frames

“What are your management goals and objectives for the project area?”

The first step is to describe the project area and your management objectives before considering the potential effects of climate change. This may include identifying:

- ▶ Any ecosystem types, stands, or other distinct areas that you want to consider individually.
- ▶ Any short- or long-term milestones that can be used to evaluate progress.

Step 2: ASSESS site-specific climate change impacts and vulnerabilities

“What climate change impacts and vulnerabilities are most important to this particular site?”

Climate change will have a wide variety of effects on the landscape, and not all places will respond similarly. List site-specific factors that may increase or reduce the effects of climate change in your project area, such as:

- ▶ Site conditions, including topographic position, soils, or hydrology.
- ▶ Past and current management.
- ▶ Forest composition and structure.
- ▶ Increasing exposure to pests, diseases, or other stressors.

Step 3: EVALUATE management objectives given projected impacts and vulnerabilities

“What management challenges and opportunities may occur as a result of climate change?”

This step explores management challenges and opportunities that may arise under changing conditions. For each of your management objectives, consider:

- ▶ Management challenges and opportunities given the climate impacts you identified previously.
- ▶ The feasibility of meeting each management objective under current management.
- ▶ Other considerations (e.g., administrative, legal, or social considerations) beyond climate change that may affect your ability to meet your management objectives.

Step 4: IDENTIFY adaptation approaches and tactics for implementation

“What actions can enhance the ability of the ecosystem to adapt to anticipated changes and meet management goals?”

Generate a list of adaptation tactics—prescriptive actions specifically designed for your project area or property and your unique management objectives. Use the Urban Forest Climate and Health Adaptation Menu as a starting point for identifying specific management tactics (e.g., what, how, and when) that you can implement. As you develop tactics, consider the following:

- ▶ Benefits, drawbacks, and barriers associated with each tactic.
- ▶ Effectiveness and feasibility of each tactic.

Step 5: MONITOR and evaluate effectiveness of implemented actions.

“What information can be used to evaluate if the selected actions were effective and inform future management?”

Monitoring metrics can help you determine if you are making progress on your management goals and evaluate the effectiveness of those actions. When identifying monitoring items, work to identify monitoring items that:

- ▶ Identify if your management goals and objectives were achieved.
- ▶ Evaluate if the adaptation tactics had the intended effect.
- ▶ Are realistic to implement.

APPENDIX 2

Adaptation Demonstration: Climate and Human Health Adaptation on a Neighborhood Scale in Providence, Rhode Island

Adaptation demonstrations are examples of organizations applying the Adaptation Workbook²¹ process and adaptation menus to their real-world natural resource management projects to generate explicit adaptation tactics in alignment with their objectives. More than 400 adaptation demonstration projects have been generated using other adaptation menus and are available online. The following demonstration was used in December 2020 to evaluate the Urban Forest Climate and Health Adaptation Menu (Menu).

Project Area and Management Goals

As public and private urban forest managers working in partnership to build a more equitable and robust urban forest, the Providence Parks Department and the Providence Neighborhood Planting Program (PNPP) aim to engage residents and neighborhood stakeholders in developing and implementing community-driven tree planting and stewardship solutions focused on climate adaptation and human health in Upper & Lower South Providence, two low-canopy and low-income neighborhoods disproportionately burdened by the impacts of climate change and environmental injustice. The general project area is primarily residential and light commercial, bordered by Interstate 95 (I-95) and the industrial port of Providence to the east and hospital campuses to the north, which are surrounded by large areas of surface parking lots.

The primary management focus is on right-of-way and front-of-property planting sites along streets between Broad Street, Eddy Street, Dudley Street, and I-95. The existing canopy cover is relatively low (20 percent) and lacks diversity. The most common tree species include callery pear (*Pyrus calleryana*), red maple, Japanese zelkova (*Zelkova serrata*), and honey locust (*Gleditsia triacanthos*). There are many highly disturbed and polluted sites in the neighborhood; residents battle poor air quality and high asthma rates. The eight census block groups that make up the project area have a population of 8,493, with 94 percent identifying as non-Caucasian. The average unemployment rate is 17 percent, and 69 percent of the population has a household income that falls below 200 percent of the poverty line.

Natural resource professionals from the City of Providence and the PNPP used the Adaptation Workbook and the Menu to consider climate change effects on the project area as part of meeting their management goals and objectives. The Northern Institute of Applied Climate Science (NIACS) professionals worked with the Providence team to translate the goals and climate change impacts into adaptation tactics.

The team identified five primary management goals:

- ▶ Increase and enhance canopy cover throughout the project area by involving community stakeholders in tree planting and ensuring newly planted street trees survive, ultimately improving the health environment (heat and air quality) for people who reside, work, and attend school in the project area.
- ▶ Decrease localized street flooding and stormwater runoff.
- ▶ Engage residents, community groups, and institutions within the project area in planning and implementation of tree planting and stewardship activities.
- ▶ Expand on existing program models and pilot new initiatives and practices (e.g., community youth tree watering and job-training partnerships) that will increase and improve tree canopy while also serving as tools for outreach and education regarding the urban forest. Outreach and education tools will link to climate and health, increasing awareness of the benefits trees provide to our communities. They will also support the development of the citywide PVD Tree Plan, which is a collaborative, equity-focused action plan for Providence's urban forest. The PVD Tree Plan is currently in development by PNPP in partnership with a coalition of core stakeholders.
- ▶ Protect existing tree canopy in the project area and prevent future canopy loss due to extreme weather events, such as heavy storms and high winds.

Managers from the Providence Parks Department and PNPP plan to address these goals by selecting and planting climate-adapted trees (30-50 in the project area through PNPP's Neighborhood Street Tree Planting model), identifying the most vulnerable sites and implementing tree pit projection, installing stormwater tree pits (five tree filter pits in the project area), and implementing community and stakeholder engagement strategies. Such strategies could include developing partnerships, cultivating new resident PNPP Tree Leaders, and involving residents, students, and community groups in tree stewardship.

Climate Change Impacts and Providence

The Providence team used their knowledge of the local landscape to examine regional climate change impacts and assess how these impacts will increase or decrease vulnerabilities in the project area. The team identified four key climate change impacts affecting the project area.

- ▶ Temperatures in New England are projected to increase 3.5 to 8.5 °F by the end of the 21st century, with the greatest warming expected to occur during winter.⁵³⁵ This increase could have detrimental effects on human and ecosystem health, leading to a chain reaction of changes among plants, animals, weather patterns, and more. The project area is affected by Providence's urban heat islands and battles poor air quality that amplifies breathing issues. In addition, the area has relatively low access to cooling via public cooling centers and air conditioning, which may further exacerbate heat-related health problems.
- ▶ Altered hydrology is another concern. Annual precipitation is projected to increase and there is potential for reduced growing season precipitation. Intense precipitation events will continue to become more frequent, and the timing and amount of stream flow is

expected to change. Compacted soil, fewer lawn strips, and more pavement may result in lower water capture and higher runoff containing pollutants.

- ▶ Storm events will continue to become more frequent and severe, including the increased possibility of tornados, hurricanes, and other tropical storms. Older trees in the existing canopy become susceptible to breakage due to wind as well as uprooting due to high water saturation, while localized flooding events, paired with a combined sewer system, could lead to water issues.
- ▶ Warmer temperatures, combined with varied precipitation, may alter soil moisture and increase drought risk. Changes in temperature and precipitation are projected to alter soil moisture patterns throughout the year, with the potential for both wetter and drier conditions depending on the location and season. In addition, forest vegetation may face increased risk of moisture deficit and drought during the growing season.

A comparison of the climate and health tree species list ([appendix 3](#)) with street tree inventory data from the project area indicates that none of the tree species planted are considered highly vulnerable to climate change, and most trees have low or low-moderate vulnerability. Red maple and Norway maple (*Acer platanoides*) are some of the most common species in the project area and are projected to experience a decline in habitat suitability. Although these species are beneficial for carbon storage and shade, they also have high allergenicity and volatile organic carbon emissions that could have negative human health implications. Thus, the large presence of Norway and red maple could contribute further to vulnerability of both the tree canopy and the human population in the neighborhood.

Overall, neighborhood vulnerability is higher compared to some others in Providence due to lower income, low canopy, low home ownership rates, high impervious surfaces, and urban heat island effects. In terms of adaptive capacity factors, the project area does not have a lot of connected green space or biodiversity. The low-income neighborhood and low rates of home ownership result in lower private tree care compared to other areas in the city because trees are costly to maintain. On the other hand, age class diversity of tree canopy is varied, and the City of Providence already has a pruning cycle established through the 15 wards in the city, each divided into 10 areas based on tree inventory. Ten percent of each ward is pruned each year. The Division of Forestry and PNPP jointly manage a program called Providence Community Tree Keepers that trains residents to act as stewards of the city's young trees. While the program currently has little participation within the project area, there are opportunities to build that capacity in the near future.

Challenges and Opportunities for Management

A changing climate will present challenges and opportunities for accomplishing the management objectives. Each management objective was evaluated keeping the list of climate change impacts in mind and many challenges were based on the vulnerabilities identified in the previous step.

CHALLENGES

The primary challenge for meeting the project goals is the overall difficulty and uncertainty of tree establishment in the face of increased drought risk and increased storm and wind events. It may be difficult to identify tree species that meet all criteria, such as climate adaptability, resilience, health concerns, and the conditions of bioswale pits. Planting days may be impacted with additional storms, and hot/dry weather may make it more difficult to get residents outside for tree watering and other maintenance. It may also be difficult to ensure the survival of trees in bioswales and maximize tree size and longevity in stormwater installations.

A changing climate can create more challenges for community engagement. Increased storms and wind events may result in more tree damage, which can increase hesitation and negative views toward trees. As a result, recruiting participants for activities such as tree planting and maintenance may be more difficult.

OPPORTUNITIES

Climate change may also create opportunities in the project area. When selecting tree species, there will be an opportunity to select and grow nursery stock that is more adaptable to climate change. Providence has well-established relationships with local nursery growers that can adjust their stock based on city requests. In addition, urban forest managers may be able to extend the planting season with a longer growing season that results in earlier springs and delayed autumns. A longer growing season may also aid in growth and establishment of young trees.

Burgeoning concern about climate change among residents and professionals alike may create opportunities to collaborate with different partners, pilot ideas, and work with young residents who are concerned by and interested in climate change. Providence's newly established Green Justice Zones may be used to build more capacity and community organizing around climate. Stormwater runoff, one of the primary concerns, is a "hot topic" due to climate change, which may present an additional opportunity to obtain more funding for management goals. Lastly, heat and flood impacts in the neighborhood can be used to deliver more effective messaging and make connections to urban trees.

Adapting to Climate Change

Adaptation actions were identified with the previous information in mind (Table 3). Several adaptation approaches were selected that addressed the biggest climate change impacts and challenges while also capitalizing on opportunities to meet project goals. Approaches and tactics were selected related to four main themes. The first theme focused on tree species

Table 3.—Selected proposed adaptation actions identified for the neighborhood project area in Providence, Rhode Island

Management Area/Topic	Climate Change Impacts	Adaptation Approach	Proposed Adaptation Tactics
Expanding tree canopy	<ul style="list-style-type: none"> ▶ Extreme heat, reduced soil moisture in summer 	<ul style="list-style-type: none"> ▶ Approach 7.3: Introduce species, genotypes, and cultivars that are expected to be adapted to future conditions 	<ul style="list-style-type: none"> ▶ Select drought-adapted and wind-tolerant trees by examining adaptive capacity scores for individual tree species
Stormwater/ street flooding	<ul style="list-style-type: none"> ▶ Increased heavy rain events and flooding, reduced soil moisture in summer 	<ul style="list-style-type: none"> ▶ Approach 4.1: Maintain or restore soils and nutrient cycling in urban areas ▶ Approach 5.1: Reduce impacts from extreme rainfall and enhance stormwater infiltration and storage ▶ Approach 9.1: Co-design large-scale green infrastructure and built systems to promote health ▶ Approach 9.2: Provide micro-scale nature experiences to promote health and healing 	<ul style="list-style-type: none"> ▶ Select flood-adapted trees for use in bioswales ▶ Increase biodiversity of ground cover and widen tree pits around existing trees ▶ Plant multiple trees in larger beds and change soil within the beds over time ▶ Implement more lawn strips ▶ Provide homeowners with education/guidance for planting and recommend planting a layer under their trees ▶ Look into products such as mycorrhizae packets, hydrogel, and biochar
Maintaining existing tree canopy	<ul style="list-style-type: none"> ▶ More extreme storms and wind, more severe hurricanes 	<ul style="list-style-type: none"> ▶ Approach 5.2: Reduce risk of damage from extreme storms and wind 	<ul style="list-style-type: none"> ▶ Maintain current systematic pruning program ▶ Consider planting windbreaks that help other trees survive on the property and reduce drying ▶ Conduct additional tree risk assessments to identify hazardous trees for removal in priority areas ▶ Treat trees that could become a hazard
Community/ stakeholder engagement (mixed land-use types)	<ul style="list-style-type: none"> ▶ Extreme heat and air quality effects on people, climate-related trauma, and stress 	<ul style="list-style-type: none"> ▶ Approach 1.3: Address climate and health challenges of disadvantaged communities and vulnerable populations ▶ Approach 8.1: Provide nature experiences to ease stress and support mental function 	<ul style="list-style-type: none"> ▶ Develop communication strategies using the local school and organizations (e.g., Boys & Girls Clubs), and send the children home flyers ▶ Implement additional door-to-door canvassing and events/programming around trees ▶ Identify heritage trees and bring attention to them ▶ Create small green spaces that incorporate nature and are designed by the community

selection to align with future climate conditions, making use of a more diverse palette of future-adapted trees in tree-planting initiatives. The second theme focused on increasing water infiltration and soil quality. The urban forest land managers were interested in exploring several tactics to increase herbaceous ground cover as a way of increasing soil organic matter and infiltration. The third theme is related to increasing concern about extreme storms and wind affecting the existing tree canopy, which prompted the project partners to consider doing a local tree canopy risk assessment to identify trees that may need additional care or removal. Finally, concerns about climate-change effects to the area's residents prompted the urban forest land managers to consider exploring the idea of creating a small space for engagement with nature as a source of community healing. This would require additional input from residents regarding what type of space would be wanted, where it could be located, and how to engage local artists to help design the space.

Monitoring

The project partners identified several monitoring items that could help inform future management. PNPP will monitor the number of new species planted and inventory the diversity and survival of trees over 2 years. Organic matter content and microbial biomass in the soil also can be measured in addition to bioswale monitoring that is already conducted, which evaluates the infiltration of stormwater through a partnership with researchers at the Rhode Island School of Design.

Stewardship and maintenance activities will be tracked by measuring the number of volunteers who participate in stewardship within the neighborhood and the number of trees maintained by volunteers. The number of risk assessments and emerald ash borer treatments conducted also can be documented.

Community engagement will be measured by the number of residential sites that request a tree planting, and the number of presentations, events, pamphlets, and general outreach activities can be documented as well. Depending on the green space designed as a pocket park, urban forest land managers can determine community surveying methods to measure the impact, such as the number of engagements with a particular aspect of the space.

More Information

Information on this adaptation demonstration is available at www.forestadaptation.org/pnpp.

APPENDIX 3

Climate and Health Species List for Rhode Island Urban Trees

The tree species list for Rhode Island (Table 5) was compiled to show some of the benefits and concerns when selecting trees to reduce climate change vulnerability, reduce carbon dioxide in the atmosphere, and provide benefits to human health. The list is meant to show the complexity of tree selection but should not be considered a recommended species list. Other factors not included in this list may also be important to species selection in forested areas, such as native species benefits, species natural ranges, site conditions, and goals. Urban and landscape decisions may also need to consider “right tree, right place”, site conditions, moisture availability, and root space. While this list can show some species with identified climate and health benefits, this is not the complete list of choices available for Rhode Island and may not be suitable for planting for your specific site or needs. Contact the State Stewardship Forester for species recommendations for natural areas and the State Urban Forester for recommendations for urban areas.

Climate Vulnerability

Trees can be vulnerable to a variety of climate-related stressors such as intense heat, drought, flooding, and changing pest and disease patterns. Climate vulnerability is a function of the impacts of climate change on a species and its adaptive capacity. Species with negative impacts on habitat suitability and low adaptive capacity will have high vulnerability and vice versa. The following factors were used to determine climate vulnerability:

- ▶ **Hardiness and heat zone tolerance:** Tree species ranges were recorded from government, university, and arboretum websites. Species tolerance ranges were compared to current and projected heat and hardiness zones for Rhode Island using downscaled climate models (Table 4).⁵³⁶
- ▶ **Habitat suitability modeling projections:** Modeled projections for native species were summarized from the [Climate Change Atlas](#) website under low and high emissions for the 1-degree latitude/longitude grid cell that covers Rhode Island (east of 71W and south of 41N).⁵³⁷⁻⁵³⁹
- ▶ **Adaptability:** Adaptability scores were generated for each species based on literature describing its tolerance to disturbances such as drought, flooding, pests, and disease, as well as its growth requirements such as shade tolerance, soil needs, and ease of nursery propagation. Scores were assigned to Rhode Island species using methods developed in an urban forest vulnerability assessment for Chicago.⁵⁴⁰

Table 4.—Current and projected USDA Hardiness Zones and AHS Heat Zones for the state of Rhode Island.⁵³⁶ Hardiness zone is determined by the average lowest temperature over a 30-year period. Heat zones are determined by the number of days above 86 °F.

Time Period	Hardiness Zone Range Rhode Island		Heat Zone Range Rhode Island	
	Low Emissions	High Emissions	Low Emissions	High Emissions
1980-2009	6 to 7		2 to 4	
2010-2039	6 to 7	7	3 to 5	4 to 6
2040-2069	7	7 to 8	4 to 6	6 to 8
2070-2099	7	8	4 to 6	8 to 9

Carbon

Trees provide benefits by reducing greenhouse gases in the atmosphere by directly storing carbon in their leaves, wood, and roots, and by helping to reduce energy use for heating and cooling. Benefits provided by each species were modeled for the city of Providence, RI, and binned into categories based on their relative carbon benefits to one another using methods developed for the [i-Tree Species selector](#).⁵⁴² The following factors were combined to assess carbon benefits:

- ▶ **Carbon storage:** the total of all carbon stored during the average lifespan for the species. Larger trees tend to store more carbon.
- ▶ **Carbon sequestration rate:** carbon absorption per year. Species that gain a lot of growth per year will have higher sequestration rates.
- ▶ **Carbon savings from energy use:** the total amount of carbon saved from reduced heating and cooling energy use. Large shade trees tend to reduce cooling energy use and large conifers tend to reduce heating energy use.

Human Health

Trees can reduce risks to human health that may be faced under a changing climate, such as heat stress and reduced air quality, by providing shade, cooling through transpiration, and absorption of pollutants. Benefits provided by each species were modeled for the city of Providence, RI and binned into categories based on their relative health benefits to one another using methods developed for the [i-Tree Species selector](#).⁵⁴¹ The following factors were combined to assess human health benefits:

- ▶ **Leaf area:** the maximum leaf area reached over the species' lifespan. Trees with greater leaf area provide more shade and can typically absorb more pollutants.
- ▶ **Transpiration:** average transpiration rate per year, which is influenced in part by tree size and differences in water-use efficiency. Trees that transpire more can be better at evaporative cooling and mitigating flooding.

- ▶ **Pollutants removed:** weighted sum of the pollutants NO₃, O₃, PM_{2.5} and SO₂ removed over a species' lifespan.

Some trees may need to be considered for their potential negative effects on human health. In particular, some trees produce allergenic pollen or volatile organic compounds such as isoprene or monoterpenes that can reduce air quality. Isoprene and monoterpene emissions for each species were modeled for the city of Providence, RI, and binned into categories based on their relative health benefits to one another using methods developed for the i-Tree Species selector.⁵⁴¹ Allergenicity was based on Ogren Plant Allergy Scale.⁵⁴² The following factors were combined to assess human health disservices:

- ▶ **Allergenicity:** how likely the tree is to cause allergies. Wind-pollinated trees tend to be more allergenic.
- ▶ **Isoprene emissions:** total emissions of isoprene over a species' lifespan. Certain species of broadleaved trees, such as oaks, are known for high isoprene emissions.
- ▶ **Monoterpene emissions:** total emissions of monoterpenes over a species' lifespan. Some species, and many conifers in particular, can be high emitters of monoterpenes.

Table 5.—Tree species list developed to aid Rhode Island community forestry practitioners in selecting trees to reduce climate change vulnerability, reduce carbon dioxide in the atmosphere, and provide benefits to human health. It is meant to be a complement to other tree selection resources. Other factors may also need to be considered, such as aesthetics, local site conditions, wildlife value, or nursery availability. Some species may have climate and health benefits but may not be suitable for planting for other reasons, such as having invasive potential or susceptibility to pests or pathogens.

Scientific name	Common name	Climate vulnerability	Carbon benefit	Health benefit	Health disservices	Notes
<i>Abies balsamea</i>	Balsam fir	moderate-high	moderate	moderate-high	moderate	
<i>Acer campestre</i>	Hedge maple	low	low	low	moderate	can be invasive
<i>Acer ginnala</i>	Amur maple	moderate-high	low-moderate	moderate-high	moderate	can be invasive
<i>Acer griseum</i>	Paperbark maple	moderate	low	low-moderate	moderate-high	
<i>Acer negundo</i>	Boxelder	moderate-high	moderate	moderate	moderate	can be invasive
<i>Acer rubrum</i>	Red maple	moderate	high	high	moderate-high	
<i>Acer saccharinum</i>	Silver maple	moderate	moderate	moderate-high	moderate-high	
<i>Acer saccharum</i>	Sugar maple	low-moderate	moderate-high	high	moderate-high	
<i>Acer tartaricum</i>	Tatarian maple	moderate-high	n/a	n/a	moderate	
<i>Acer truncatum</i>	Shantung maple	low-moderate	low	low	moderate-high	
<i>Acer x freemanii</i>	Freeman maple	low-moderate	n/a	n/a	moderate	
<i>Aesculus hippocastanum</i>	Horse chestnut	low-moderate	moderate-high	high	low	can be invasive

Scientific name	Common name	Climate vulnerability	Carbon benefit	Health benefit	Health disservices	Notes
<i>Aesculus x carnea</i>	Red horsechestnut	low	low	low-moderate	low	
<i>Amelanchier canadensis</i>	Shadblow/ Canadian serviceberry	moderate-high	low	low	low	
<i>Amelanchier laevis</i>	Serviceberry	low	low	low-moderate	low	
<i>Betula alleghaniensis</i>	Yellow birch	moderate	moderate	moderate-high	low-moderate	significant pest/ disease issues
<i>Betula lenta</i>	Sweet birch	moderate-high	low-moderate	moderate	low-moderate	
<i>Betula nigra</i>	River birch	low-moderate	moderate-high	moderate	low-moderate	
<i>Betula papyrifera</i>	Paper birch	moderate-high	moderate-high	moderate	low-moderate	significant pest/ disease issues
<i>Betula pendula</i>	Silver birch	moderate-high	low-moderate	low-moderate	moderate	significant pest/ disease issues, can be invasive
<i>Betula populifolia</i>	Gray birch	moderate-high	low	low	low-moderate	significant pest/ disease issues
<i>Carpinus betulus</i>	European hornbeam	low	low-moderate	low-moderate	low-moderate	
<i>Carpinus caroliniana</i>	American hornbeam	low	low	low-moderate	moderate	
<i>Carya alba</i>	Mockernut hickory	low	moderate-high	moderate-high	moderate-high	
<i>Carya glabra</i>	Pignut hickory	low-moderate	moderate-high	high	moderate-high	
<i>Carya ovata</i>	Shagbark hickory	low-moderate	moderate-high	moderate-high	moderate-high	
<i>Carya texana</i>	Black hickory	moderate	high	high	moderate-high	
<i>Catalpa bignonioides</i>	Southern catalpa	moderate	moderate-high	moderate-high	low-moderate	
<i>Celtis laevigata</i>	Sugarberry	low-moderate	low-moderate	high	low-moderate	
<i>Celtis occidentalis</i>	Hackberry	moderate	moderate	high	low-moderate	
<i>Cercidiphyllum japonicum</i>	Katsura tree	moderate-high	low	low-moderate	moderate-high	
<i>Cercis canadensis</i>	Eastern redbud	low-moderate	low	low	low	
<i>Chamaecyparis thyoides</i>	Atlantic white-cedar	moderate	low-moderate	moderate-high	low-moderate	
<i>Cladrastis kentukea</i>	Yellowwood	low	low	low-moderate	low	
<i>Cornus florida</i>	Flowering dogwood	low-moderate	low	low	low-moderate	significant pest/ disease issues
<i>Cornus kousa</i>	Kousa dogwood	low	low	low	low-moderate	
<i>Corylus colurna</i>	Turkish filbert	moderate	low-moderate	low-moderate	low-moderate	

Scientific name	Common name	Climate vulnerability	Carbon benefit	Health benefit	Health disservices	Notes
<i>Crataegus crus-galli var. inermis</i>	Thornless cockspur hawthorn	moderate	low	low-moderate	low	significant pest/disease issues
<i>Diospyros virginiana</i>	Common persimmon	low	low	low	low	
<i>Eucommia ulmoides</i>	Hardy rubber tree	moderate	low-moderate	low-moderate	moderate	
<i>Fagus grandifolia</i>	American beech	low-moderate	moderate-high	high	low-moderate	Beech leaf disease (BLD) present
<i>Fagus sylvatica</i>	European beech	moderate	moderate	high	low-moderate	Beech leaf disease (BLD) present
<i>Ginkgo biloba</i>	Ginkgo	low	low-moderate	moderate-high	moderate-high	
<i>Gleditsia triacanthos var. inermis</i>	Honeylocust	low-moderate	moderate-high	low-moderate	low	
<i>Gymnocladus dioica</i>	Kentucky coffeetree	low	low	low	low	
<i>Ilex opaca</i>	American holly	low	low	moderate	low	
<i>Juglans nigra</i>	Black walnut	low-moderate	high	high	moderate-high	
<i>Juniperus virginiana</i>	Eastern redcedar	low	low	low-moderate	moderate	
<i>Koelreuteria paniculata</i>	Golden raintree	low	low	low-moderate	low	
<i>Larix decidua</i>	European larch	moderate-high	moderate	moderate-high	moderate	
<i>Liquidambar styraciflua</i>	Sweetgum	low-moderate	high	high	high	
<i>Liriodendron tulipifera</i>	Tulip tree	moderate	high	high	low-moderate	
<i>Maackia amurensis</i>	Amur maackia	moderate	low	low	moderate	
<i>Metasequoia glyptostroboides</i>	Dawn redwood	low-moderate	moderate	moderate-high	low-moderate	
<i>Nyssa sylvatica</i>	Black tupelo	low	high	high	moderate	
<i>Ostrya virginiana</i>	Hop-hornbeam	low	low	low-moderate	low-moderate	
<i>Oxydendrum arboreum</i>	Sourwood	low	low	low	low-moderate	
<i>Parrotia persica</i>	Persian iron-wood	low	low	low	high	
<i>Pinus resinosa</i>	Red pine	high	moderate-high	moderate	low	significant pest/disease issues
<i>Pinus rigida</i>	Pitch pine	moderate-high	high	moderate-high	moderate	significant pest/disease issues
<i>Pinus strobus</i>	White pine	high	high	high	moderate	

Scientific Name	Common name	Climate vulnerability	Carbon benefit	Health benefit	Health disservices	Notes
<i>Pinus sylvestris</i>	Scots pine	moderate-high	moderate-high	moderate-high	moderate	significant pest/disease issues
<i>Pinus taeda</i>	Loblolly pine	low-moderate	moderate-high	moderate-high	moderate	significant pest/disease issues
<i>Pinus virginiana</i>	Virginia pine	moderate	moderate	moderate	moderate	significant pest/disease issues
<i>Platanus x acerifolia</i>	London planetree	low-moderate	moderate-high	high	low-moderate	
<i>Platanus occidentalis</i>	American sycamore	low-moderate	high	high	moderate	
<i>Populus deltoides</i>	Eastern cottonwood	moderate	moderate	moderate	moderate	prone to structural failure
<i>Populus grandidentata</i>	Bigtooth aspen	high	low-moderate	low-moderate	moderate	
<i>Populus tremuloides</i>	Quaking aspen	high	low-moderate	low	moderate	
<i>Prunus cerasifera</i>	Flowering plum	low-moderate	low	low	low	significant pest/disease issues, prone to structural failure
<i>Prunus padus</i>	Bird cherry	low-moderate	n/a	n/a	low	
<i>Prunus pensylvanica</i>	Pin cherry	high			low	
<i>Prunus sargentii</i>	Sargent cherry	moderate	low	low-moderate	low	
<i>Prunus serotina</i>	Black cherry	moderate	moderate-high	moderate	low	
<i>Prunus serrulata 'Kwanzan'</i>	Kwanzan cherry	low-moderate	low	low	low	
<i>Prunus virginiana 'Schubert'</i>	Schubert cherry	moderate-high	low	low	low	
<i>Quercus alba</i>	White oak	low-moderate	moderate-high	moderate-high	moderate-high	
<i>Quercus bicolor</i>	Swamp white oak	low-moderate	high	high	moderate-high	
<i>Quercus coccinea</i>	Scarlet oak	low-moderate	high	high	high	
<i>Quercus falcata</i>	Southern red oak	low-moderate	high	high	moderate-high	
<i>Quercus imbricaria</i>	Shingle oak	low	low	moderate-high	moderate-high	
<i>Quercus macrocarpa</i>	Bur oak	low	low-moderate	moderate-high	moderate-high	
<i>Quercus marilandica</i>	Blackjack oak	moderate	low-moderate	low-moderate	moderate-high	
<i>Quercus michauxii</i>	Swamp chestnut oak	low-moderate	moderate	moderate	moderate-high	
<i>Quercus nigra</i>	Water oak	low-moderate	moderate	moderate-high	moderate-high	
<i>Quercus palustris</i>	Pin oak	moderate	moderate-high	high	moderate-high	

Scientific Name	Common name	Climate vulnerability	Carbon benefit	Health benefit	Health disservices	Notes
<i>Quercus phellos</i>	Willow oak	low	high	high	moderate-high	
<i>Quercus prinus</i>	Chestnut oak	low	moderate	moderate	moderate-high	
<i>Quercus robur</i>	English oak	low-moderate	moderate-high	high	high	can be invasive
<i>Quercus rubra</i>	Northern red oak	low-moderate	high	moderate-high	high	
<i>Quercus stellata</i>	Post oak	moderate	moderate	moderate	moderate-high	
<i>Quercus velutina</i>	Black oak	low-moderate	moderate	low-moderate	moderate-high	
<i>Sassafras albidum</i>	Sassafras	low-moderate	moderate	moderate	low	
<i>Sorbus alnifolia</i>	Korean mountain ash	moderate	low-moderate	low-moderate	low	
<i>Styphnolobium japonicum</i>	Sophora/ Japanese pagoda	moderate	low	low-moderate	low	
<i>Styrax japonicus</i>	Japanese snowbell	low-moderate	low	low	low	
<i>Syringa reticulata</i>	Japanese tree lilac	moderate	n/a	n/a	low	
<i>Taxodium distichum</i>	Bald cypress	low	high	high	moderate	
<i>Tilia americana</i>	American basswood	low-moderate	low-moderate	moderate	low-moderate	
<i>Tilia cordata</i>	Littleleaf linden	moderate	low-moderate	moderate	low-moderate	
<i>Tilia tomentosa</i>	Silver linden	moderate	low-moderate	moderate-high	low-moderate	
<i>Tilia x euchlora</i>	Caucasian linden	low-moderate	low-moderate	moderate	low-moderate	
<i>Ulmus alata</i>	Winged elm	low-moderate	low-moderate	low	low-moderate	
<i>Ulmus americana</i>	American elm	low-moderate	high	high	low-moderate	significant pest/ disease issues
<i>Ulmus 'Homestead'</i>	Homestead elm	low	n/a	moderate-high	low	can be invasive
<i>Ulmus parvifolia</i>	Chinese elm	low	moderate-high	moderate-high	low-moderate	can be invasive
<i>Zelkova serrata</i>	Japanese zelkova	low	low	low-moderate	low	

Janowiak, Maria K.; Brandt, Leslie A.; Wolf, Kathleen L.; Brady, Mattison; Darling, Lindsay; Lewis, Abigail Derby; Fahey, Robert T.; Giesting, Kristen; Hall, Eboni; Henry, Molly; Hughes, Maise; Miesbauer, Jason W.; Marcinkowski, Kailey; Ontl, Todd; Rutledge, Annamarie; Scott, Lydia; Swanston, Christopher W. 2021.

Climate adaptation actions for urban forests and human health. Gen. Tech. Rep. NRS-203. Madison, WI: U.S. Department of Agriculture, Forest Service, Northern Research Station. 115 p. <https://doi.org/10.2737/NRS-GTR-203>.

Urban areas can be particularly vulnerable to climate change due to extensive impervious cover, increased pollution, greater human population densities, and a concentration of built structures that intensify impacts from urban heat, drought, and extreme weather. Urban residents are at risk from a variety of climate stressors, which can cause both physical and mental harm. Urban forests and tree cover provide a critical role in helping cities address climate change by supporting greenhouse gas mitigation, reducing the impacts of extreme heat and altered climate that impair human health, and helping communities to adaptively respond through engagement with nature. At the same time, urban forests are vulnerable to changes in climate and in need of robust strategies to adapt to those changes.

As climate change impacts increase, efforts to “green” cities and adapt urban forests to changing conditions take on greater importance to support human health and well-being. Urban forest managers and allied professionals are looking for information to reduce climate risks to urban forests and secure their benefits for people and ecosystems. This report, *Climate Adaptation Actions for Urban Forests and Human Health*, synthesizes adaptation actions to address climate change in urban forest management and promote human health and well-being through nature-based solutions. It compiles and organizes information from a wide range of peer-reviewed research and evidence-based reports on climate change adaptation, urban forest management, carbon sequestration and storage, and human health response to urban nature.

This report includes the Urban Forest Climate and Health Adaptation Menu, which presents information and ideas for optimizing the climate and human health outcomes of urban forestry projects and provides professionals who are working at the intersection of climate, public health, and urban forestry with resources to support climate adaptation planning and activities. Notably, it *does not* provide specific recommendations or guidance for any particular place; rather, it offers a range of action opportunities at different scales that can be incorporated into either comprehensive or specific climate adaptation initiatives. The Menu can be used with an existing, tested adaptation process to help managers consider climate risks and explore the benefits and drawbacks of potential adaptation actions within the context of a particular situation or project. It also can be useful for generating productive discussions about community needs and values to guide planning, education and outreach, research, or changes in policy or infrastructure within communities.

Keywords: adaptive management, ecosystem services, green infrastructure, human health, urban forests

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident.

Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotope, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English.

To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at http://www.ascr.usda.gov/complaint_filing_cust.html and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov.



Northern Research Station

<https://www.nrs.fs.fed.us>